

## **Chapter 2**

**Protein and mRNA levels of IgM H- and L-chains artificially accumulated in HDAC2-deficient DT40 mutants are gradually reduced via a lot of generations during continuous cultivation**

by Masami Nakayama and Tatsuo Nakayama

## **SUMMARY**

We analyzed HDAC2-deficient DT40 mutant cells HDAC2(-/-), which were cultivated for varying periods, by 2D-PAGE, Western blotting, RT-PCR and immuno-electron microscopy. During cultivation, protein and mRNA levels of IgM H- and L-chains were dramatically and considerably increased at the early stage, and thereafter until the late stage gradually reduced to comparable levels in DT40 cells. Further, the mRNA level of PCAF was dramatically increased from the early to late stages, and those of HDAC7 and HDAC9 were certainly changed. The mRNA level of EBF1 was almost completely decreased at the early stage and thereafter remained unchanged as undetectable level, but that of E2A was slightly increased at the early stage and thereafter remained unchanged. Remarkably, the mRNA level of Pax5, which should down-regulate gene expressions of IgM H- and L-chains, was surely reduced at the early stage and thereafter certainly increased via the middle to late stages in anti-parallel with those of these two immunoglobulin genes. In addition, bulk acetylation levels of K9, K14, K18, K23 and K27 of histone H3 were gradually increased. Subsequently, we cloned ~4.9 kb 5'-upstream region of the Pax5 gene from DT40 genomic DNA. Chromatin immuno-precipitation assay suggested that acetylation levels of Lys-9 residues of histone H3 (K9/H3) within some chromatin segments surrounding proximal ~2.0 kb 5'-upstream region of the Pax5 gene were decreased at the early stage in HDAC2(-/-) and thereafter at the late stage increased to comparable levels in DT40 cells.

These results revealed not only that the HDAC2-deficiency induces dramatic accumulations in mRNA and protein levels of IgM H- and L-chains, and these accumulated mRNAs and proteins are gradually reduced during cultivation, but also that alterations in gene expressions of IgM H- and L-chains may be coupled with changes in those of Pax5, Aiolos, EBF1, PCAF, HDAC7 and HDAC9. In addition, changing patterns in acetylation levels of K9/H3 within proximal 5'-upstream chromatin region of the Pax5 gene in HDAC2(-/-) should qualitatively agree with those in its transcription level.

## **INTRODUCTION**

In eukaryotic cells, alterations in chromatin structure are preferentially involved in controls of gene expressions, and replication, repair and recombination of DNA and so on [1-3]. Naturally, such alterations are concerned in regulations of lymphocyte development and differentiation [4-7]. Regulations of the lymphocyte development require a number of transcription factors, including Ikaros, E2A, PU.1, GATA-3, Pax5, EBF and so on [8-13]. Moreover, transcription regulation of IgM H-chain requires USF, TFEB, Ig/EBP, NF-IL6, OCA-b and others as promoter binding proteins, and Ig/EBP, NF-IL6, YY-1, E2A, PU.1 and others as intron enhancer binding proteins. On the other hand, of various epigenetic modifications of chromatin structure with acetyl, methyl and phosphate groups and/or so on, acetylation and deacetylation of particular Lys residues (K) of core histones (H2A, H2B, H3 and H4) may

be major ones [14-21]. They are cooperatively controlled in combination of several kinds of histone acetyltransferase(s) (HATs) and deacetylase(s) (HDACs) [22-25]. To clarify individual roles of members of HATs and HDACs in expressions of cell functions, we have systematically generated a number of homozygous (or conditional) chicken DT40 mutant cell lines lacking appropriate genes [26-42] by gene targeting techniques [43-53]. Our previous analysis of HDAC2-deficient DT40 mutant cells, HDAC2(-/-), revealed that HDAC2 controls the amount of IgM H-chain at the steps of transcription of its gene plus alternative processing of its pre-mRNA [26] and down-regulates IgM L-chain gene promoter activity [29]. Moreover, the HDAC2-deficiency decreases gene expressions of HDAC7, Pax5, Aiolos, Ikaros plus EBF1, increases those of HDAC4, HDAC9, PCAF plus E2A, and changes bulk acetylation levels of several particular Lys residues of core histones H2A, H2B, H3 and H4 [33].

Subsequently, to know individual roles of these altered transcription factors in regulations of IgM H- and L-chain gene expressions, we generated homozygous DT40 mutant cells; EBF1(-/-), Aiolos(-/-), E2A(-/-) and Helios(-/-), devoid of two alleles of EBF1, Aiolos, E2A and Helios genes, respectively [33, 54-58]. In addition, we generated Pax5-deficient mutant cells, Pax5(-), devoid of the Pax5 gene located on Z chromosome that is monosomy in chickens (USCS Genome Browser data base) [see Chap. 3], but could generate only Ikaros-down mutant cells, Ikaros(-/+), devoid of two alleles of the gene located on chromosome 2 that is trisomy, since it should probably be very important or essential for DT40 cells (data will be shown elsewhere). Analyses of these resultant mutants revealed that Pax5, EBF1, Aiolos plus Ikaros down-regulate IgM H- and L-chain gene expressions, and E2A up-regulates transcriptions of these two immunoglobulin genes [33]. Thus, these results indicated that HDAC2 indirectly controls transcriptions of IgM H- and L-chain genes through opposite transcription regulations of Pax5, EBF1, Aiolos plus Ikaros, and E2A genes [33, 36].

In this study, we analyzed HDAC2(-/-) at different cultivation periods by 2D-PAGE, Western blotting, immuno-electron microscopy and RT-PCR. During cultivation of HDAC2(-/-), protein and mRNA levels of IgM H- and L-chains were dramatically and considerably accumulated at the early stage, and thereafter gradually reduced and finally at the late stage reached to comparable levels in DT40 cells. In addition, mRNA levels of HDAC7, HDAC9 and PCAF were gradually elevated, and those of Pax5, Aiolos, E2A and EBF1 were certainly altered in different patterns. Further, we cloned ~4.9 kb 5'-upstream region of the Pax5 gene from DT40 genomic DNA by our original gene walking techniques. Chromatin immuno-precipitation (ChIP) assay suggested that acetylation levels of Lys-9 residues of histone H3 (K9/H3) within the proximal ~2.0 kb 5'-upstream chromatin region of the Pax5 gene were slightly decreased at the early stage in HDAC2(-/-) and at the late stage increased to comparable levels in DT40 cells. We discussed on these changing patterns in gene expressions of IgM H- and L-chains, particular transcription factors and chromatin modifying enzymes, and acetylation levels of K9/H3 within the proximal ~2.0 kb 5'-upstream chromatin region of the Pax5 gene in HDAC2(-/-) mutants during

continuous cultivation.

## RESULTS

### **Artificially accumulated IgM H- and L-chains in HDAC2(-/-) are gradually reduced during continuous cultivation**

To know changing patterns in cellular protein levels of HDAC2(-/-) mutants during cultivation, we first analyzed total cellular proteins prepared from clone cl.33-12 of HDAC2(-/-) mutants at the early (E; ~10 days), middle (M; ~30 days) and late (L; ~60 days) stages of cultivation and DT40 cells (W) as control by 2D-PAGE (Fig. 2-1). As reported previously [26], the amounts of IgM H-chain and L-chain (detected as two spots) were dramatically and certainly increased at the early cultivation stage in mutant clone cl.33-12 compared with those in DT40 cells. Surprisingly, the elevated amounts of these two immunoglobulin proteins in clone cl.33-12 were gradually reduced via the middle to late cultivation stages to almost same levels in DT40 cells, whereas changes in cellular levels of most of other major proteins were insignificant during cultivation.

Moreover, Western blotting carried out at shorter intervals up to 56 days, using antibody for chicken IgM L-chain that cross-reacts with IgM H-chain, revealed that protein levels of IgM H-chain and L-chain (detected as two bands) were dramatically increased at the early stage in clone cl.33-28 (Fig. 2-2). Thereafter, these increased protein levels were gradually reduced via the middle to late stages to almost similar levels in DT40 cells. These results obtained in clones cl.33-12 and cl.33-28 by 2D-PAGE and Western blotting were sufficiently confirmed in two other independent clones (cl.33-30 and cl.45-28) (Suppl. Fig. 2-S1).

Next, we examined both subcellular localization and changing pattern of heightened intracellular IgM H-chains in clone cl.33-28 during cultivation by immunocytochemistry analysis with anti-chicken IgM H-chain antibody. Electron microscopy showed that the electron-dense materials were accumulated in irregularly elaborated and vesiculated rough endoplasmic reticula only at the early stage in HDAC2(-/-) mutants (data not shown). Moreover, immuno-electron microscopy revealed that colloidal gold immuno-labeling for chicken IgM H-chain was vividly observed in the electron-dense materials accumulating in the dilated and vesiculated rough endoplasmic reticula only at the early (~10 days) stage, and thereafter the heightened protein levels were obviously reduced at the late (~60 days) stage to almost similar levels in DT40 cells (Fig. 2-3).

Taken together, these results revealed not only that IgM H- and L-chains are dramatically and considerably accumulated at the early stage of cultivation in HDAC2(-/-) mutant cells, but also that these accumulated immunoglobulin proteins are gradually reduced during cultivation and finally at the late stage reached to comparable levels in DT40 cells. In addition, majority of artificially accumulated IgM

H-chains, which were indicated to exist as a native soluble form to be capable of building a high molecular weight complex with IgM L-chains in HDAC2(-/-) mutants [33], should be localized in specific subcellular organelle, i.e., rough endoplasmic reticula.

**Alterations in gene expressions of IgM H-chain, IgM L-chain and core histones in HDAC2(-/-), respectively, are significant, moderate and insignificant during continuous cultivation**

To know alterations in gene expressions of IgM H- and L-chains and core histones in HDAC2(-/-) mutants during cultivation, we prepared total RNAs from three independent clones (cl.33-28, cl.33-30 and cl.45-28) at the early (~20 days), middle (~40 days) and late (~60 days) stages of cultivation and DT40 cells. First, we carried out RT-PCR on total RNAs, using primers IgM Hc, IgM Hs plus IgM Hm and IgM L specific for whole, secreted plus membrane-bound forms of IgM H-chain mRNA and IgM L-chain mRNA, respectively, and  $\beta$ -actin was used as control (Fig. 2-4). The levels of whole and secreted forms of IgM H-chain mRNA, respectively, were obviously and dramatically elevated at the early stage and thereafter gradually reduced via the middle to late stages in these mutant clones, whereas alterations in the levels of membrane-bound form of IgM H-chain mRNA and IgM L-chain mRNA were insignificant and moderate, respectively. These findings in three mutant clones sufficiently agreed with those in clone cl.33-12 (data not shown). Next, we performed RT-PCR using appropriate primers specific for core histones H2A, H2B, H3 and H4 on total RNAs. In three mutant clones alterations in the mRNA levels of these four core histones were insignificant from the early to late stages (Fig. 2-4).

**Gene expressions of HDAC7, HDAC9 and PCAF in HDAC2(-/-) change during continuous cultivation**

To clarify whether alterations in gene expressions of chromatin modifying enzymes (HDACs and HATs) are linked with those of IgM H- and L-chains during cultivation, we carried out RT-PCR on total RNAs, using appropriate primers specific for HDAC1, HDAC3, HDAC4, HDAC7, HDAC8, HDAC9 (equivalent to HDAC5 in previous paper [33]), PCAF, GCN5, HAT1, ELP3, TIP60, MORF, MOZ and p300, together with  $\beta$ -actin as control (Fig. 2-5A). In three clones cl.33-28, cl.33-30 and cl.45-28, the HDAC9 mRNA level was certainly increased from the early (~20 days) to middle (~40 days) stages of cultivation and thereafter slightly reduced until the late (~60 days) stage. The HDAC7 mRNA level was slightly decreased at the early stage and remained unchanged until the late stage. However, gene expressions of residual HDACs did not change. On the other hand, in these mutant clones, the PCAF mRNA level was dramatically increased from the early to late stages of cultivation, whereas those of residual HAT members remained unchanged at any cultivation stages.

### **Gene expressions of E2A, EBF1, Pax5 and Aiolos in HDAC2(-/-) change differently during continuous cultivation**

To clarify whether alterations in gene expressions of various transcription factors are associated with those of IgM H- and L-chains during cultivation, we carried out RT-PCR on total RNAs, using appropriate primers specific for Oct1, Oct2, OBF1, NF- $\kappa$ B, RelB, YY1, NF-AT, PU.1, E2A, EBF1, Pax5 and Aiolos, together with  $\beta$ -actin as control (Fig. 2-5B). In three clones cl.33-28, cl.33-30 and cl.45-28, the gene expression of E2A was slightly increased at the early (~20 days) stage of cultivation and remained unchanged via the middle (~40 days) until late (~60 days) stages. The gene expression of EBF1 was almost completely decreased at the early stage and remained unchanged as undetectable level until the late stage. Interestingly, the gene expression of Pax5 was obviously reduced at the early stage but thereafter certainly elevated via the middle to late stages. The gene expression of Aiolos was observed to be low at the early stage, and further decreased from the middle to late stages. On the other hand, the gene expressions of residual factors (Oct1, Oct2, OBF1, NF- $\kappa$ B, RelB, YY1, NF-AT and PU.1) remained unchanged as a whole during cultivation.

### **Bulk acetylation levels of some particular Lys residues of core histones in HDAC2(-/-) change during continuous cultivation**

To know whether bulk acetylation levels of particular Lys residues (K) of core histones H2A, H2B, H3 and H4 are altered associated with those of gene expressions of IgM H- and L-chains during cultivation, we carried out immuno-blotting on total cellular proteins prepared from three clones (cl.33-28, cl.33-30 and cl.45-28) at the early (~20 days), middle (~40 days) and late (~60 days) stages of cultivation, together with those from DT40 cells as control. Site-specific antibodies for various acetylated Lys residues of core histones H2A, H2B, H3 and H4 were used (Fig. 2-6). Concerning histone H2A, bulk acetylation levels of K5, K7 and K9 remained unchanged (or insignificantly changed) during cultivation as a whole. Among Lys residues of histone H2B tested, bulk acetylation level of K16 was certainly reduced at the early stage and thereafter remained unchanged until the late stage, whereas those of remaining Lys residues unchanged at any cultivation stages. Bulk acetylation level of K5 of histone H4 was elevated at the early stage and remained unchanged until the late stage, but those of remaining Lys residues unchanged as a whole. Interestingly, concerning histone H3, bulk acetylation levels of K9, K14, K18, K23 and K27 were obviously elevated from the early via middle to late stages, whereas the methylation level of K9 did not change. These results obtained qualitatively here were uncertain yet because some antibodies used were relatively low specificity.

### **Cloning of ~4.9 kb 5'-upstream region of chicken Pax5 gene**

As will be mentioned in our next article [Chap. 3], in spite of insufficient information concerning the

Pax5 gene (as a single allele) located on Z chromosome that is monosomy in chickens, fortunately, we could generate the Pax5-deficient DT40 mutant cell line, Pax5(-), and clarify severe and moderate influences of its deficiency on the gene expressions of IgM H- and L-chains and others. In order to know transcription manner of the chicken Pax5 gene, sufficient information on its proximal 5'-upstream region should be essential. However, as mentioned above, nucleotide sequences of the gene were not fully deposited in any database, whereas those on its homologues could be cited from GenBank. Therefore, we cloned the ~4.9 kb 5'-upstream region of the Pax5 gene from DT40 genomic DNA by our original gene walking techniques, including Southern blotting, colony hybridization and sub-cloning as follows.

To clone the ~4.9 kb 5'-upstream fragment of the Pax5 gene, we first amplified the 85 bp fragment, corresponding to positions -55 to +30 in exon 1 and including its translation initiation codon, from DT40 genomic DNA by PCR. Since several screenings using the resultant PCR product as probe yielded no positive clones from a DT40  $\lambda$ FIXII genomic library [26], we directly screened DT40 genomic DNA to obtain the Pax5 5'-upstream fragment in combination with Southern blotting and colony hybridization using the above-mentioned PCR amplified product. As a first step, we digested DT40 genomic DNA in various different combinations of two of numerous restriction enzymes, each of which recognizes multiple cloning sites of pBluescript II SK(+) plasmid, and subsequently Southern blotting using the 85 bp PCR amplified fragment as probe was performed on digested DNA fragments several times. Of numerous positive fragments being different in length and restriction enzyme sites, we focused our attention on ~2.1 kb XbaI/ClaI fragment containing the 85 bp fragment. To obtain the desired ~2.1 kb fragment, large amounts of genomic DNA (200  $\mu$ g) were digested with XbaI plus ClaI and separated in 0.8% agarose gel, followed by excising the gel piece expected to contain ~2.1 kb fragments. The DNA fragments were extracted using filter cartridge from the gel piece, recovered by phenol/chloroform extraction plus ethanol precipitation and inserted into XbaI/ClaI sites of pBluescript II SK(+) vector. The ligated plasmids were introduced into XL-1 blue MRF competent cells, and colony hybridization using the 85 bp fragment as probe was carried out to yield positive colonies. Then, recombinant plasmid DNA was isolated and its nucleotide sequences were determined. The XbaI/ClaI fragment consisted of 1995 bp 5'-upstream and 241 bp downstream regions from the transcription initiation site +1.

As a second step, we obtained further extended 5'-upstream fragment of the Pax5 gene, using colony hybridization and sub-cloning techniques as follows. By PCR using appropriate sense and antisense primers constructed based on the above-mentioned nucleotide sequences, the 1626 bp fragment corresponding to positions -1656 to -32 of the gene was amplified from DT40 genomic DNA. The resultant PCR products were digested by BssHII and separated in 0.8% agarose gel. The desired 263 bp fragment was excised from the gel, purified and sequenced. The obtained BssHII fragment corresponded to the region from positions -1656 to -1394 of the Pax5 gene was used as probe in

subsequent sub-cloning.

To obtain the ~4.9 kb 5'-upstream region of the Pax5 gene, ~4.9 kb BamHI/ClaI fragment hybridized with both of the 85 bp and 263 bp fragments as probes was inserted into pBluescript II SK(+) vector in frame. Screening and cloning techniques were essentially similar to those described above. Finally, we cloned and sequenced 4950 bp 5'-upstream fragment and 241 bp open reading frame of the chicken Pax5 gene (Suppl. Fig. 2-S2). As a matter of course, nucleotide sequences of XbaI/ClaI fragment agreed fully with those of BamHI/ClaI fragment at positions -1995 to +241. The nucleotide sequences of the ~4.9 kb 5'-upstream region of the chicken Pax5 gene will be registered to a database. Our nucleotide sequences from positions -635 to +46 were surely coincident with those very recently reported [12]. Dual-luciferase assay with several 3'-deletion and 5'-deletion mutant vectors in HeLa and DT40 cells preliminarily suggested not only that as a whole the ~4.9 kb 5'-upstream region affected negatively but very slightly on the Pax5 gene expression, but also that two 5'-upstream regions from positions -1561 to -974 and positions -521 to -40 were possibly involved in negative and bare regulation of the gene expression, though these findings are still uncertain (our unpublished data).

#### **Acetylation levels of Lys-9 residues of histone H3 within proximal ~2.0 kb 5'-upstream chromatin region of chicken Pax5 gene in HDAC2(-/-) probably change during continuous cultivation**

In general, chromatin structure surrounding the proximal 5'-upstream region of a certain gene is thought to participate directly/closely in its transcription, regardless of the existence or nonexistence of transcriptional element(s) within the region. The above-mentioned unpublished data suggested that at least ~1.6 kb 5'-upstream region of the Pax5 gene was necessary for its transcription, though those were uncertain. Furthermore, remarkably, as shown in Figure 2-5B, in HDAC2(-/-) mutants the gene expression of Pax5 was decreased at the early cultivation stage and thereafter increased until the late stage. Therefore, to know whether chromatin structure changes during cultivation, we carried out chromatin immuno-precipitation (ChIP) assay, using site-specific antibody for acetylated Lys-9 residue of histone H3 (K9/H3) and appropriate primers (Suppl. Table 2-SII), on the proximal ~2.0 kb 5'-upstream chromatin region of the Pax5 gene in three clones (cl.33-28, cl.33-30 and cl.45-28) at the early (~20 days) and late (~60 days) cultivation stages and DT40 cells as control (Fig. 2-7). Acetylation levels of K9/H3 within chromatin surrounding the ~2.0 kb 5'-upstream and open reading frame regions (from positions -1923 to +30) of the Pax5 gene were high in DT40 cells. Noticeably, in spite of the HDAC2-deficiency, as a whole, acetylation levels of K9/H3 within the ~2.0 kb 5'-upstream chromatin region of the gene in these mutant clones were observed to be slightly decreased at the early stage and thereafter increased at the late stage, whereas those within the open reading frame remained unchanged. In addition, in clone cl.33-28, acetylation levels of K14/H3 and K18/H3 within the ~2.0 kb 5'-upstream chromatin region of the gene were observed to change slightly at the early and late stages (our unpublished data).

These findings suggested that acetylation levels of at least K9/H3 within the proximal ~2.0 kb 5'-upstream chromatin region of the Pax5 gene in HDAC2(-/-) mutants were reduced at the early cultivation stage and thereafter elevated at the late stage to almost the same levels in DT40 cells. Thus, these results qualitatively agreed with those on alterations in the transcription level of the Pax5 gene.

## DISCUSSION

To know individual roles of transcription factors altered associated with the HDAC2-deficiency on regulation of gene expressions of IgM H- and L-chains, we generated homozygous DT40 mutant cell lines EBF1(-/-), Aiolos(-/-), E2A(-/-) plus Helios(-/-) [33, 36, 54-58], and Pax5-deficient mutant cell line Pax5(-) [Chap. 3] plus Ikaros-down mutant cell line Ikaros(-/+)(its generation will be shown elsewhere). Systematic analyses of these resultant mutants showed that Pax5, EBF1, Aiolos plus Ikaros down-regulate transcriptions of IgM H- and L-chain genes, and E2A up-regulates transcriptions of these two immunoglobulin genes, indicating that HDAC2 regulates indirectly transcriptions of IgM H- and L-chain genes through opposite regulations of gene expressions of Pax5, EBF1, Aiolos plus Ikaros, and E2A [33, 36].

In this study, through qualitative analyses of HDAC2(-/-) mutant cells, which were collected at different cultivation periods, we casually noticed following surprising phenomena. 2D-PAGE showed not only that amounts of IgM H- and L-chains were dramatically and certainly elevated at the early stage of cultivation, but also that the elevated amounts of these two immunoglobulin proteins were gradually decreased and at the late stage reached to comparable levels in DT40 cells (Fig. 2-1). On the other hand, changes in cellular levels of most of other major proteins were insignificant during cultivation. Western blotting carried out at shorter intervals, using antibody for chicken IgM L-chain that cross-reacts with IgM H-chain, sufficiently confirmed the above-mentioned results obtained by 2D-PAGE (Fig. 2-2 and Suppl. Fig. 2-S1). Immuno-electron microscopy using antibody specific for chicken IgM H-chain also showed that in HDAC2(-/-) the immunoglobulin proteins were clearly accumulated in endoplasmic reticula at the early stage, and the accumulated proteins were reduced at the late stage to almost similar level in DT40 cells (Fig. 2-3). Further, RT-PCR using primers IgM Hc and Hs showed that whole and secreted forms of IgM H-chain mRNA were considerably and dramatically increased at the early stage, and the elevated amounts of these two type mRNAs were gradually reduced via the middle to late stages to very close levels in DT40 cells (Fig. 2-4).

Interestingly, RT-PCR using appropriate primers specific for various genes of chromatin modifying enzymes (HDACs and HATs) and transcription factors showed noticeable results as follows (Fig. 2-5). Concerning HATs, in HDAC2(-/-) the gene expression of PCAF was dramatically elevated from the early via middle to late stages of cultivation. On the other hand, regarding HDACs, the gene expression of

HDAC9 was surely increased from the early to middle stages but thereafter slightly decreased at the late stage, contrary to this, that of HDAC7 was slightly decreased at the early stage and remained unchanged during further cultivation. As for transcription factors, in HDAC2(-/-) the gene expression of EBF1 was almost completely decreased at the early stage and thereafter remained undetectable level until the late stage, but that of E2A was slightly elevated at the early stage and remained unchanged during further cultivation. Surprisingly, the gene expression of Pax5 was obviously decreased at the early stage and thereafter certainly increased via the middle to late stages. That is, as a whole, in HDAC2(-/-) the changing pattern of the Pax5 gene expression during cultivation was in anti-parallel with those of IgM H- and L-chain gene expressions. Moreover, immuno-blotting using site-specific antibodies for various acetylated Lys residues of core histones H2A, H2B, H3 and H4 showed that bulk acetylation levels of K9, K14, K18, K23 and K27 residues of histone H3 were gradually increased during cultivation, although insignificant alterations were observed in those of remaining Lys residues (except K16 of histone H2B and probably K5 of histone H4) of core histones tested (Fig. 2-6).

To explore the manner for the transcription of the chicken Pax5 gene, first, we directly cloned and sequenced its ~4.9 kb 5'-upstream fragment from DT40 genomic DNA by our original gene walking techniques including Southern blotting, colony hybridization and sub-cloning (Suppl. Fig. 2-S2), since detailed information on its nucleotide sequences was not deposited in any database at that time. Dual-luciferase assay showed that the ~4.9 kb 5'-upstream region and especially its two regions of positions -1561 to -974 and positions -521 to -40 are thought to be possibly necessary for negative control of the Pax5 gene expression (our unpublished data), although critical promoter region(s) or element(s) has not been determined yet. Finally, to know how the transcription level of the Pax5 gene is altered in HDAC2(-/-) during cultivation, we analyzed chromatin structure surrounding the proximal 5'-upstream region from positions -1923 to +30 of the gene at the early and late stages by the ChIP assay. Interestingly, acetylation levels of Lys-9 residues of histone H3 (K9/H3) within the limited chromatin region were decreased at the early stage and thereafter increased at the late stage to almost similar levels in DT40 cells (Fig. 2-7).

Taken together, these results indicated that the HDAC2-deficiency dramatically and considerably induces accumulations of IgM H- and L-chains based on dramatic and considerable increases in their gene expressions (mRNAs) at the early stage of cultivation, and thereafter these accumulated immunoglobulin mRNAs and proteins are gradually reduced via lots of generations during cultivation. Such alterations in gene expressions of IgM H- and L-chains should be coupled with those of gene expressions of some of Pax5, Aiolos, EBF1, E2A, PCAF, HDAC7 and HDAC9 and others. In addition, changes in acetylation levels of at least Lys-9 residues of histone H3 (K9/H3) within chromatin structure surrounding the proximal ~2.0 kb 5'-upstream region of the Pax5 gene in HDAC2(-/-) should be a basis for changes in the transcription level of the gene through its chromatin structural change, resulting in

alterations in gene expressions of IgM H- and L-chains. However, unfortunately, the early cultivation stage in both of RT-PCR and CHIP assay was considerably delayed from that in other four experiments (2D-PAGE, Western blotting, immuno-blotting and immuno-electron microscopy), since the mutant cells used for the former two experiments had been already cultivated some times before use of the latter four studies and others. Therefore, to further elucidate the above-mentioned interesting biological phenomena, we must newly generate HDAC2-deficient mutant cells, collect them at early (as quickly as possible), middle and late cultivation stages, and analyze their cellular characteristics in detail at these three stages, each of which is the same cultivation period (as exactly as possible) for all of experiments that will be tested.

## **METHODS**

### **Cell cultures**

Generation of HDAC2(-/-) mutants was described in our previous report [26]. All positive mutant clones were picked up on 96-well plates at ~10 days after knockout of two HDAC2 alleles, cultivated for another several days to be  $\sim 10^6$  cells/ml and stocked at  $-80^{\circ}\text{C}$  until use. HDAC2(-/-) mutant clones were continuously cultivated as described [26, 33] and collected at indicated interval periods, including the early (E; ~10 or 20 days), middle (M; ~30 or 40 days) and late (L; ~56 or 60 days) stages of cultivation. The cultivation periods and/or stages were practically counted from the first stock day.

### **2D-PAGE**

Whole cellular proteins were prepared from HDAC2(-/-) mutant cells at the early (E; ~10 days), middle (M; ~30 days) and late (L; ~60 days) stages of cultivation and DT40 cells (W), all of which were exponentially growing and separated on 2D-PAGE using an automated apparatus Multiphor II (Amersham Pharmacia Biotech.), followed by staining with SYPRO Red (Molecular Dynamics, Inc.) as described [26, 46-48, 59].

### **Western blotting**

Cells were cultivated for ~56 days, and whole cellular proteins were prepared from HDAC2(-/-) mutant cells at different cultivation periods and DT40 cells (W) as described [27, 33]. Briefly, cells ( $1 \times 10^7$ ) were treated with 10% trichloroacetic acid (TCA) and lysed in 100  $\mu\text{l}$  of SDS buffer. Aliquots (10  $\mu\text{l}$ ) of 1:50 dilutions of the resultant cell extracts were subjected to 12% SDS-PAGE and electro-transferred to membranes, and proteins were detected with rabbit anti-chicken IgM L-chain antiserum A30-100A (Bethyl Laboratories Inc., TX, USA) that cross-reacts with IgM H-chain as described [30, 33]. Data analyses were carried out by Multi Gauge Ver3.X software using a luminescent

image analyzer LAS-1000plus (FUJIFILM) [30, 33]. The relative amounts of IgM H-chain, and large (high) and small (low) forms of IgM L-chain were measured at indicated times.

### **Immuno-electron microscopy**

Post-embedding immunocytochemistry was carried out as described [60]. Exponentially growing DT40 (W) and HDAC2(-/-) mutant cells at the early (E; ~10 days) and late (L; ~60 days) cultivation stages were collected by a light centrifugation, and the resultant cell pellets were fixed with 4% paraformaldehyde and 0.1% glutaraldehyde in 0.1 M phosphate buffer (PB) for 30 min. After washing with PB, the samples were postfixed with 1% osmium tetroxide in PB for 60 min, washed with PB, dehydrated in ethanol and embedded in Epon. Ultrathin sections were picked up on 200-mesh gold grids coated with Formvar film and treated with 5% sodium meta-periodate in distilled water for 30 min. After rinsing in distilled water, the sections were treated with 5% normal horse serum (NHS) and 1% bovine serum albumin (BSA) in phosphate buffered saline (PBS) for 10 min to block nonspecific binding and incubated with goat anti-chicken IgM H-chain antibody A30-102A (Bethyl Laboratories Inc., TX, USA; diluted 1:1500 with 5% NHS, 1% BSA in PBS) at room temperature for 60 min. After rinsing in PBS, the sections were incubated with biotinylated horse anti-goat IgG (Vector Laboratories, Burlingame, CA, USA; diluted 1:200 with 1% BSA in PBS) for 40 min. After washing with PBS, the sections were incubated with 8 nm colloidal gold conjugated anti-biotin antibody (Vector Laboratories, Burlingame, CA, USA; diluted with 1% BSA in PBS) for 30 min. After washing with distilled water and drying, the sections were contrasted with 2% uranyl acetate in 70% methanol and Reynolds' lead citrate and observed in a JEOL 1200EX transmission electron microscope operating at 80 kV (JEOL, Tokyo, JAPAN). As control, primary antibody was omitted or replaced by normal goat serum.

### **RT-PCR**

Total RNAs were isolated from three independent HDAC2(-/-) mutant clones (cl.33-28, cl.33-30 and cl.45-28) at the early (E; ~20 days), middle (M; ~40 days) and late (L; ~60 days) stages of cultivation and DT40 cells (W) as described [26, 33, 61]. RT-PCR was carried out using appropriate sense and anti-sense primers used in our previous papers [30, 33] and listed in Supplementary Table 2-SI.  $\beta$ -actin was used as control. RT-PCR products were subjected to 15% agarose gel electrophoresis. Data analyses were carried out as described above. Nucleotide sequences of all amplified products were confirmed by the PCR sequencing protocol as described [30, 33].

### **Immunoblotting**

Whole cellular proteins were prepared from three independent HDAC2(-/-) clones at the early (E; ~20 days), middle (M; ~40 days) and late (L; ~60 days) stages of cultivation and DT40 cells (W) (three

different cultures) as described [33]. Briefly, cells ( $1 \times 10^7$ ) were treated with 10% TCA and lysed in 100  $\mu$ l of SDS buffer. Aliquots (10  $\mu$ l) of the resultant cell extracts were subjected to 15% SDS-PAGE and electro-transferred to membranes. Bulk acetylation levels of particular Lys (K) residues (indicated by appropriate designations) of core histones H3, H4, H2A and H2B, and the bulk methylation level of Lys-9 residue of histone H3 were measured by immunoblotting using site-specific anti-acetylated and anti-methylated histone antibodies as described [33]. Antibodies used were: anti-AcK9H3 (#06-942), anti-MeK9H3 (#07-212), anti-AcK14H3 (#06-911), anti-AcK18H3 (#07-354), anti-AcK23H3 (#07-355), anti-AcK27H3 (#07-360), anti-AcK8H4 (#06-760), anti-AcK12H4 (#07-323), anti-AcK16H4 (#07-329), anti-AcK5H2A (#07-290), anti-AcK7H2A (#07-386), anti-AcK9H2A (#07-289), anti-AcK5H2B (#07-382), anti-AcK12H2B (#07-336), anti-AcK15H2B (#07-343), anti-AcK16H2B (#07-341) and anti-AcK20H2B (#07-347) from Upstate and anti-AcK5H4 (AB3064) from CHEMICON.

#### **Cloning of ~4.9 kb 5'-upstream fragment of chicken Pax5 gene**

Although the chicken Pax5 gene has been reported to be located on Z chromosome that is monosomy and to consist of 10 exons at the present time, the detailed information, including nucleotide sequences of the 5'-upstream region plus 5'-untranslational region of its first exon, was not available in USCS Genome Browser database until we had started this study. Only nucleotide sequences of chicken Pax5 cDNA [8] and 5'-upstream genomic DNA sequences of exon 1 up to position -109 (GenBank accession number; NM\_204424) had been reported. To clone and identify the ~4.9 kb 5'-upstream fragment of the Pax5 gene, we first screened the DT40 genomic DNA library in a  $\lambda$ FIXII by PCR using sense primer (5'-GCACACTACGGACAACTTTTCC-3') and antisense primer (5'-CGGGGCCGCGTACATCTTCTCCAAATCCAT-3'), which were designed to amplify a part of exon 1 of the gene. However, no positive clones were yielded.

Therefore, we directly cloned the ~4.9 kb 5'-upstream region of the Pax5 gene from DT40 genomic DNA by our original gene walking techniques, including Southern blotting [62], sub-cloning and colony hybridization (details will be shown later). First, the genomic DNA was digested with appropriate restriction enzymes, recognizing particular positions in multiple cloning site of pBluescript II SK(+) vector, and hybridized with the PCR amplicon (as probe), corresponding to the 85 bp fragment of exon 1 (positions -55 ~ +30), which was obtained from the genomic DNA using the above-mentioned primers. Among positive fragments obtained, the ~2.1 kb XbaI/ClaI fragment was sub-cloned into the multiple cloning site of pBluescript II SK(+) in frame as mentioned below. Enough amounts of genomic DNA (200  $\mu$ g) were digested with XbaI and ClaI at 37°C for 18 hr and separated in a 0.8% agarose gel, and the gel area containing the desired DNA fragment of ~2.1 kb was excised. The ~2.1 kb DNA fragments were extracted using the filter cartridge SUPEC<sup>TM</sup>-01 (TaKaRa) following the manufacture's protocol and recovered by phenol/chloroform extraction and ethanol precipitation. The purified DNAs were

dissolved in TE buffer (10 mM Tris-HCl, 1 mM EDTA; pH 8.0) and inserted into the XbaI/ClaI site of pBluescript II SK(+) in molar ratio of 1 to 10 (for vector to insert) using DNA Ligation Kit Ver.1 (TaKaRa) at 16°C for 21 hr. The ligated plasmid was introduced into XL-1 blue MRF Competent cells, plated onto LB/amp plates, incubated at 37°C for overnight, cooled at 4°C for 1 hr, and subjected to colony hybridization with the 85 bp PCR amplicon as probe. Our screening yielded one positive colony. Then, we sequenced the cloned the ~2.1 kb DNA fragment by the dye terminator method using Big Dye Terminator V3.1 Sequencing Standard kit (Applied Biosystems).

As the second step, using the similar techniques, we cloned and identified nucleotide sequences of more additional ~3.0 kb 5'-upstream region of the Pax5 gene as follows. Based on the 5'-upstream genomic sequences mentioned above, sense 5'-TGCCATTTTCAAACACGCGC-3' and antisense 5'-GGAAAAGTTTGTCCGTAGTGTGC-3' primers were constructed. The 1626 bp PCR amplified product (corresponding to positions -1656 ~ -32) was obtained from DT40 genomic DNA by PCR using these primers and digested with BssHII. After separation in 0.8% agarose gel, the band of desired 263 bp fragment was excised from the gel and purified using Wizard SV Gel and PCR Clean-Up System (Promega).

Resultant 263 bp DNA fragment, corresponding to the region from positions -1656 ~ -1394, was used as probe for subsequent hybridization to obtain more additional 5'-upstream genomic fragment cut with appropriate restriction enzymes. Among positive fragments obtained, the ~4.9 kb BamHI/ClaI fragment was inserted into the BamHI/ClaI site of pBluescript II SK(+) in molar ratio of 1 to 5 (for vector to insert), introduced into XL-1 blue MRF Competent cells, plated onto LB/amp plates and subjected to colony hybridization. Finally, we isolated three positive clones, possessing the ~4.9 kb 5'-upstream fragment from nucleotide number +241, and sequenced this isolated DNA fragment as described above.

### **Colony hybridization**

We carried out colony hybridization on Hybond-N+ nylon membrane filter following the manufacture's protocol (Amersham Biosciences). In brief, the membrane was laid on the surface of agar plates for 45 sec until it became thoroughly wetted, peeled carefully, and then its colony side was placed in upper direction onto a 10%(w/v) SDS saturated 3MMChr paper (Whatman) for 3 min. DNA was denatured and fixed to the membrane by successive treatments with denaturing buffer (0.5 M NaOH, 1.5 M NaCl) for 7 min and neutralization buffer (1.5 M NaCl, 0.5 M Tris-HCl; pH 7.4) for 3 min twice. Finally, the membrane was vigorously washed in 2X SSC (0.3 M NaCl, 30 mM C<sub>6</sub>H<sub>5</sub>O<sub>7</sub>Na<sub>3</sub>·2H<sub>2</sub>O; pH7.0) and its surface was gently scraped using KIMWIPE soaked in 2X SSC to remove cell debris. DNA was fixed to the membrane by a UV cross-linking procedure and hybridized with <sup>32</sup>P-labeled probe by means of conventional methods.

### **Chromatin immunoprecipitation assay**

Chromatin immunoprecipitation (ChIP) assay was done with ChIP assay kit (Millipore) according to manufacturer's instructions. Briefly, DT40 (W) and HDAC2(-/-) mutant clones (cl.33-28, cl.33-30 and cl.45-28) ( $1 \times 10^6$  cells) at the early (E; ~20 days) and late (L; ~60 days) stages of cultivation were cross-linked by the addition of 37% formaldehyde to a final concentration of 1% and incubated at 37°C for 10 min, and the cross-linking was stopped with the addition of one-tenth volume of 1.25 M glycine to a final concentration of 0.125 M at 37°C for 5 min. After removing medium, cells were washed with ice-cold PBS twice and resuspended in 200  $\mu$ l of SDS Lysis Buffer freshly supplemented with 1 mM phenylmethylsulfonyl fluoride (PMSF) (Wako, Osaka, Japan) and 1  $\mu$ g/ml aprotinin (Sigma, St Louis, MO, USA), followed by sonication to solubilize and to shear cross-linked DNA. At this step, we used a Bioruptor UCD-250 (Cosmo Bio) at power L for ten 30-s pulses (30-s pause between pulses) at 4°C to shear chromatin DNA into lengths of approximately 200 ~ 1000 bp. The samples were centrifuged at 13,000 rpm at 4°C for 10 min to remove cell debris from the crude chromatin lysate. Sonicated cell supernatant was diluted to 1/10-fold by Chip Dilution Buffer. For input, a portion (40  $\mu$ l; 2%) of the diluted cell supernatant was kept to quantify the amount of DNA present in each of different samples at PCR step. Antibody (2  $\mu$ l) was added to 2 ml of the supernatant fraction and incubated at 4°C for overnight. Incubation with 60  $\mu$ l of Protein Agarose/Salmon Sperm DNA was continued for 1 hr, followed by centrifugation at 1,000 rpm at 4°C for 1 min. The immuno-precipitated complexes were sequentially washed with low-salt solution, high-salt solution, LiCl solution and TE (pH 8.0) twice. The antibody/histone/DNA complexes were eluted from Protein Agarose beads by adding 250  $\mu$ l of the elution buffer (1% SDS, 0.1% NaHCO<sub>3</sub>) twice, and the cross-linking was reversed at 65°C for 6.5 hr by the addition of 5 M NaCl to a final concentration of 0.2 M. Samples were treated with proteinase K at 37°C for 1 hr to remove proteins. DNA was recovered by phenol/chloroform extraction and ethanol precipitation using 40  $\mu$ g glycogen (NAKARAI TESQUE, INC.) as inner carrier. Anti-AcK9H3 (#07-352) (Millipore) was used for pulldown. Recovered DNA fragments were subjected to PCR for 25 ~ 35 cycles, using several appropriate primers specific for approximately 150 ~ 250 bp segments, corresponding to proximal 5'-upstream region of the Pax5 gene (from the start codon) and first exon, which are listed in Supplementary Table 2-SII. PCR was carried out using the same master reaction mixture in 25- $\mu$ l scale. Immuno-precipitated chromatin treated with non-immune rabbit serum (Vector Laboratories) (as negative control) produced no bands following PCR (data not shown).

### **REFERENCES**

- [1] Kornberg, R. D.: Structure of chromatin. *Annu. Rev. Biochem.* 46, 931-954, 1977.
- [2] van Holde, K. E.: *Chromatin*. New York: Springer-Verlag, 1989.

- [3] Workman, J. L. and Kingston, R. E.: Alteration of nucleosome structure as a mechanism of transcriptional regulation. *Annu. Rev. Biochem.* 67, 545-579, 1998.
- [4] Peterson, M. L.: RNA processing and expression of immunoglobulin genes. In *Handbook of B and T Lymphocytes* (Snow, E. C. ed.) pp321-342, 1994, Academic press, San Diego.
- [5] Busslinger, M.: Transcriptional control of early B cell development. *Annu. Rev. Immunol.* 22, 55-79, 2004.
- [6] Nutt, S. L. and Kee, B. L.: The transcriptional regulation of B cell lineage commitment. *Immunity* 26, 715-725, 2007.
- [7] Ramirez, J., Lukin, K. and Hagman, J.: From hematopoietic progenitors to B cells: mechanisms of lineage restriction and commitment. *Curr. Opin. Immunol.* 22, 177-184, 2010.
- [8] Funahashi, J., Okufuji, T., Ohuchi, H., Noji, S., Tanaka, H. and Nakamura, H.: Role of Pax-5 in the regulation of a mid-hindbrain organizer's activity. *Dev. Growth Differ.* 41, 59-72, 1999.
- [9] Stevens, S., Ong, J., Kim, U., Eckhardt, L. A. and Roeder, R. G.: Role of OCA-B in 3'-IgH enhancer function. *J. Immunol.* 164, 5306-5312, 2000.
- [10] Savitsky, D. and Calame, K.: B-1 B lymphocytes require Blimp-1 for immunoglobulin secretion. *J. Exp. Med.* 203, 2305-2314, 2006.
- [11] Nera, K.-P., Kohonen, P., Narvi, E., Peippo, A., Mustonen, L., Terho, P., Koskele, K., Buerstedde, J.-M. and Lassila, O.: Loss of Pax5 promotes plasma cell differentiation. *Immunity* 24, 283-293, 2006.
- [12] Fujita, T. and Fujii, H.: Species-specific 5'-genomic structure and multiple transcription start sites in the chicken Pax5 gene. *Gene* 477, 24-31, 2011.
- [13] John, L. B. and Ward, A. C.: The Ikaros gene family: transcriptional regulators of hematopoiesis and immunity. *Mol. Immunol.* 48, 1272-1278, 2011.
- [14] Brownell, J. E., Zhou, J., Rannali, T., Kobayashi, R., Edmondson, D. G., Roth, S. Y. and Allis, C. D.: Tetrahymena histone acetyltransferase A: a homolog to yeast Gcn5 linking histone acetylation to gene activation. *Cell* 84, 843-851, 1996.
- [15] Ogryzko, V. V., Schiltz, R. L., Russanova, V., Howard, B. H. and Nakatani, Y.: The transcriptional coactivators p300 and CPB are histone acetyltransferases. *Cell* 87, 953-959, 1996.
- [16] Taunton, J., Hassig, C. A. and Schreiber, S. L.: A mammalian histone deacetylase related to the yeast transcriptional regulator Rpd3p. *Science* 272, 408-411, 1996.
- [17] Brown, C. E., Lechner, T., Howe, L. and Workman, J. L.: The many HATs of transcription coactivators. *Trends Biochem. Sci.* 25, 15-19, 2000.
- [18] Cheung, W. L., Briggs, S. D. and Allis, C. D.: Acetylation and chromosomal functions. *Curr. Opin. Cell Biol.* 12, 326-333, 2000.
- [19] Turner, B. M.: Histone acetylation and an epigenetic code. *Bioessays* 22, 836-845, 2000.
- [20] Jenuwein, T. and Allis, C. D.: Translating the histone code. *Science* 293, 1074-1080, 2001.

- [21] Roth, S. Y. Denu, J. M. and Allis, C. D.: Histone acetyltransferases. *Annu. Rev. Biochem.* 70, 81-120, 2001.
- [22] Carrozza, M. J., Utley, R. T., Workman, J. L. and Cote, J.: The diverse functions of histone acetyltransferase complexes. *Trend. Genet.* 19, 321-329, 2003.
- [23] Yang, X. J. and Seto, E.: Collaborative spirit of histone deacetylases in regulating chromatin structure and gene expression. *Curr. Opin. Genet. Dev.* 13, 143-153, 2003.
- [24] Allis, C. D., Berger, S. L., Cote, J., Dent, S., Jenuwien, T., Kouzarides, T., Pillus, L., Reinberg, D., Shi, Y., Shiekhatar, R., Shilatifard, A., Workman, J. and Zhang, Y.: New nomenclature for chromatin-modifying enzymes. *Cell* 131, 633-636, 2007.
- [25] Lee, K. K. and Workman, J. L.: Histone acetyltransferase complexes: one size doesn't fit all. *Nat. Rev. Mol. Cell Biol.* 8, 284-295, 2007.
- [26] Takami, Y., Kikuchi, H. and Nakayama, T.: Chicken histone deacetylase-2 controls the amount of the IgM H-chain at the steps of both transcription of its gene and alternative processing of its pre-mRNA in the DT40 cell line. *J. Biol. Chem.* 274, 23977-23990, 1999.
- [27] Takami, Y. and Nakayama, T.: N-terminal region, C-terminal region, nuclear export signal and deacetylation activity of histone deacetylase-3 are essential for the viability of the DT40 chicken B cell line. *J. Biol. Chem.* 275, 16191-16201, 2000.
- [28] Nakayama, T. and Takami, Y.: Participation of histones and histone-modifying enzymes in cell functions through alterations in chromatin structure. *J. Biochem.* 129, 491-499, 2001.
- [29] Takechi, S., Adachi, M. and Nakayama, T.: Chicken HDAC2 down-regulates IgM light chain gene promoter activity. *Biochem. Biophys. Res. Commun.* 299, 263-267, 2002.
- [30] Kikuchi, H., Takami, Y. and Nakayama, T.: GCN5: a supervisor in all-inclusive control of vertebrate cell cycle progression through transcription regulation of various cell cycle-related genes. *Gene* 347, 83-97, 2005.
- [31] Kikuchi, H., Barman, H. K., Nakayama, M., Takami, Y. and Nakayama, T.: Participation of histones, histone modifying enzymes and histone chaperones in vertebrate cell functions. *Reviews and Protocols in DT40 Research*, Springer-Verlag, Berlin, pp225-243, 2006.
- [32] Barman, H. K., Takami, Y., Ono, T., Nishijima, H., Sanematsu, F., Shibahara, K. and Nakayama, T.: Histone acetyltransferase 1 is dispensable for replication-coupled chromatin assembly but contributes to recover DNA damages created following replication blockage in vertebrate cells. *Biochem. Biophys. Res. Commun.* 345, 1547-1557, 2006.
- [33] Nakayama, M., Suzuki, H., Yamamoto-Nagamatsu, N., Barman, H. K., Kikuchi, H., Takami, Y., Toyonaga, K., Yamashita, K. and Nakayama, T.: HDAC2 controls IgM H and L-chain gene expressions via EBF1, Pax5, Ikaros, Aiolos and E2A gene expressions. *Genes Cells* 12, 359-373, 2007.

- [34] Kikuchi, H. and Nakayama, T.: GCN5 and BCR signaling collaborate to induce pre-mature B cell apoptosis through depletion of ICAD and IAP2 and activation of caspase activities. *Gene* 419, 48-55, 2008.
- [35] Barman, H. K., Takami, Y., Nishijima, H., Shibahara, K., Sanematsu, F. and Nakayama, T.: Histone acetyltransferase-1 regulates integrity of cytosolic histone H3-H4 containing complex. *Biochem. Biophys. Res. Commun.* 373, 624-630, 2008.
- [36] Kikuchi, H., Barman, H. K., Nakayama, M., Takami, Y. and Nakayama, T.: Studies on epigenetic control of B cell functions using the DT40 cell line. *Advances in Genetics Research* 2, Urbano K. V. (Ed.), Nova Science Publishers, Inc. NY, pp153-166, 2010.
- [37] Kikuchi, H., Kuribayashi, F., Takami, Y., Imajoh-Ohmi, S. and Nakayama, T.: GCN5 regulates the activation of PI3K/Akt survival pathway in B cells exposed to oxidative stress via controlling gene expressions of Syk and Btk. *Biochem. Biophys. Res. Commun.* 405, 657-661, 2011.
- [38] Kikuchi, H., Kuribayashi, F., Kiwaki, N., Takami, Y. and Nakayama, T.: GCN5 regulates the superoxide-generating system in leukocytes via controlling gp91-phox gene expression. *J. Immunol.* 186, 3015-3022, 2011.
- [39] Kikuchi, H., Kuribayashi, F., Imajoh-Ohmi, S., Nishitoh, Y., Takami, Y. and Nakayama, T.: GCN5 protects vertebrate cells against UV-irradiation via controlling gene expression of DNA polymerase  $\eta$ . *J. Biol. Chem.* 287, 39842-39849, 2012.
- [40] Kikuchi, H., Nakayama, M., Kuribayashi, F., Imajoh-Ohmi, S., Nishitoh, H., Takami, Y. and Nakayama T.: GCN5 is essential for IRF-4 gene expression followed by transcriptional activation of Blimp-1 in immature B cells. *J. Leukoc. Biol.* 95, 399-404, 2014.
- [41] Kikuchi, H., Nakayama, M., Kuribayashi, F., Imajoh-Ohmi, S., Nishitoh, H., Takami, Y. and Nakayama T.: GCN5 is involved in regulation of immunoglobulin heavy chain gene expression in immature B cells. *Gene* 544, 19-24, 2014.
- [42] Escaffit, F., Vaute, O., Chevillard-Briet, M., Segui, B., Takami, Y., Nakayama, T. and Trouche, D.: Cleavage and cytoplasmic relocalization of histone deacetylase 3 are important for apoptosis progression. *Mol. Cell. Biol.* 27, 554-567, 2007.
- [43] Takami, Y., Takeda, S. and Nakayama, T.: Targeted disruption of an H3-IV/H3-V gene pair causes increased expression of the remaining H3 genes in the chicken DT40 cell line. *J. Mol. Biol.* 250, 420-433, 1995.
- [44] Seguchi, K., Takami, Y. and Nakayama, T.: Targeted disruption of 01H1 encoding a particular H1 histone variant causes changes in protein patterns in the DT40 chicken B cell line. *J. Mol. Biol.* 254, 869-880, 1995.

- [45] Takami, Y., Takeda, S. and Nakayama, T.: Targeted disruption of H2B-V encoding a particular H2B histone variant causes changes in protein patterns on two-dimensional polyacrylamide gel electrophoresis in the DT40 chicken B cell line. *J. Biol. Chem.* 270, 30664-30670, 1995.
- [46] Takami, Y., Takeda, S. and Nakayama, T.: An approximately half set of histone genes is enough for cell proliferation and a lack of several histone variants causes protein pattern changes in the DT40 chicken B cell line. *J. Mol. Biol.* 265, 394-408, 1997.
- [47] Takami, Y. and Nakayama, T.: One allele of the major histone gene cluster is enough for cell proliferation of the DT40 chicken B cell line. *Biochim. Biophys. Acta* 1354, 105-115, 1997.
- [48] Takami, Y. and Nakayama, T.: A single copy of linker H1 genes is enough for proliferation of the DT40 chicken B cell line, and linker H1 variants participate in regulation of gene expression. *Genes Cells* 2, 711-723, 1997.
- [49] Sanematsu, F., Takami, Y., Barman, H. K., Fukagawa, T., Ono, T., Shibahara, K. and Nakayama, T.: Asf1 is required for viability and chromatin assembly during DNA replication in vertebrate cells. *J. Biol. Chem.* 281, 13817-13827, 2006.
- [50] Takami, Y., Ono, T., Fukagawa, T., Shibahara, K. and Nakayama, T.: Essential role of CAF-1-mediated rapid nucleosome assembly for DNA replication and cell division in vertebrate cells. *Mol. Biol. Cell* 18, 129-141, 2007.
- [51] Baba, T. W., Giroir, B. P. and Humphries, E. H.: Cell lines derived from avian lymphomas exhibit two distinct phenotypes. *Virology* 144, 139-151, 1985.
- [52] Buerstedde, J.-M. and Takeda, S.: Increased ratio of targeted to random integration after transfection of chicken B cell lines. *Cell* 67, 179-188, 1991.
- [53] Takeda, S., Masteller, E. L., Thompson, C. B. and Buerstedde, J.-M.: RAG-2 expression is not essential for chicken immunoglobulin gene conversion. *Proc. Natl. Acad. Sci. USA* 89, 4023-4027, 1992.
- [54] Toyonaga, K., Kikuchi, H., Yamashita, K., Nakayama, M., Chijiwa, K. and Nakayama, T.: E2A participates in a fine control of pre-mature B cell apoptosis mediated by B cell receptor signaling via transcription regulations of survivin, IAP2 and caspase-8 genes. *FEBS J.* 276/5, 1418-1428, 2009.
- [55] Kikuchi, H., Yamashita, K., Nakayama, M., Toyonaga, K., Tsuneyoshi, I., Takasaki, M. and Nakayama, T.: Lacking of Aiolos accelerates pre-mature B cell apoptosis mediated by BCR signaling through elevation in cytochrome c release. *BBA-Molecular Cell Research* 1793, 1304-1314, 2009.
- [56] Kikuchi, H., Nakayama, M., Takami, Y., Kuribayashi, F. and Nakayama, T.: Possible involvement of Helios in controlling the immature B cell functions via transcriptional regulation of protein kinase Cs. *Results Immunol.* 1, 88-94, 2011.
- [57] Kikuchi, H., Nakayama, M., Takami, Y., Kuribayashi, F. and Nakayama, T.: EBF1 acts as a powerful repressor of Blimp-1 gene expression in immature B cells. *Biochem. Biophys. Res. Commun.* 422, 780-785, 2012.

- [58] Kikuchi, H., Nakayama, M., Kuribayashi, F., Imajoh-Ohmi, S., Nishitoh, H., Takami, Y. and Nakayama, T.: Protein kinase C  $\theta$  gene expression is oppositely regulated by EBF1 and GCN5 in immature B cells. *FEBS Lett.* 588, 1739-1742, 2014.
- [59] Takami, Y., Nishi, R. and Nakayama, T.: Histone H1 variants play individual roles in transcription regulation in the DT40 chicken B cell line. *Biochem. Biophys. Res. Commun.* 268, 501-508, 2000.
- [60] Bendayan, M. and Zollinger, M.: Ultrastructural localization of antigenic sites on osmium-fixed tissues applying the protein A-gold technique. *J. Histochem. Cytochem.* 31, 101-109, 1983.
- [61] Chomczynski, P. and Sacchi, N.: Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* 162, 156-159, 1987.
- [62] Southern, E. M.: Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 98, 503-517, 1975.

## FIGURE LEGENDS

### **Figure 2-1. Alterations in amounts of IgM H- and L-chains in HDAC2(-/-) DT40 mutant cells during cultivation**

Total cellular proteins were prepared from HDAC2(-/-) mutant cells (clone cl.33-12) at the early (E; ~10 days), middle (M; ~30 days) and late (L; ~60 days) stages of cultivation and compared with those from DT40 cells (W) by 2D-PAGE. Iso-electrofocusing in the first dimension (pI) and SDS-PAGE in the second dimension (MW) were performed on Immobiline DryStrip gel (pH 4 - 7) and ExcelGel XL SDS gel (gradient 12 - 14), followed by the fluorostaining method. H and L indicate IgM H- and L-chains, respectively.

### **Figure 2-2. Alterations in protein levels of IgM H- and L-chains in HDAC2(-/-) DT40 mutant cells during cultivation**

Total cellular proteins were prepared from HDAC2(-/-) mutant cells (clone cl.33-28) collected at indicated cultivation periods and compared with those from DT40 cells (W) by Western blotting. Aliquots of cell extracts were subjected to 12% SDS-PAGE and transferred to membrane filter. Proteins were detected with anti-chicken IgM L-chain antiserum (as primary antibody) that cross-reacts with IgM H-chain. Relative levels of IgM H-chain and large (high) and small (low) forms of IgM L-chain are shown in the lower panel. IgM H and IgM L indicate IgM H- and L-chains, respectively.

### **Figure 2-3. Alterations in amounts of IgM H-chain proteins in HDAC2(-/-) DT40 mutant cells during cultivation**

HDAC2(-/-) mutant cells (clone cl.33-28) collected at the early (E; ~10 days) and late (L; ~60 days) stages of cultivation and DT40 cells (W) were fixed, and immuno-electron microscopy with anti-chicken IgM H-chain antiserum was performed. A) Study in wide range. A number of cells were observed in (W), (E) and (L), but accumulated IgM H-chain proteins were detected only in mutant cells in (E). B) Study for single cell. Each of cells indicated by arrows in A) was magnified and shown in (W), (E) or (L). A large amount of accumulated IgM H-chain proteins were observed only in (E).

### **Figure 2-4. Alterations in gene expressions of IgM H- and L-chains, and core histones in HDAC2(-/-) DT40 mutant cells during cultivation**

Total RNAs were extracted from three HDAC2(-/-) mutant clones (cl.33-28, cl.33-30 and cl.45-28) at the early (E; ~20 days), middle (M; ~40 days) and late (L; ~60 days) stages of cultivation and DT40 cells (W). RT-PCR was performed using equal amounts of total RNAs and appropriate primers for HDAC2, whole (IgM Hc), secreted (IgM Hs) plus membrane-bound forms (IgM Hm) of IgM H-chain and IgM L-chain (IgM L), and four core histones (H2A, H2B, H3 and H4). Chicken  $\beta$ -actin was used as control.

Some data for DT40 (W) and HDAC2(-/-) at the early (E) stage were the same as those in our previous paper [33].

**Figure 2-5. Alterations in gene expressions of HDACs, HATs, and transcription factors in HDAC2(-/-) DT40 mutant cells during cultivation**

Total RNAs used were the same ones as in Fig. 2-4. RT-PCR was performed in the same manner, using appropriate primers for members of HDACs and HATs (A) and transcription factors (B), instead of primers used in Fig. 2-4. Some data for DT40 (W) and HDAC2(-/-) at the early (E) stage were the same as those in our previous paper [33].

**Figure 2-6. Alterations in bulk acetylation and methylation levels of Lys residues of histone H3 in HDAC2(-/-) DT40 mutant cells during cultivation**

Whole cellular proteins were extracted from three HDAC2(-/-) mutant clones (cl.33-28, cl.33-30 and cl.45-28) at the early (E; ~20 days), middle (M; ~40 days) and late (L; ~60 days) stages of cultivation and DT40 cells (W) (three separate cultures) and subjected to 15% SDS-PAGE. Bulk acetylation levels of particular Lys (K) residues (indicated by appropriate designations) of core histones H3, H4, H2A and H2B were measured by immunoblotting using various site-specific anti-acetylated histone antibodies. Bulk methylation level of K9/H3 was measured using site-specific anti-methylated histone antibody. Most data for DT40 (W) and HDAC2(-/-) at the early (E) stage were the same as those in our previous paper [33].

**Figure 2-7. Alterations in acetylation levels of Lys-9 residues of histone H3 within proximal 5'-upstream chromatin region of Pax5 gene in HDAC2(-/-) DT40 mutant cells during cultivation**

Chromatin immunoprecipitation (ChIP) assay was carried out using Chromatin Immunoprecipitation Assay Kit. Cross-linked chromatins prepared from cell lysates of three HDAC2(-/-) mutant clones (cl.33-28, cl.33-30 and cl.45-28) at the early (E; ~20 days) and late (L; ~60 days) stages of cultivation and DT40 cells (W) were co-precipitated by antiserum specific for acetylated Lys-9 residue of histone H3 (K9/H3). After decross-linking, co-precipitated chromatins were amplified by PCR using appropriate primers A to Q (Suppl. Table 2-SII) for the proximal 5'-upstream region of the Pax5 gene (see Suppl. Fig. 2-S2). PCR products were analyzed by 1.5% agarose gel electrophoresis.

**Supplementary Figure 2-S1. Alterations in protein levels of IgM H- and L-chains in two HDAC2(-/-) DT40 mutant clones (cl.33-30 and cl.45-28) during cultivation**

Total cellular proteins were prepared from HDAC2(-/-) mutant clones (cl.33-30 and cl.45-28) collected at indicated cultivation periods and analyzed in the same manner in Fig. 2-2. However, these

periods were not coincident with those in Fig. 2-2.

**Supplementary Figure 2-S2. Nucleotide sequences of the 5'-upstream region of chicken Pax5 gene**

Nucleotide sequences of the 5'-upstream region of the chicken Pax5 gene up to the position -4950 (including exon 1, putative transcription start site and translation initiation triplet) are shown.