

# Zebrafish tyrosylprotein sulfotransferase: molecular cloning, expression, and functional characterization

Emi Mishiro, Ming-Yih Liu, Yoichi Sakakibara, Masahito Suiko, and Ming-Cheh Liu

**Abstract:** By employing the reverse transcriptase – polymerase chain reaction technique in conjunction with 3' rapid amplification of cDNA ends, a full-length cDNA encoding a zebrafish tyrosylprotein sulfotransferase (TPST) was cloned and sequenced. Sequence analysis revealed that this zebrafish TPST is, at the amino acid sequence level, 66% and 60% identical to, respectively, the human and mouse TPST-1 and TPST-2. The recombinant form of the zebrafish TPST, expressed in COS-7 cells, exhibited a pH optimum at 5.75. Manganese appeared to exert a stimulatory effect on the zebrafish TPST. The activity of the enzyme determined in the presence of 20 mM MnCl<sub>2</sub> was more than 2.5 times that determined in the absence of MnCl<sub>2</sub>. Of the other nine divalent metal cations tested at a 10 mM concentration, Co<sup>2+</sup> also showed a considerable stimulatory effect, while Ca<sup>2+</sup>, Pb<sup>2+</sup>, and Cd<sup>2+</sup> exerted some inhibitory effects. The other four divalent cations, Fe<sup>2+</sup>, Cu<sup>2+</sup>, Zn<sup>2+</sup>, and Hg<sup>2+</sup>, inhibited completely the sulfating activity of the zebrafish TPST. Using the wild-type and mutated P-selectin glycoprotein ligand-1 N-terminal peptides as substrates, the zebrafish TPST was shown to exhibit a high degree of substrate specificity for the tyrosine residue on the C-terminal side of the peptide. These results constitute a first study on the cloning, expression, and characterization of a zebrafish cytosolic TPST.

*Key words:* XXXXXXXXX.

**Résumé :** À l'aide de la technique d'amplification en chaîne par polymérase après transcription inverse (RT-PCR) et de l'amplification rapide des extrémités 3' d'ADNc (3'-RACE), un ADNc pleine longueur codant une tyrosylprotéine sulfotransférase (TPST) du poisson-zèbre a été cloné et séquencé. L'analyse de la séquence des acides aminés montre que cette TPST du poisson-zèbre est, respectivement, 66% et 60% identique aux TPST-1 et TPST-2 de la souris et humaines. La TPST recombinante du poisson-zèbre, exprimée dans des cellules COS-7, a un pH optimum de 5,75. Le manganèse semble avoir un effet stimulant sur la TPST du poisson-zèbre. L'activité de l'enzyme en présence de MnCl<sub>2</sub> 20 mM est 2,5 fois plus élevée qu'en absence de MnCl<sub>2</sub>. Des neuf autres cations divalents métalliques évalués à une concentration de 10 mM, le Co<sup>2+</sup> a également un effet stimulant considérable, alors que le Ca<sup>2+</sup>, le Pb<sup>2+</sup> et le Cd<sup>2+</sup> ont un petit effet inhibiteur. Les quatre autres cations divalents, le Fe<sup>2+</sup>, le Cu<sup>2+</sup>, le Zn<sup>2+</sup> et le Hg<sup>2+</sup>, inhibent complètement l'activité de sulfatation de la TPST du poisson-zèbre. En utilisant les peptides N-terminaux du ligand glycoprotéique 1 de la P-sélectine (PSGL-1) de type sauvage ou muté comme substrats, nous montrons que la TPST du poisson-zèbre a une très grande spécificité de substrat envers le résidu tyrosine du côté C-terminal du peptide. Ceci constitue la première étude sur le clonage, l'expression et la caractérisation d'une TPST cytosolique du poisson-zèbre.

*Mots clés :* (non transmis).

[Traduit par la Rédaction]

## Introduction

The structure of a mature protein is dependent on not only its amino acid sequence but also on various covalent modifications that take place after ribosomal protein synthesis. These posttranslational protein modifications, including glycosylation,

proteolytic cleavage, phosphorylation, acylation, adenylation, farnesylation, ubiquitination, and sulfation, abound among proteins in eukaryotic cells (Tuboi et al. 1992). First discovered in a peptide (fibrinopeptide B) derived from fibrinogen (Bettelheim 1954), protein tyrosine sulfation has emerged as a widespread post-translational modification in multicellular

Received 1 August 2003. Revision received 18 November 2003. Accepted 1 December 2003. Published on the NRC Research Press Web site at <http://beb.nrc.ca> on XX XXXX 2004.

E. Mishiro, Y. Sakakibara, and M. Suiko. Biomedical Research Center, University of Texas Health Center, Tyler, TX 75708, U.S.A., and Department of Biochemistry and Applied Biosciences, Miyazaki University, Miyazaki 889-2192, Japan.

M.-Y. Liu. Department of Biochemistry and Molecular Biology, University of Georgia, Athens, GA 30602, U.S.A.

M.-C. Liu.<sup>1</sup> Biomedical Research Center, University of Texas Health Center, Tyler, TX 75708, U.S.A.

<sup>1</sup>Corresponding author (e-mail: [ming.liu@uthct.edu](mailto:ming.liu@uthct.edu)).

eukaryotes (Huttner 1982). It has been shown to be involved in the alteration of biological activity of proteins (Jensen et al. 1981; Brand et al. 1984), proteolytic processing of bioactive peptides (Rosa et al. 1985), change in half-life of proteins in circulation (Pauwels et al. 1987), intracellular transport of secretory proteins (Hille et al. 1984), and modulation of extracellular protein-protein interactions (Kehoe and Bertozzi 2000). More recently, tyrosine sulfation has been demonstrated to be a key modulator of protein-protein interactions that mediate inflammatory leukocyte adhesion (Pouyani and Seed 1995; Sako et al. 1995; Wilkins et al. 1995). The recent discovery of tyrosine-sulfated chemokine receptors (Farzan et al. 1999) suggests an even broader role in the inflammatory response.

Tyrosylprotein sulfotransferase (TPST), the enzyme responsible for protein tyrosine sulfation, catalyzes the transfer of a sulfonyl group from 3'-phosphoadenosine 5'-phosphosulfate (PAPS), the "active" sulfate, to specific tyrosine residues of proteins (Lee and Huttner 1983). This enzyme, following its first description in rat pheochromocytoma 12 cells (Lee and Huttner 1983), has been found in numerous cell lines and tissues and has been characterized from the bovine adrenal medulla (Lee and Huttner 1985) and heart (Suiko et al. 1997), rat brain (Vargas et al. 1985), liver (Rens-Domiano and Roth 1989; Rens-Domiano et al. 1989; Ramaprasad and Kashinathan 1998), gastric mucosa (Kashinathan et al. 1992), submandibular salivary glands (Sundaram et al. 1992; William et al. 1997), human liver (Lin and Roth 1990; Young 1990), and platelets (Sane and Baker 1993). Comparison of the properties reported for TPSTs from the mammalian tissues mentioned above reveals some common properties including acidic pH optima, stimulation by  $Mn^{2+}$ , and the use of PAPS as the sulfate donor. Biochemical evidence indicates that TPST is an integral membrane protein with a lumenally oriented active site localized in the trans-Golgi network. Tyrosine sulfation was demonstrated to take place in the same compartment as galactosylation and sialylation (i.e., the trans-Golgi), occurring shortly after these terminal glycosylation reactions (Baeuerle and Huttner 1987). A major advance in the field came with the molecular cloning of two distinct TPSTs, designated TPST-1 and TPST-2, from both human and mouse (Ouyang et al. 1998; Beisswanger et al. 1998; Ouyang and Moore 1998). Both enzymes were shown to be membrane-bound, *N*-glycosylated Golgi enzymes with a luminal catalytic domain, a single-span transmembrane domain, and a short cytosolic tail. The two human TPSTs share 67% sequence identity. Northern analysis indicated that both human TPSTs are expressed in many tissues (Ouyang et al. 1998; Beisswanger et al. 1998; Ouyang and Moore 1998). The existence of more than one TPST might explain the diversity of sequences that are tyrosine sulfated. The two enzymes might have differential substrate specificity and act upon different subsets of target proteins. A recent gene-knockout study revealed that mice deficient in TPST-1 had reduced body mass and increased postimplantation fetal death (Ouyang et al. 2002).

Zebrafish has in recent years emerged as a popular animal model for a wide range of studies (Briggs 2002; Ward and Lieschke 2002). Its advantages, compared with mouse, rat, or other vertebrate animal models, include the small size (3–4 cm length for adult fish), the availability of relatively large

number of eggs (>200 eggs per week per mature female), rapid development externally of virtually transparent embryo (in 2–4 days), and short generation time (~3 months). These unique characteristics of the zebrafish make it an excellent model for a systematic investigation on the ontogeny of the expression of various proteins and their tissue- and cell-type-specific distribution as well as physiological relevance.

We report here the molecular cloning of a full-length zebrafish TPST cDNA. Using COS-7 cells as the host, the zebrafish TPST was expressed in a recombinant form. The enzymatic properties of the recombinant zebrafish TPST were analyzed using as substrates wild-type and a series of mutated P-selectin glycoprotein ligand-1 (PSGL-1) N-terminal peptides.

## Experimental procedures

### Materials

Adenosine 5'-triphosphate (ATP), sodium dodecyl sulfate, *N*-2-hydroxylpiperazine-*N'*-2-ethanesulfonic acid (Hepes), 3-[*N*-tris-(hydroxymethyl)methylamino]-propanesulfonic acid, Trizma base, and dithiothreitol were products of the Sigma Chemical Company (St. Louis, Mo.). A zebrafish cDNA clone, ID 2644043 (GenBank accession No. AW344308), encoding the N-terminal region of a TPST was obtained from Genome Systems, Inc. AmpliTaq DNA polymerase was a product of Perkin Elmer. Takara *Ex Taq* DNA polymerase and the 3'-Full RACE core set were purchased from PanVera Corporation.  $T_4$  DNA ligase and all restriction endonucleases were from New England Biolabs. XL1-Blue MRF' *Escherichia coli* host strain was purchased from Stratagene (La Jolla, Calif.). Eukaryotic TOPO TA cloning kit and Lipofectamine 2000 were products of Invitrogen (Carlsbad, Calif.). TNT coupled reticulocyte lysate system was from Promega (Madison, Wis.). Oligonucleotide primers were synthesized by MWG Biotech. Recombinant human bifunctional ATP sulfurylase/adenosine 5'-phosphosulfate kinase was prepared as described previously (Yanagisawa et al. 1998). COS-7 SV40 transformed African green monkey kidney cells (ATCC CRL1651) were from the American Type Culture Collection. TRI reagent was from Molecular Research Center, Inc. Total RNA from zebrafish was prepared using the TRI reagent according to the manufacturer's instructions. Sep-Pak Plus C18 cartridges were from Waters. Cellulose thin-layer chromatography plates were products of EM Science. Carrier-free sodium [ $^{35}$ S]sulfate and [ $^{35}$ S]methionine were from ICN Biomedicals. All other reagents were of the highest grades commercially available.

### Molecular cloning of the zebrafish TPST

By searching the expressed sequence tag database, a zebrafish cDNA clone (GenBank accession No. AW344308) encoding the N-terminal region of a TPST was identified. This zebrafish TPST cDNA was purified and subjected to nucleotide sequencing based on the cycle sequencing method using, respectively, M13 forward and M13 reverse as primers. The nucleotide sequences, as well as the deduced amino acid sequences, of the cDNA were analyzed using BLAST search for sequence homology to known TPSTs.

To obtain the 3'-coding region and the untranslated sequence further downstream, 3' rapid amplification of cDNA

ends (3'-RACE) was performed using the Takara 3'-Full RACE core kit. First-strand cDNA was synthesized using AMV reverse transcriptase with zebrafish total RNA as the template in conjunction with an oligo dT-3 sites adaptor primer. Afterwards, a polymerase chain reaction (PCR) was carried out using an oligonucleotide (5'-CGTAGTGGCACCACGCTA-ATGC-3') designed based on the sequence determined for the above-mentioned zebrafish TPST cDNA and a three sites adaptor primer as the primer pair with the first-strand cDNA as the template. Amplification conditions were 25 cycles of 30 s at 94 °C, 30 s at 59 °C, and 1 min at 72 °C. The reaction mixture was analyzed by agarose electrophoresis. A discrete PCR product detected was isolated and subcloned into pSTBlue-1 cloning vector and subjected to nucleotide sequencing (Sanger et al. 1977). The nucleotide sequences, as well as the deduced amino acid sequences, of the cDNA were analyzed using BLAST search for sequence homology to known TPSTs.

To amplify the full-length zebrafish TPST cDNA for subcloning into the pcDNA3.1 eukaryotic expression vector, a sense primer (5'-CCCTGACTTTTGCCCCACCTGC-3') corresponding to a region 65 residues upstream from the initiation codon of the open reading frame (see Fig. 1) and an antisense primer (5'-GGTAGTTGTGACTCCG-3') corresponding to a region 116 residues downstream from the stop codon were synthesized. With these two oligonucleotides as primers, PCR in a 50- $\mu$ L reaction mixture was carried out under the action of *Ex Taq* DNA polymerase using zebrafish first-strand cDNA as the template. Amplification conditions were 25 cycles of 40 s at 94 °C, 45 s at 47.5 °C, and 1 min and 20 s at 72 °C. The final reaction mixture was applied onto a 1.2% agarose gel and separated by electrophoresis. The discrete PCR product band, visualized upon ethidium bromide staining, was excised from the gel and the DNA fragment therein was purified by spin filtration. The purified zebrafish TPST cDNA, containing 3' A-overhangs, was subcloned into the 3' T-overhangs-containing pcDNA3.1 eukaryotic expression vector at a cloning site located between *Bst*X I and *Eco*RV restriction sites. To verify its authenticity and direction of cloning, the cDNA insert was subjected to nucleotide sequencing (Sanger et al. 1977).

#### Transient expression of the zebrafish TPST in COS-7 cells

COS-7 cells, routinely maintained in Dulbecco's modified Eagle's medium containing 10% fetal bovine serum, were used as the host cells for the expression of recombinant zebrafish TPST protein. Dishes (100 mm) of COS-7 cells were individually transfected with 10  $\mu$ g of pcDNA3.1 vector only or pcDNA3.1 harboring the zebrafish TPST cDNA using the Lipofectamine 2000 mediated procedure. Transfection was for 18 h at 37 °C. Afterwards, the transfected cells were incubated at 37 °C in Dulbecco's modified Eagle's medium containing 10% fetal bovine serum. At the end of a 48-h incubation, the cells were rinsed twice with phosphate-buffered saline and lysed in an ice-cold lysis buffer containing 20 mM Hepes-NaOH (pH 7.0), 1% Triton X-100, and 10  $\mu$ g aprotinin/mL. The lysate was subjected to centrifugation and the supernatant was used in the TPST assays described below. Lysates of untransfected COS-7 cells or COS-7 cells transfected with pcDNA 3.1 alone were prepared as controls.

#### Preparation of wild-type and mutated PSGL-1 N-terminal peptides as substrates for the zebrafish TPST

For the preparation of cDNA encoding the N-terminal region (ATEYELYLDYDFL) of PSGL-1, two complementary 42-mer oligonucleotides (sense strand 5'-GATCCGCCACCGAAT-ATGAGTACCTAGATTATGATTTCTGG-3' and antisense strand 5'-AATTCCAGGAAATCATAATCTAGGTACTCA-TATTCGGTGGCG-3') were synthesized with, respectively, *Bam*HI and *Eco*RI restriction sites incorporated at the ends. The annealed oligonucleotides were subcloned into the *Bam*HI/*Eco*RI site of pGEX-4T-1 and transformed into *E. coli* XL1-Blue MRF'. For the preparation of cDNA encoding mutated PSGL-1 N-terminal peptides, sense and antisense mutagenic primers with the TAT or TAC codon changed to TTT or TTC were synthesized and similarly processed and subcloned into pGEX-4T-1 and transformed into *E. coli* XL1-Blue MRF'. Figure 2 shows the amino acid sequences of the wild-type and mutated PSGL-1 N-terminal peptides. Competent *E. coli* BL21 cells transformed with pGEX-4T-1 harboring cDNA encoding the wild-type or mutated PSGL-1 N-terminal sequence were grown to  $A_{600nm} = 0.5$  in 100 mL of LB medium supplemented with 100  $\mu$ g ampicillin/mL. Upon induction with 0.1 mM isopropyl  $\beta$ -D-thiogalactopyranoside overnight at 25 °C, the cells were collected by centrifugation at 1000g for 10 min at 4 °C and homogenized in 15 mL of an ice-cold lysis buffer (containing 20 mM Tris-HCl (pH 8.0), 150 mM NaCl, and 1 mM EDTA) using an Aminco French press. The crude homogenate thus prepared was subjected to centrifugation at 10 000g for 20 min at 4 °C. The collected supernatant that includes the fusion protein was fractionated using 0.5 mL of glutathione-Sepharose for 1 h, and the Sepharose was washed three times with lysis buffer and twice with thrombin digestion buffer (containing 20 mM Tris-HCl (pH 8.0), 150 mM NaCl, and 2.5 mM CaCl<sub>2</sub>). Afterwards, the fusion protein bound to the Sepharose was treated with 1 mL of a thrombin digestion buffer containing 5 units thrombin/mL. After a 2-h incubation at room temperature, the preparation was subjected to centrifugation. The recombinant peptide released into the supernatant was purified using a Waters Sep-Pak Plus C18 cartridge according to the manufacturer's instructions. Purified peptide was used as substrate in the enzymatic assay.

#### Enzymatic assay

The standard assay for TPST activity was carried out in a reaction mixture (25  $\mu$ L final volume) consisting of 50 mM MES (pH 5.75), 50 mM NaF, 20 mM MnCl<sub>2</sub>, 0.1% Triton X-100, 40  $\mu$ M peptide substrate, and 15  $\mu$ M PAP[<sup>35</sup>S]. The reaction was started by the addition of the enzyme, allowed to proceed for 30 min at 28 °C, and terminated by heating at 100 °C for 3 min. The precipitates formed were cleared by centrifugation, and the supernatants collected were analyzed for [<sup>35</sup>S]sulfated product using a previously developed thin-layer chromatography procedure (Liu and Lipmann 1984) with *n*-butanol - isopropanol - 88% formic acid - water (3:1:1:1 by volume) as the solvent system. To examine the pH dependence, different buffers (50 mM MES at pH 5.5, 5.75, or 6.0; MOPS at pH 6.25, 6.5, or 7.0) instead of 50 mM MES (pH 5.75) were used in the reactions. For the kinetic studies on the sulfation of peptide substrates, varying

**Fig. 1.** Nucleotide and deduced amino acid sequences of the zebrafish TPST cDNA. Nucleotides are numbered in the 5' to 3' direction. The translation stop codon is indicated by an asterisk. The putative transmembrane segment and the residues involved in the binding of the 5'-phosphosulfate group and the 3'-phosphate group of PAPS are underlined.

```

1   CTGTTGGCCTACTGGAGCGATTATTTACATACAGACAGAATTCACGAGAAATGTGCTCGC
61  GGTGTCAGATTCTGAGTCTCCTGAGAGGTTTATAGCTCCATTTACAGATCTGTTCTGCTCT
121 GCATTTTCATCACGCGATTCTGGAGAATGTAGAATTCGAGGAGTTGTGCTTTCTGTACGGC
181 ACAATGGTGCTGACAGAGAATATATGGCTTGATTTACCATGTCTGAGTTTATGAAGCCT
241 CAAGCAAAGCCCAGAAGACCTGCTGACTGAGGAAAAAAATCTCCCATCCCTTCCCTCCCC
301 CCAAAAAAATTTTACACTTCGCCTTTTACATTCCCTTACCCTGACGCTTTTGGAAACA
361 TTCACTCCGCTTTCCCTTCCCTGACTTTTGCCCCACCTGCAAAATCTGTTAACTCA
421 TTTGTTATTTTCCCTTACAAGCAAGTACTTCCCTAACTGTAAGTCATGAGAAAAACA
1   M R K Q
481 ACGTGAATGTTCTCCTAGTCTGTGGGGTCATCAGCTCCATCACAGTATTTTATCTTGGC
5   T C N V L L V C G V I S S I T V F Y L G
541 CTCAGTACGATAGAGTGTCCGAATGCCCGTTCTCGTGCATCACAGCATGGGTGGGTGTA
25  L S T I E C P N A R S R A S Q H G W V V
601 AACCTACAGCTGGCAGAAACCTTAGTGACCCATTACAGCTCCCTGAGGAGTACAATGAG
45  N L H A G R N L S D P L Q L P E E Y N E
661 GAACTCCTCTCATTTTGTGGCGGAGTCCCTCGTAGTGGCACCACGCTAATGCGGGCT
65  E T P L I F V G G V P R S G T T L M R A
721 ATGTTAGATGCTCACCCATCGTGGCGGAGAGACCCGGTTCATCCCTCGGTG
85  M L D A H P I V R C G E E T R V I P R L
781 TTAGCCATGCAGCAACCTGGAGTCACTCGGCACGAGAGCGGGTCCGTCTGGATGAGGCT
105 L A M Q A T W S H S A R E R V R L D E A
841 GGTGTCAGTATGATGTTTTGGACTCTGCTGTACGTGCGTTTCTTTGGAGATCATAGTA
125 G V T D D V L D S A V R A F L L E I I V
901 GGGCATGGGAGCCAGCGCCGAGGCTCTGCAACAAGGACCCATTTGCTCTGAAGTCCATG
145 G H G E P A P R L C N K D P F A L K S M
961 TCCTACCTCTCAAACTCTTCCCAAAGGCGAAGTTTATTCTCATGCTTCGTGATGGCAGG
165 S Y L S K L F P K A K F I L M L R D G R
1021 GCCACCGTTCACTCGATGATCTCCCGAAGGTTACTATTACTGGGTTTGACTTGACAAGT
185 A T V H S M I S R K V T I T G F D L T S
1081 TACCGGGATTGTTGGTAAAGTGAACCGGGCGGTGAAGTGTACGACCAGTGCCTG
205 Y R D C L V K W N R A V E V M Y D Q C L
1141 GCTGCAGTGGATGGCAACTGTCTGCCTGTCCATTATGAGCAGCTTGTGCTGCATCCGAG
225 A A V D G N C L P V H Y E Q L V L H P E
1201 CGGGTATGCGAGGCTCCTTCAGTCTTGGATCTGCCATGGGACACTGTGTGCTGCAT
245 R V M R R L L Q F L D L P W D T A V L H
1261 CATGAACAGCTAATGGGAAAGTCCGAGGAGTTTCGCTGTCAAAGGTGGAACCTGTCAACA
265 H E Q L I G K V G G V S L S K V E L S T
1321 GATCAAGTAGTGAAGCCAGTGAATACAGAGGCTCTGTCTAAATGGGTGGCAAGATTCT
285 D Q V V K P V N T E A L S K W V G K I P
1381 GCTGATGTAGTGAAGGACATGCCAGCCTTGCCCCATGCTGAGTGCCTTGGGTTATGAC
305 A D V V K D M P S L A P M L S R L G Y D
1441 CCTCTGGCCAACCCACCAACTACAACAAGCCTGATCTTATATCTGAACAACAAAAA
325 P L A N P P N Y N K P D L L Y L N N T K
1501 ATAGTAAGGCCGATGTAAGTGAAGGTCATCTCTGTGGATACAACCTGTGAATATTTG
345 I V R P M *
1561 GAAGCAATAACAATGTGCATTTTTTGTAGCACGCAAAGTCTCTACTTGAATTAAGAACA
1621 TGTATTGACAAAACGGAGTCACAACTACCATTGTTTACATACAAAGTGAAGTCTAGTTT
1681 ATGTTGATTTTAAGTCATTTCTCAATTAAGTGCATTTTCAGATGTTTGTGCTGTTTGTACC
1741 GATGTTATGATGTTGTTATTAATGTGTCAGTCCGATGACTATAAGGTGGCTTTTGTCTTGTG
1801 GTTCTATTTGTATGCATGCATGTCACAGTAAATGTGCTCCTAAAGAGTGATATCACACAG
1861 GAACTATGTTTGTGAAAGAGGAAGTCAAAGTTGCACAGAAATATCACTATTTTTCTCT
1921 GACCGCAGGAGAACTAAGTAAGTTCCATCTAC

```

concentrations of the substrate compounds and 50 mM MES at pH 5.75 were used.

#### Miscellaneous methods

PAP[<sup>35</sup>S] (15 Ci/mmol (1 Ci = 37 GBq)) was synthesized from ATP and [<sup>35</sup>S]sulfate using recombinant human

bifunctional ATP sulfurylase/adenosine 5'-phosphosulfate kinase as described previously (Yanagisawa et al. 1998). In vitro transcription/translation of the zebrafish TPST was performed using the TNT coupled reticulocyte lysate system according to the manufacturer's instructions. Sodium dodecyl sulfate – polyacrylamide gel electrophoresis (SDS–

**Fig. 2.** Amino acid sequences of the wild-type (wt) and mutated PSGL-1 N-terminal peptides. The three tyrosine residues and the mutated phenylalanine residues are boldfaced.

PSGL-1 wt	ATEYEYLDYDFL
PSGL-1 1, 2 Y	ATEYEYLD <b>DF</b> DFL
PSGL-1 1, 3 Y	ATEY <b>EF</b> LDYDFL
PSGL-1 2, 3 Y	ATE <b>FE</b> YLDYDFL
PSGL-1 1Y	ATEY <b>EF</b> LD <b>DF</b> DFL
PSGL-1 2Y	ATE <b>FE</b> YLD <b>DF</b> DFL
PSGL-1 3Y	ATE <b>FE</b> <b>EF</b> LDYDFL
PSGL-1 F	ATE <b>FE</b> <b>EF</b> LD <b>DF</b> DFL

PAGE) was performed on 12% polyacrylamide gels using the method of Laemmli (1970). Protein determination was based on the method of Bradford (1976) with bovine serum albumin as the standard.

## Results and discussion

Although considerable progress has been made in recent years in the study of the TPST enzymes, some fundamental questions concerning their ontogeny, regulation, and physiological involvement remain unanswered. The present study was prompted by an attempt to develop a zebrafish model to address these important issues. As a first step toward achieving this goal, we have decided to clone, express, and characterize the TPST enzymes present in zebrafish.

### Molecular cloning of the zebrafish TPST

The zebrafish TPST cDNA cloned by reverse transcription – PCR in conjunction with 3'-RACE was subjected to nucleotide sequencing, and the nucleotide sequence obtained was deposited at the GenBank database under accession No. AY263386. Figure 1 shows the nucleotide and deduced amino acid sequences of this zebrafish enzyme. The open reading frame encompasses 1047 nucleotides and encodes a 349 amino acid polypeptide with a calculated molecular mass of 39 030 Da (without considering posttranslational modifications). Sequence analysis based on BLAST search revealed that the deduced amino acid sequence of the zebrafish TPST displayed, respectively, 66% and 60% identity to those of human and mouse TPST-1 (Ouyang et al. 1998) and TPST-2 (Beisswanger et al. 1998; Ouyang and Moore 1998). It should be pointed out that although the cloned zebrafish TPST showed higher percent homology to human or mouse TPST-1, it may actually correspond to TPST-2. We have recently cloned and sequenced another zebrafish TPST (Liu et al., unpublished data), and sequence analysis revealed it to resemble more closely human or mouse TPST-1. Moreover, it is worthwhile mentioning that TPSTs have also been found to be present in lower animals such as *Caenorhabditis elegans* and *Drosophila melanogaster* but not in yeast (Moore 2003). The zebrafish TPST cloned in the present study, while displaying higher percent identity in amino acid sequence to human and mouse TPSTs, also exhibited significant homology to TPSTs from these latter animals. Hydropathy analysis revealed that, similar to human or

mouse TPSTs, the zebrafish TPST cloned in the present study contains a putative transmembrane segment (as underlined) located near the N terminus. The zebrafish TPST therefore appears to be a type II transmembrane protein with a short N-terminal cytoplasmic tail with the bulk of this putative Golgi protein being present in the Golgi lumen. Previous studies employing X-ray crystallography in conjunction with sequence alignment have revealed that all cytosolic and Golgi sulfotransferases contain two conserved sequence elements (Negishi et al. 2001). These two sequence elements, designated the 5'-phosphosulfate binding motif and the 3'-phosphate binding motif, are responsible for binding to, respectively, the 5'-phosphosulfate group and the 3'-phosphate group of PAPS, a cosubstrate for sulfotransferase-catalyzed sulfation reactions (Lipmann 1958). Examination of the amino acid sequence of the zebrafish TPST also revealed residues 76–80 (RSGTT, underlined) that correspond to the 5'-phosphosulfate binding motif. And, similar to human and mouse TPSTs previously reported, the zebrafish TPST also contains two conserved residues (Arg<sup>181</sup> and Ser<sup>189</sup>, boldfaced and underlined) proposed to be involved in binding the 3'-phosphate group of PAPS (Ouyang et al. 1998; Beisswanger et al. 1998; Ouyang and Moore 1998).

### Expression of recombinant zebrafish TPST in COS-7 cells

The full-length zebrafish TPST cDNA packaged in pcDNA3.1, a eukaryotic expression vector, was used. To ensure that the cDNA can be used as a template for transcription followed by translation for the synthesis of TPST protein, an in vitro transcription/translation experiment was first carried out. As shown in Fig. 3, a 39-kDa protein band was detected in the reaction mixture upon SDS-PAGE, indicating the production of the TPST protein. pcDNA3.1 harboring the zebrafish TPST cDNA was then used to transfect COS-7 cells for the expression of the recombinant enzyme. Lysates of untransfected cells and cells transfected with pcDNA3.1 vector alone or pcDNA3.1 harboring the zebrafish TPST cDNA were then assayed for TPST activity using the wild-type PSGL-1 N-terminal peptide as substrate. As shown in Table 1, lysates of untransfected cells and cells transfected with pcDNA3.1 alone showed approximately the same level of TPST activity. This basal TPST activity is due to the endogenous TPST of COS-7 cells. In contrast, there was a six-fold increase in TPST activity in lysate of COS-7 cells transfected with pcDNA3.1 harboring the zebrafish TPST cDNA, indicating clearly the production of functionally active zebrafish TPST.


### Characterization of the recombinant zebrafish TPST

The wild-type and a series of mutated PSGL-1 N-terminal peptides (see Fig. 2) were prepared for the characterization of the recombinant zebrafish TPST.

#### Temperature dependence, pH optimum, and divalent cation requirement

We first examined the temperature dependence of the activity of the zebrafish TPST using the wild-type PSGL-1 N-terminal peptide as substrate. As shown in Fig. 4, the zebrafish TPST exhibited approximately the same level of activity at 28 and 37 °C. At 21 °C, there was a 50% decrease in the sulfating activity of the enzyme, and only a very low level of activity

**Fig. 3.** Production of the zebrafish TPST protein by in vitro transcription/translation. The figure shows the autoradiograph taken from the dried polyacrylamide gel used for the SDS-PAGE of the reaction mixtures. Samples analyzed: lane 1, in vitro transcription/translation using pcDNA3.1 vector only as the template; lane 2, in vitro transcription/translation using pcDNA3.1 harboring the zebrafish TPST cDNA as the template.




**Table 1.** Expression of recombinant zebrafish TPST in COS-7 cells.

	Specific activity (pmol·min <sup>-1</sup> ·mg <sup>-1</sup> )	Relative activity (fold)
None	2.03±0.67	1.0
pcDNA3.1	2.07±0.38	1.0
pcDNA3.1-TPST-2 cDNA	11.7±0.20	5.8


Note: Values are means ± SD derived from three determinations.

was detected at 44 °C. As indicated in Table 1, the untransfected COS-7 cell lysate contained a low level of TPST activity due to the endogenous enzyme. It is likely that this endogenous TPST activity may account for the bulk of the activities detected at 37 and 44 °C. It is worthwhile mentioning that although zebrafish are subjected to fluctuation in body temperature in their natural habitat, they have been shown to be best maintained in aquaria heated to 28 °C (Westerfield 2000). Taking into consideration their optimal temperature of growth and the results from the temperature dependence study, we therefore decided to characterize the other enzymatic properties of the zebrafish TPST at 28 °C. Another important property is with regard to the pH optimum of the enzyme. In a pH dependence experiment, the zebrafish TPST displayed a pH optimum of 5.75 (Fig. 5). Previous studies have revealed TPSTs as Golgi enzymes with the catalytic domain being located in the Golgi lumen (Ouyang et al. 1998; Beisswanger et al. 1998; Ouyang and Moore 1998). The acidic pH optimum determined for the zebrafish TPST, therefore, is compatible with the acidic environment in the

**Fig. 4.** Temperature dependence of the sulfating activity of the zebrafish TPST with the wild-type PSGL-1 N-terminal peptide as substrate. The enzymatic assays were carried out at designated temperatures under standard assay conditions as described in Materials and methods. The data represent calculated mean values derived from three experiments.




**Fig. 5.** pH dependency of the sulfating activity of the zebrafish TPST with the wild-type PSGL-1 N-terminal peptide as substrate. The enzymatic assays were carried out under standard assay conditions, as described in Materials and methods, using different buffer systems. The data represent calculated mean values derived from three experiments.



Golgi lumen (Mellman et al. 1986). Previous studies have revealed stimulatory effects of manganese on the activity of some mammalian TPSTs (Lin et al. 1994). We were interested in examining whether manganese is also capable of stimulating the activity of the zebrafish TPST. In a concentration dependence experiment (Fig. 6), it was found that in the presence of 20 mM MnCl<sub>2</sub>, the sulfating activity of the zebrafish TPST was more than 2.5 times that determined in the absence of MnCl<sub>2</sub>. At concentrations higher than 50 mM,

Fig. 6. Manganese dependence of the sulfating activity of the zebrafish TPST with the wild-type PSGL-1 N-terminal peptide as substrate. The enzymatic assays were carried out in the presence of different concentrations of  $MnCl_2$  under standard assay conditions as described in Materials and methods. The data represent calculated mean values derived from three experiments.




however, manganese became inhibitory. The mechanism underlying the stimulatory effects of manganese remains to be clarified. We further tested nine other divalent cations for their stimulatory/inhibitory effects on the activity of the zebrafish TPST. As shown in Fig. 7, at a 10 mM concentration,  $Co^{2+}$  showed a significant stimulatory effect, while  $Ca^{2+}$ ,  $Pb^{2+}$ , and  $Cd^{2+}$  exerted some inhibitory effects. In contrast, the other four divalent cations,  $Fe^{2+}$ ,  $Cu^{2+}$ ,  $Zn^{2+}$ , and  $Hg^{2+}$ , inhibited completely the sulfating activity of the zebrafish TPST.

#### Substrate specificity

Previous studies have led to a consensus that a key feature of the tyrosine sulfation site is the presence of acidic residues on the N-terminal side of the sulfatable tyrosine (Huttner 1984; Hortin et al. 1986). As shown in Fig. 2, the wild-type PSGL-1 N-terminal peptide contains three tyrosine residues and all of them fulfill such a requirement for sulfation. We were interested in finding out whether the three tyrosine residues of the wild-type PSGL-1 N-terminal peptide are equally or differentially sulfated by the zebrafish TPST. Mutated peptides with one or two of the three tyrosine residues being replaced by phenylalanine were prepared and used as substrates. As shown in Fig. 8, the zebrafish TPST appeared to be most active toward mutated peptides that retained the C-terminal tyrosine residue. In contrast, mutated peptides where this C-terminal tyrosine residue was mutated served as very poor substrates for the zebrafish enzyme even when the N-terminal and (or) middle tyrosine residue(s) were intact. These results indicated clearly the specificity of the zebrafish TPST for the C-terminal tyrosine

Fig. 7. Effects of divalent metal cations on the sulfating activity of the zebrafish TPST with the wild-type PSGL-1 N-terminal peptide as substrate. The sulfating activity of the zebrafish TPST was assayed in the presence of different divalent metal cations or NaCl (as a control for the counter ion  $Cl^-$ ) under standard conditions as described in Materials and methods. The concentration of the divalent metal cations tested was 10 mM, and the concentration of NaCl tested was 20 mM.



residue in the wild-type PSGL-1 N-terminal peptide. Kinetic constants for the wild-type and the three mutated PSGL-1 N-terminal peptide that have the C-terminal tyrosine residue intact were determined. Table 2 shows the differential  $K_m$




**Table 2.** Kinetic constants of zebrafish TPST with wild-type and mutant PSGL1 N-terminal peptides as substrates.

Substrate	$K_m$ ( $\mu\text{M}$ )	$V_{max}$ ( $\text{pmol}\cdot\text{min}^{-1}\cdot\text{mg}^{-1}$ )	$V_{max}/K_m$ ( $\text{pmol}\cdot\text{min}^{-1}\cdot\text{mg}^{-1}/\mu\text{M}$ )
PSGL-1 (wild type)	35.9	4.1	0.11
3Y	31.5	42.2	1.34
1,3Y	5.2	17.5	3.37
2,3Y	8.5	10.0	1.18

Note: Values are means  $\pm$  SD derived from three determinations.

**Fig. 8.** Substrate specificity of the zebrafish TPST with the wild-type and mutated PSGL-1 N-terminal peptides as substrates. The enzymatic assays were carried out under standard assay conditions as described in Experimental procedures. The data represent calculated mean values derived from three experiments. The solid bars correspond to the activities detected using the zebrafish TPST-expressing COS-7 cell lysate, and the open bars correspond to the activities detected using the control COS-7 cell lysate.



and  $V_{max}$  values of these peptide substrates. Interestingly, in the absence of both the N-terminal and the middle tyrosine residues, the C-terminal tyrosine residue was sulfated 10 times faster than in the presence of those two tyrosine residues.

To summarize, we have cloned a zebrafish TPST and expressed and characterized the recombinant enzyme. The recombinant zebrafish TPST exhibited some properties, including acidic pH optimum and stimulation by manganese, that are similar to those previously determined for mammalian TPSTs. Further studies concerning the ontogeny, regulation, and physiological involvement of the zebrafish TPST are now in progress.

### Acknowledgments

This work was supported in part by a grant from the American Heart Association (Texas Affiliate) (M.-C.L.), a grant-in-aid for Encouragement of Young Scientists (No. 11760072) from Munbusho, Japan (Y.S.), and an award from the Naito Foundation (M.S.).

### References

- Bauerle, P.A., and Huttner, W.B. 1987. Tyrosine sulfation is a trans Golgi-specific protein modification. *J. Cell Biol.* **105**: 2655–2664.
- Beisswanger, R., Corbeil, D., Vannier, C., Thiele, C., Dohrmann, U., Kellner, R., Ashman, K., Niehrs, C., and Huttner, W.B. 1998. Existence of distinct tyrosylprotein sulfotransferase genes: molecular characterization of tyrosylprotein sulfotransferase-2. *Proc. Natl. Acad. Sci. U.S.A.* **95**: 11 134 – 11 139.
- Bettelheim, F.R. 1954. Tyrosine-*O*-sulfate in a peptide from fibrinogen. *J. Am. Chem. Soc.* **76**: 2838–2839.
- Bradford, M.M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**: 248–254.
- Brand, S.J., Andersen, B.N., and Rehfeld, J.F. 1984. Complete tyrosine-*O*-sulphation of gastrin in neonatal rat pancreas. *Nature (Lond.)*, **309**: 456–458.
- Briggs, J.P. 2002. The zebrafish: a new model organism for integrative physiology. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **282**: R3–R9.
- Farzan, M., Mirzabekov, T., Kolchinsky, P., Wyatt, R., Cayabyab, M., Gerard, N.P., Gerard, C., Sodroski, J., and Choe, H. 1999. Tyrosine sulfation of the amino terminus of CCR5 facilitates HIV-1 entry. *Cell*, **96**: 667–676.
- Hille, A., Rosa, P., and Huttner, W.B. 1984. Tyrosine sulfation: a post-translational modification of proteins destined for secretion? *FEBS Lett.* **117**: 129–134.
- Hortin, G., Folz, R., Gordon, J.I., and Strauss, A.W. 1986. Characterization of sites of tyrosine sulfation in proteins and criteria for predicting their occurrence. *Biochem. Biophys. Res. Commun.* **141**: 326–333.
- Huttner, W.B. 1982. Sulphation of tyrosine residues — a widespread modification of proteins. *Nature (Lond.)*, **299**: 273–276.
- Huttner, W.B. 1984. Determination and occurrence of tyrosine *O*-sulfate in proteins. *Methods Enzymol.* **107**: 200–223.
- Jensen, S.L., Holst, J.J., Nielsen, O.V., and Rehfeld, J.F. 1981. Effect of sulfation of CCK-8 on its stimulation of the endocrine and exocrine secretion from the isolated perfused porcine pancreas. *Digestion*, **22**: 305–309.
- Kashinathan, C., Sundaram, P., Slomiany, B.L., and Slomiany, A. 1992. Identification of tyrosylprotein sulfotransferase in rat gastric mucosa. *Enzyme*, **46**: 179–187.
- Kehoe, J.W., and Bertozzi, C.R. 2000. Tyrosine sulfation: a modulator of extracellular protein-protein interactions. *Chem. Biol.* **7**: R57–R61.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (Lond.)*, **227**: 680–685.
- Lee, R.W.H., and Huttner, W.B. 1983. Tyrosine-*O*-sulfated proteins of PC12 pheochromocytoma cells and their sulfation by a



- tyrosylprotein sulfotransferase. *J. Biol. Chem.* **258**: 11 326 – 11 334.
- Lee, R.W.H., and Huttner, W.B. 1985. (Glu62, Ala30, Tyr8)*n* serves as high-affinity substrate for tyrosylprotein sulfotransferase: a Golgi enzyme. *Proc. Natl. Acad. Sci. U.S.A.* **82**: 6143–6147.
- Lin, W.-H., and Roth, J.A. 1990. Characterization of a tyrosylprotein sulfotransferase in human liver. *Biochem. Pharmacol.* **40**: 629–635.
- Lin, W.H., Marcucci, K., and Roth, J.A. 1994. Effect of manganese on tyrosylprotein sulfotransferase activity in PC12 cells. *Biochem. Pharmacol.* **47**: 1575–1580.
- Lipmann, F. 1958. Biological sulfate activation and transfer. *Science (Wash., D.C.)*, **128**: 575–580.
- Liu, M.-C., and Lipmann, F. 1984. Decrease of tyrosine-*O*-sulfate-containing proteins found in rat fibroblasts infected with Rous sarcoma virus or Fujinami sarcoma virus. *Proc. Natl. Acad. Sci. U.S.A.* **81**: 3695–3698.
- Mellman, I., Fuchs, R., and Helenius, A. 1986. Acidification of the endocytic and exocytic pathways. *Annu. Rev. Biochem.* **55**: 663–700.
- Moore, K.L. 2003. The biology and enzymology of protein tyrosine *O*-sulfation. *J. Biol. Chem.* **278**: 24 243 – 24 246.
- Negishi, M., Pedersen, L.G., Petrotchenko, E., Shevtsov, S., Gorokhov, A., Kakuta, Y., and Pedersen, L.C. 2001. Structure and function of sulfotransferases. *Arch. Biochem. Biophys.* **390**: 149–157.
- Ouyang, Y.B., and Moore, K.L. 1998. Molecular cloning and expression of human and mouse tyrosylprotein sulfotransferase-2 and a tyrosylprotein sulfotransferase homologue in *Caenorhabditis elegans*. *J. Biol. Chem.* **273**: 24 770 – 24 774.
- Ouyang, Y., Lane, W.S., and Moore, K.L. 1998. Tyrosylprotein sulfotransferase: purification and molecular cloning of an enzyme that catalyzes tyrosine *O*-sulfation, a common posttranslational modification of eukaryotic proteins. *Proc. Natl. Acad. Sci. U.S.A.* **95**: 2896–2901.
- Ouyang, Y.B., Crawley, J.T., Aston, C.E., and Moore, K.L. 2002. Reduced body weight and increased postimplantation fetal death in tyrosylprotein sulfotransferase-1-deficient mice. *J. Biol. Chem.* **277**: 23 781 – 23 787.
- Pauwels, S., Dockray, G.J., and Walker, R. 1987. Comparison of the metabolism of sulfated and unsulfated heptadecapeptide gastrin in humans. *Gastroenterology*, **92**: 1220–1225.
- Pouyani, T., and Seed, B. 1995. PSGL-1 recognition of P-selectin is controlled by a tyrosine sulfation consensus at the PSGL-1 amino terminus. *Cell*, **83**: 333–343.
- Ramaprasad, P., and Kashinathan, C. 1998. Isolation of tyrosylprotein sulfotransferase from rat liver. *Gen. Pharmacol.* **30**: 555–559.
- Rens-Domiano, S., and Roth, J.A. 1989. Characterization of tyrosylprotein sulfotransferase from rat liver and other tissues. *J. Biol. Chem.* **264**: 899–905.
- Rens-Domiano, S., Hortin, G.L., and Roth, J.A. 1989. Sulfation of *tert*-butoxycarbonylcholecystokinin and other peptides by rat liver tyrosylprotein sulfotransferase. *Mol. Pharmacol.* **36**: 647–653.
- Rosa, P., Hille, A., Lee, R.W.H., Zanini, A., De Camilli, P., and Huttner, W.B. 1985. Secretogranins I and II: two tyrosine-sulfated secretory proteins common to a variety of cells secreting peptides by the regulated pathway. *J. Cell Biol.* **101**: 1999–2011.
- Sako, D., Comess, K.M., Barone, K.M., Camphausen, R.T., Cumming, D.A., and Shaw, G.D. 1995. A sulfated peptide segment at the amino terminus of PSGL-1 is critical for P-selectin binding. *Cell*, **83**: 323–331.
- Sane, D.C., and Baker, M.S. 1993. Human platelets possess tyrosylprotein sulfotransferase (TPST) activity. *Thromb. Haemostasis*, **69**: 272–275.
- Sanger, F., Nicklen, S., and Coulson, A.R. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. U.S.A.* **74**: 5463–5467.
- Suiko, M., Fernando, P.H., Sakakibara, Y., Kudo, H., Nakamura, T., and Liu, M.-C. 1997. Characterization of bovine heart sulfotransferase catalyzing the sulfation of tyrosine-containing peptides. *J. Nutr. Sci. Vitaminol.* **43**: 485–490.
- Sundaram, P., Slomiany, A., Slomiany, B.L., and Kashinathan, C. 1992. Tyrosylprotein sulfotransferase in rat submandibular salivary glands. *Int. J. Biochem.* **24**: 663–667.
- Tuboi, S., Taniguchi, N., and Katunuma, N. 1992. The post-translational modification of proteins: roles in molecular and cellular biology. CRC Press, Boca Raton, Fla.
- Vargas, F., Frerot, O., Tuong, M.D., and Schwartz, J.C. 1985. Characterization of a tyrosine sulfotransferase in rat brain using cholecystokinin derivatives as acceptors. *Biochemistry*, **24**: 5938–5943.
- Ward, A.C., and Lieschke, G.J. 2002. The zebrafish as a model system for human disease. *Front. Biosci.* **7**: d827–d833.
- Westerfield, M. 2000. The zebrafish book. University of Oregon Press, Eugene, Oreg.
- Wilkins, P.P., Moore, K.L., McEver, R.P., and Cummings, R.D. 1995. Tyrosine sulfation of P-selectin glycoprotein ligand-1 is required for high affinity binding to P-selectin. *J. Biol. Chem.* **270**: 22 677 – 22 680.
- William, S., Ramaprasad, P., and Kashinathan, C. 1997. Purification of tyrosylprotein sulfotransferase from rat submandibular salivary glands. *Arch. Biochem. Biophys.* **338**: 90–96.
- Yanagisawa, K., Sakakibara, Y., Suiko, M., Takami, Y., Nakayama, T., Nakajima, H., Takayanagi, K., Natori, Y., and Liu, M.-C. 1998. cDNA cloning, expression, and characterization of the human bifunctional ATP sulfurylase/adenosine 5'-phosphosulfate kinase enzyme. *Biosci. Biotechnol. Biochem.* **62**: 1037–1040.
- Young, W.F., Jr. 1990. Human liver tyrosylsulfotransferase. *Gastroenterology*, **99**: 1072–1078.