

Protein and mRNA levels of IgM H- and L-chains artificially and excessively accumulated in HDAC2-deficient DT40 mutants are dramatically reduced through various generations during continuous cultivation**

Masami Nakayama[#] and Tatsuo Nakayama^{#, \$, *}

[#]Section of Biochemistry and Molecular Biology, Department of Medical Sciences, Faculty of Medicine, ^{\$}Department of Life Science, Frontier Science Research Center, University of Miyazaki, 5200, Kihara, Kiyotake, Miyazaki, 889-1692, Japan

***Corresponding author: tnakayam@med.miyazaki-u.ac.jp (T. Nakayama)**

****The article is the revision of the original paper, which was published in 2017 and is available from following URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10145170>**

****The article is also the modified version of Chapter 2 of the revised monograph (the original monograph was published in 2015 and is available from following URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10123995>) as follows: Nakayama, M. and Nakayama, T.: Protein and mRNA levels of IgM H- and L-chains artificially accumulated in HDAC2-deficient DT40 mutants are dramatically reduced through various generations during continuous cultivations. In: Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations, Nakayama, T. and Nakayama, M. (Eds.), pp. 11-44, 2018. The revised monograph is available from following URL:**

<http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10169365>

Abstract

We analyzed histone deacetylase-2 (HDAC2)-deficient DT40 mutant cells, HDAC2(-/-), which were cultivated for varying periods, by two-dimensional polyacrylamide gel electrophoresis (2D-PAGE), Western blotting, reverse transcription-polymerase chain reaction (RT-PCR) and immuno-electron microscopy. Protein and mRNA levels of IgM H- and L-chains were increased at the early stage of cultivation and thereafter until the later stage reduced to comparable levels as in DT40 cells. The mRNA level of PCAF was increased from the early to later stages, and those of HDAC7 and HDAC9 changed during cultivation. The mRNA level of EBF1 was almost completely decreased at the early stage and thereafter remained unchanged, but that of E2A was slightly increased at the early stage and thereafter remained unchanged. The mRNA level of Pax5, which down-regulates gene expressions of IgM H- and L-chains, was reduced at the early stage and thereafter increased through the middle to later stages in anti-parallel with those of these two immunoglobulin proteins. Bulk acetylation levels of Lys-9, Lys-14, Lys-18, Lys-23 and Lys-27 residues of histone H3 (K9/H3, K14/H3, K18/H3, K23/H3 and K27/H3) were increased during cultivation. Chromatin immuno-precipitation (ChIP) assay suggested that acetylation levels of K9/H3 within some chromatin segments surrounding proximal ~2.0 kb 5'-upstream region of the Pax5 gene were decreased at the early stage in HDAC2(-/-) and thereafter at the later stage increased to comparable levels as in DT40 cells.

These results revealed not only that the HDAC2-deficiency induces dramatic increases in mRNA and protein levels of IgM H- and L-chains, and thereafter these accumulated mRNAs and proteins are gradually reduced during cultivation, but also that alterations in gene expressions of IgM H- and L-chains may be coupled with changes in those of Pax5, EBF1, PCAF, HDAC7 and HDAC9. In addition, the changing pattern in acetylation levels of K9/H3 within proximal 5'-upstream chromatin region of the Pax5 gene in HDAC2(-/-) qualitatively agrees with that in the transcription level of the gene.

Keywords

HDAC2-deficient DT40 mutants, continuous cultivation, decreases in IgM H/L chain protein/mRNA levels, changes in mRNA levels of transcription factors, gene targeting techniques, ChIP assay, changes in acetylation levels of K9/H3 surrounding Pax5 5'-upstream chromatin region

Introduction

The modulation of chromatin topology with epigenetic modifications is one of the most important ways for expression of normal cell functions in eukaryotes. Alterations in the chromatin structure are preferentially involved in controls of gene expression (transcription) and DNA replication, repair and recombination, etc. [1-3]. Naturally, such alterations in the chromatin structure are concerned in regulations of lymphocyte development and differentiation [4-7]. Moreover, the regulation of the

lymphocyte development requires a number of transcription factors, including Ikaros, E2A, PU.1, GATA-3, Pax5, EBF and others, and the transcriptional regulation of IgM H-chain requires USF, TFEB, Ig/EBP, NF-IL6, OCA-B and others as promoter binding proteins and Ig/EBP, NF-IL6, YY-1, E2A, PU.1 and others as intron enhancer binding proteins [8-13]. On the other hand, of various epigenetic modifications of the chromatin structure with acetyl, methyl, phosphate, ubiquitin, ADP ribose groups and others, acetylation and deacetylation of specific Lys residues (K) of core histones (H2A, H2B, H3 and H4) may be major ones [14-21]. They are cooperatively controlled in combination of specific members of chromatin-modifying enzymes, i.e., histone acetyltransferase(s) (HATs) and histone deacetylase(s) (HDACs) [22, 23]. To clarify *in vivo* roles of individual members of HATs and HDACs in expressions of cell functions, we have systematically generated a number of homozygous (or conditional) chicken DT40 mutant cell lines lacking appropriate genes [24-33] by gene targeting techniques [34-39]. Our previous analysis of HDAC2-deficient DT40 mutant cells, HDAC2(-/-), revealed that HDAC2 controls the amount of IgM H-chain at the steps of its gene expression plus pre-mRNA alternative processing [24] and down-regulates IgM L-chain gene promoter activity [27]. Moreover, the HDAC2-deficiency decreases gene expressions of HDAC7, Pax5, Aiolos, Ikaros plus EBF1, increases those of HDAC4, HDAC9, PCAF plus E2A, and changes bulk acetylation levels of several specific Lys residues of core histones H2A, H2B, H3 and H4 [30].

Subsequently, to know individual roles of these changed transcription factors in regulations of IgM H- and L-chain gene expressions, we generated homozygous DT40 mutant cells; EBF1(-/-), Aiolos(-/-), E2A(-/-) and Helios(-/-), devoid of two alleles of EBF1, Aiolos, E2A and Helios genes, respectively [30]. In addition, we generated Pax5-deficient mutant cells, Pax5(-), devoid of the Pax5 gene existing on Z sex chromosome that is monosomy in chickens (USCS Genome Browser data base) [40], but could generate only Ikaros-down mutant cells, Ikaros(-/+), devoid of two alleles of the Ikaros gene existing on chromosome 2 that is trisomy, probably because it is very important or essential for DT40 cells (data will be shown elsewhere). Analyses of these resultant mutants revealed that Pax5, EBF1, Aiolos plus Ikaros down-regulate IgM H- and L-chain gene expressions, and E2A up-regulates gene expressions of these two immunoglobulin proteins [30]. Thus, these results indicated that HDAC2 indirectly controls gene expressions of IgM H- and L-chains through opposite regulations of gene expressions of Pax5, EBF1, Aiolos plus Ikaros, and E2A [30, 31].

In this study, we analyzed HDAC2(-/-) at different cultivation periods by 2D-PAGE, Western blotting, immuno-electron microscopy and RT-PCR. During cultivation of HDAC2(-/-), protein and mRNA levels of IgM H- and L-chains were dramatically and considerably elevated at the early stage, and thereafter gradually reduced and finally at the later stage reached comparable levels as in DT40 cells. In addition, mRNA levels of HDAC7, HDAC9 and PCAF were gradually elevated, and those of Pax5, Aiolos, E2A and EBF1 certainly changed in different patterns. Further, we cloned proximal ~4.9 kb

5'-upstream region of the Pax5 gene from DT40 genomic DNA by our original gene walking techniques. The ChIP assay suggested that acetylation levels of Lys-9 residues of histone H3 (K9/H3) within the proximal ~2.0 kb 5'-upstream chromatin region of the Pax5 gene were slightly decreased at the early stage in HDAC2(-/-) and at the later stage increased to comparable levels as in DT40 cells. We discussed on the relationship among these changing patterns in gene expressions of IgM H- and L-chains, specific transcription factors and chromatin-modifying enzymes, and acetylation levels of K9/H3 within the proximal ~2.0 kb 5'-upstream chromatin region of the Pax5 gene in HDAC2(-/-) mutants during continuous cultivation.

Methods

Cell cultures

Generation of HDAC2(-/-) DT40 mutants was described in our previous paper [24]. All positive mutant clones had been picked up on 96-well plates at ~10 days after knockout of two HDAC2 alleles, cultivated for another several days to be ~10⁶ cells/ml and stocked at -80 °C until use. HDAC2(-/-) mutant clones were continuously cultivated as described [24, 30] and collected at varying interval periods, including the early (E), middle (M) and later (L) stages of cultivation. The cultivation periods and stages were practically counted from the first day of cultivation from the stock at -80 °C.

2D-PAGE

Whole cellular proteins were prepared from HDAC2(-/-) mutant clone (cl.33-12) at the early (E; ~10 days), middle (M; ~30 days) and later (L; ~60 days) stages of cultivation and from DT40 (W), all of which were exponentially growing, and separated on 2D-PAGE using an automated apparatus Multiphor II (Amersham Pharmacia Biotech.), followed by staining with SYPRO Red (Molecular Dynamics, Inc.) as described [24, 37-39, 41].

Western blotting

Whole cellular proteins were prepared from three HDAC2(-/-) mutant clones (cl.33-28, cl.33-30 and cl.45-28) at different cultivation periods (up to ~56 days) and from DT40 (W) as described [25, 30]. Briefly, cells (1 x 10⁷) were treated with 10% trichloroacetic acid (TCA) and lysed in 100 µl of SDS buffer. Aliquots (10 µl) of 1:50 dilutions of the resultant cell extracts were subjected to 12% SDS-PAGE and electro-transferred to membranes, and proteins were detected with rabbit anti-chicken IgM L-chain antiserum A30-100A (Bethyl Laboratories Inc., TX, USA) that cross-reacts with IgM H-chain as described [28, 30]. Data analyses were carried out by Multi Gauge Ver3.X software using a luminescent image analyzer LAS-1000plus (FUJIFILM) [28, 30]. The relative amounts of IgM H-chain and large (high) and small (low) forms of IgM L-chain were measured at indicated periods.

Immuno-electron microscopy

Post-embedding immunocytochemistry was carried out as described [42]. Exponentially growing DT40 (W) and HDAC2(-/-) mutant clone (cl.33-28) at the early (E; ~10 days) and later (L; ~60 days) cultivation stages were collected by a light centrifugation, and the resultant cell pellets were fixed with 4% paraformaldehyde and 0.1% glutaraldehyde in 0.1 M phosphate buffer (PB) for 30 min. After washing with PB, the samples were post-fixed with 1% osmium tetroxide in PB for 60 min, washed with PB, dehydrated in ethanol and embedded in Epon. Ultrathin sections were picked up on 200-mesh gold grids coated with Formvar film and treated with 5% sodium meta-periodate in distilled water for 30 min. After rinsing in distilled water, the sections were treated with 5% normal horse serum (NHS) and 1% bovine serum albumin (BSA) in phosphate buffered saline (PBS) for 10 min to block nonspecific binding and incubated with goat anti-chicken IgM H-chain antibody A30-102A (Bethyl Laboratories Inc., TX, USA; diluted 1:1500 with 5% NHS, 1% BSA in PBS) at room temperature for 60 min. After rinsing in PBS, the sections were incubated with biotinylated horse anti-goat IgG (Vector Laboratories, Burlingame, CA, USA; diluted 1:200 with 1% BSA in PBS) for 40 min. After washing with PBS, the sections were incubated with 8 nm colloidal gold conjugated anti-biotin antibody (Vector Laboratories, Burlingame, CA, USA; diluted with 1% BSA in PBS) for 30 min. After washing with distilled water and drying, the sections were contrasted with 2% uranyl acetate in 70% methanol and Reynolds' lead citrate and observed in a JEOL 1200EX transmission electron microscope operating at 80 kV (JEOL, Tokyo, Japan). As a control, primary antibody was omitted or replaced by normal goat serum.

RT-PCR

Total RNAs were isolated from three independent HDAC2(-/-) mutant clones (cl.33-28, cl.33-30 and cl.45-28) at the early (E; ~20 days), middle (M; ~40 days) and later (L; ~60 days) stages of cultivation and from DT40 (W) as described [24, 30, 43]. RT-PCR was carried out using appropriate sense and anti-sense primers used in our previous papers [28, 30] and listed in Table I. HDAC2 and β -actin were used as negative and positive controls, respectively. RT-PCR products were subjected to 15% agarose gel electrophoresis. Data analyses were carried out as described above. Nucleotide sequences of all amplified products were confirmed by the PCR sequencing protocol as described [28, 30].

Immunoblotting

Whole cellular proteins were prepared from three independent HDAC2(-/-) mutant clones (cl.33-28, cl.33-30 and cl.45-28) at the early (E; ~20 days), middle (M; ~40 days) and later (L; ~60 days) stages of cultivation and from DT40 (W) (three different cultures) as described [30]. Briefly, cells (1×10^7) were treated with 10% TCA and lysed in 100 μ l of SDS buffer. Aliquots (10 μ l) of the resultant cell extracts

were subjected to 15% SDS-PAGE and electro-transferred to membranes. Bulk acetylation levels of specific Lys residues (K) of core histones H2A and H2B, H3 and H4 (indicated by appropriate designations) and the bulk methylation level of K9/H3 were measured by immunoblotting using site-specific anti-acetylated and anti-methylated histone antibodies as described [30]. Antibodies used were: anti-AcK9H3 (#06-942), anti-MeK9H3 (#07-212), anti-AcK14H3 (#06-911), anti-AcK18H3 (#07-354), anti-AcK23H3 (#07-355), anti-AcK27H3 (#07-360), anti-AcK8H4 (#06-760), anti-AcK12H4 (#07-323), anti-AcK16H4 (#07-329), anti-AcK5H2A (#07-290), anti-AcK7H2A (#07-386), anti-AcK9H2A (#07-289), anti-AcK5H2B (#07-382), anti-AcK12H2B (#07-336), anti-AcK15H2B (#07-343), anti-AcK16H2B (#07-341) and anti-AcK20H2B (#07-347) from Upstate and anti-AcK5H4 (AB3064) from CHEMICON.

Cloning of proximal ~4.9 kb 5'-upstream fragment of chicken Pax5 gene

Although at the present time the chicken Pax5 gene has been reported to exist on Z sex chromosome that is monosomy and to consist of 10 exons, the detailed information, including nucleotide sequences of the 5'-upstream region plus 5'-untranslational region of its first exon, was not available from USCS Genome Browser database until we had started this study. Only nucleotide sequences of chicken Pax5 cDNA [8] and 5'-upstream genomic DNA sequences of exon 1 up to position -109 (GenBank accession number: NM_204424) had been reported. To clone and identify the proximal ~4.9 kb 5'-upstream fragment of the Pax5 gene, we first screened the DT40 genomic DNA library in a λ FIXII [24] by PCR using sense primer (5'-GCACACTACGGACAACTTTTCC-3') and antisense primer (5'-CGGGGCCGCGTACATCTTCTCCAAATCCAT-3'), which were designed to amplify a part of exon 1 of the gene. However, no positive clones were yielded.

Therefore, we directly cloned the proximal ~4.9 kb 5'-upstream region of the Pax5 gene from DT40 genomic DNA by our original gene walking techniques, including Southern blotting [44], sub-cloning and colony hybridization (details will be shown later). First, the genomic DNA was digested with appropriate restriction enzymes, recognizing specific positions in the multiple cloning site of pBluescript II SK(+) vector, and hybridized with the PCR amplicon (as a probe), corresponding to the 85 bp fragment of exon 1 (positions -55 ~ +30), which was obtained from the genomic DNA using the above-mentioned primers. Among positive fragments obtained, the ~2.1 kb XbaI/ClaI fragment was sub-cloned into the multiple cloning site of pBluescript II SK(+) in frame as mentioned below. Enough amounts of the genomic DNA (200 μ g) were digested with XbaI and ClaI at 37 °C for 18 hrs and separated in 0.8% agarose gel, and the gel area containing the desired DNA fragment of ~2.1 kb was excised. The ~2.1 kb DNA fragments were extracted using the filter cartridge SUPECTM-01 (TaKaRa) following the manufacture's protocol and recovered by phenol/chloroform extraction and ethanol precipitation. The purified DNAs were dissolved in TE buffer (10 mM Tris-HCl, 1 mM EDTA; pH 8.0) and inserted into the

XbaI/ClaI site of pBluescript II SK(+) in molar ratio of 1 to 10 (for vector to insert) using DNA Ligation Kit Ver.1 (TaKaRa) at 16 °C for 21 hrs. The ligated plasmid was introduced into XL-1 blue MRF Competent cells, and then the cells were plated onto LB/amp plates, incubated at 37 °C for overnight, cooled at 4 °C for 1 hr, and subjected to colony hybridization with the 85 bp PCR amplicon as a probe. Our screening yielded one positive colony. Then, we sequenced the cloned ~2.1 kb DNA fragment by the dye terminator method using Big Dye Terminator V3.1 Sequencing Standard kit (Applied Biosystems).

As the next step, using the similar techniques, we cloned and identified nucleotide sequences of more additional ~3.0 kb 5'-upstream region of the Pax5 gene as follows. Based on the 5'-upstream genomic sequences mentioned above, sense 5'-TGCCATTTTCAAACACGCGC-3' and antisense 5'-GGAAAAGTTTGTCCGTAGTGTGC-3' primers were constructed. The 1626 bp PCR amplified product (corresponding to positions -1656 ~ -32) was obtained from the genomic DNA using these primers and digested with BssHII. After separation in 0.8% agarose gel, the band of desired 263 bp fragment was excised and purified using Wizard SV Gel and PCR Clean-Up System (Promega).

Successively, the resultant 263 bp DNA fragment, corresponding to the region from positions -1656 ~ -1394, was used as a probe for subsequent hybridization to obtain more additional 5'-upstream genomic fragments cut with appropriate restriction enzymes. Among positive fragments obtained, the ~4.9 kb BamHI/ClaI fragment was inserted into the BamHI/ClaI site of pBluescript II SK(+) in molar ratio of 1 to 5 (for vector to insert) and introduced into XL-1 blue MRF Competent cells. Then, the cells were plated onto LB/amp plates and subjected to colony hybridization. Finally, we isolated three positive clones, possessing the proximal ~4.9 kb 5'-upstream fragment from nucleotide number +241, and sequenced the isolated DNA fragment as described above.

Colony hybridization

We carried out colony hybridization on the Hybond-N+ nylon membrane filter following the manufacture's protocol (Amersham Biosciences). In brief, the membrane was laid on the surface of the agar plate for 45 sec. until it became thoroughly wetted, peeled carefully, and then its colony side was placed in upper direction onto a 10%(w/v) SDS saturated 3MMChr paper (Whatman) for 3 min. DNA was denatured and fixed to the membrane by successive treatments with denaturalization buffer (0.5 M NaOH, 1.5 M NaCl) for 7 min and neutralization buffer (1.5 M NaCl, 0.5 M Tris-HCl; pH 7.4) for 3 min twice. Finally, the membrane was vigorously washed in 2X SSC (0.3 M NaCl, 30 mM C₆H₅O₇Na₃-2H₂O; pH7.0) and its surface was gently scraped using KIMWIPE soaked in 2X SSC to remove cell debris. DNA was fixed to the membrane by a UV cross-linking procedure and hybridized with a ³²P-labeled probe by means of conventional methods.

Chromatin immunoprecipitation assay

Chromatin immunoprecipitation (ChIP) assay was done with ChIP assay kit (Millipore) according to manufacturer's instructions. Briefly, DT40 (W) and HDAC2(-/-) mutant clones (cl.33-28, cl.33-30 and cl.45-28) (1×10^6 cells) at the early (E; ~20 days) and later (L; ~60 days) stages of cultivation were cross-linked by the addition of 37% formaldehyde to a final concentration of 1% and incubated at 37 °C for 10 min, and the cross-linking was stopped with the addition of one-tenth volume of 1.25 M glycine to a final concentration of 0.125 M at 37 °C for 5 min. After removing medium, cells were washed with ice-cold PBS twice and re-suspended in 200 μ l of SDS Lysis Buffer freshly supplemented with 1 mM phenylmethylsulfonyl fluoride (PMSF) (Wako, Osaka, Japan) and 1 μ g/ml aprotinin (Sigma, St Louis, MO, USA), followed by sonication to solubilize and to shear cross-linked DNA. At this step, we used a Bioruptor UCD-250 (Cosmo Bio) at power L for ten 30-s pulses (30-s pause between pulses) at 4 °C to shear chromatin DNA into fragments of approximately 200 ~ 1000 bp. The samples were centrifuged at 13,000 rpm at 4 °C for 10 min to remove cell debris from the crude chromatin lysate. Sonicated cell supernatant was diluted to 1/10-fold by Chip Dilution Buffer. For an input, a portion (40 μ l; 2%) of the diluted cell supernatant was kept to quantify the amount of DNA present in each of different samples at the PCR step. Antibody (2 μ l) was added to 2 ml of the supernatant fraction and incubated at 4 °C for overnight. Incubation with 60 μ l of Protein Agarose/Salmon Sperm DNA was continued for 1 hr, followed by centrifugation at 1,000 rpm at 4 °C for 1 min. The immuno-precipitated complexes were sequentially washed with low-salt solution, high-salt solution, LiCl solution and TE (pH 8.0) twice. The antibody/histone/DNA complexes were eluted from Protein Agarose beads by adding 250 μ l of the elution buffer (1% SDS, 0.1% NaHCO₃) twice, and the cross-linking was reversed at 65 °C for 6.5 hrs by the addition of 5 M NaCl to a final concentration of 0.2 M. Samples were treated with proteinase K at 37 °C for 1 hr to remove proteins. DNA was recovered by phenol/chloroform extraction and ethanol precipitation using 40 μ g glycogen (NAKARAI TESQUE, INC.) as an inner carrier. Anti-AcK9H3 (#07-352) (Millipore) was used for pulldown. Recovered DNA fragments were subjected to PCR for 25 ~ 35 cycles, using appropriate primers (see Table II) specific for segments A to Q of approximately 150 ~ 250 bp, corresponding to the proximal 5'-upstream region and first exon (including the start codon) of the Pax5 gene. PCR was carried out using the same master reaction mixture in 25- μ l scale. Immuno-precipitated chromatin treated with non-immune rabbit serum (Vector Laboratories) (as a negative control) produced no bands following PCR (data not shown).

Results

IgM H- and L-chains artificially accumulated in HDAC2(-/-) DT40 mutants are gradually reduced during continuous cultivation

To know changing patterns in cellular protein levels of HDAC2(-/-) mutants during cultivation, we

first analyzed total cellular proteins prepared from clone cl.33-12 of HDAC2(-/-) mutants at the early (E; ~10 days), middle (M; ~30 days) and later (L; ~60 days) stages of cultivation and from DT40 cells (W) as a control by 2D-PAGE (Fig. 1). The amounts of IgM H-chain and L-chain (detected as two spots), whose amino acid sequences had been determined in our previous paper [24], were dramatically and certainly increased at the early cultivation stage in clone cl.33-12 compared with those as in DT40 cells. Surprisingly, the elevated amounts of these two immunoglobulin proteins in clone cl.33-12 were gradually reduced through the middle to later cultivation stages to almost similar levels in DT40 cells, whereas changes in cellular levels of most of other major proteins were insignificant during cultivation.

Next, we carried out Western blotting on total cellular proteins from clone cl.33-28 of HDAC2(-/-) mutants at shorter intervals up to 56 days, using the chicken IgM L-chain antibody that cross-reacts with IgM H-chain. As shown in Figure 2, protein levels of IgM H-chain and L-chain (detected as two bands) were dramatically increased at the early stage in clone cl.33-28, and thereafter these increased protein levels were gradually decreased through the middle to later stages to almost similar levels in DT40 cells. These results obtained in clones cl.33-12 and cl.33-28 by 2D-PAGE and Western blotting were confirmed in two other independent HDAC2(-/-) mutant clones (cl.33-30 and cl.45-28) (Fig. 3).

Moreover, we examined both the subcellular localization and changing pattern of the heightened intracellular level of IgM H-chain proteins in clone cl.33-28 during cultivation by immunocytochemistry using anti-chicken IgM H-chain antibody. Electron microscopy showed that the electron-dense materials were accumulated in the irregularly elaborated and vesiculated rough endoplasmic reticula only at the early stage in HDAC2(-/-) mutants (data not shown). Immuno-electron microscopy revealed that the colloidal gold immuno-labeling for chicken IgM H-chain was vividly observed in the electron-dense materials accumulating in the dilated and vesiculated rough endoplasmic reticula only at the early (~10 days) stage, and thereafter the heightened protein level was obviously decreased at the later (~60 days) stage to almost the same level in DT40 cells (Fig. 4).

These results revealed not only that IgM H- and L-chains were dramatically and considerably accumulated at the early stage of cultivation in HDAC2(-/-) mutant cells, but also that these accumulated immunoglobulin proteins were gradually reduced during cultivation and finally at the later stage reached comparable levels in DT40 cells. In addition, majority of artificially accumulated IgM H-chain, which existed as a native soluble form to be capable of building a high molecular weight complex with IgM L-chain in HDAC2(-/-) mutants [30], should be localized in rough endoplasmic reticula.

Alterations in gene expressions of IgM H-chain, IgM L-chain and core histones in HDAC2(-/-) DT40 mutants, respectively, are significant, moderate and insignificant during continuous cultivation

To know alterations in gene expressions of IgM H- and L-chains and core histones in HDAC2(-/-) mutants during cultivation, we prepared total RNAs from three independent clones (cl.33-28, cl.33-30 and cl.45-28) at the early (~20 days), middle (~40 days) and later (~60 days) stages of cultivation and from DT40 cells. First, we carried out RT-PCR on total RNAs, using primers IgM Hc, IgM Hs plus IgM Hm and IgM L specific for whole, secreted plus membrane-bound forms of IgM H-chain mRNA and IgM L-chain mRNA, respectively (Fig. 5). The levels of whole and secreted forms of IgM H-chain mRNA, respectively, were obviously and dramatically elevated at the early stage and thereafter gradually reduced through the middle to later stages in these mutant clones, whereas alterations in the levels of membrane-bound form of IgM H-chain mRNA and IgM L-chain mRNA were insignificant and moderate, respectively. These findings in the three mutant clones sufficiently agreed with those in clone cl.33-12 (data not shown). Next, we performed RT-PCR on total RNAs, using appropriate primers specific for core histones H2A, H2B, H3 and H4. In the three mutant clones, alterations in the mRNA levels of the four core histones were insignificant from the early to later stages (Fig. 5).

Gene expressions of HDAC7, HDAC9 and PCAF in HDAC2(-/-) DT40 mutants change during continuous cultivation

To clarify whether alterations in gene expressions of various members of chromatin-modifying enzymes (HDACs and HATs) are linked with those of IgM H- and L-chains during cultivation, we carried out RT-PCR on total RNAs, using appropriate primers specific for HDAC1, HDAC3, HDAC4, HDAC7, HDAC8, HDAC9 (equivalent to HDAC5 in our previous paper [30]), PCAF, GCN5, HAT1, ELP3, TIP60, MORF, MOZ and p300 (Fig. 6A). In three clones cl.33-28, cl.33-30 and cl.45-28, the HDAC9 mRNA level was certainly increased from the early (~20 days) to middle (~40 days) stages of cultivation and thereafter slightly decreased until the later (~60 days) stage. The HDAC7 mRNA level was slightly decreased at the early stage and remained unchanged until the later stage. However, the mRNA levels of residual HDACs did not change. On the other hand, in these mutant clones, the PCAF mRNA level was dramatically increased from the early to later stages of cultivation, whereas those of residual HAT members remained unchanged at any cultivation stages.

Gene expressions of E2A, EBF1, Pax5 and Aiolos in HDAC2(-/-) DT40 mutants differently change during continuous cultivation

To clarify whether alterations in gene expressions of various transcription factors are associated with those of IgM H- and L-chains during cultivation, we carried out RT-PCR on total RNAs, using appropriate primers specific for Oct1, Oct2, OBF1, NF- κ B, RelB, YY1, NF-AT, PU.1, E2A, EBF1, Pax5 and Aiolos (Fig. 6B). In three clones cl.33-28, cl.33-30 and cl.45-28, the mRNA level of E2A was slightly increased at the early (~20 days) stage of cultivation and thereafter remained unchanged

throughout the middle (~40 days) until later (~60 days) stages. The mRNA level of EBF1 was almost completely decreased at the early stage and thereafter remained unchanged as an undetectable level until the later stage. Interestingly, the mRNA level of Pax5 was obviously reduced at the early stage but thereafter certainly elevated through the middle to later stages. The mRNA level of Aiolos was observed to be low at the early stage, and further decreased from the middle to later stages. On the other hand, the mRNA levels of residual factors (Oct1, Oct2, OBF1, NF- κ B, RelB, YY1, NF-AT and PU.1) remained unchanged as a whole during cultivation.

Bulk acetylation levels of some specific Lys residues of core histones in HDAC2(-/-) DT40 mutants change during continuous cultivation

To know whether bulk acetylation levels of specific Lys residues (K) of core histones H2A, H2B, H3 and H4 change associated with alterations in gene expressions of IgM H- and L-chains during cultivation, we carried out immuno-blotting on total cellular proteins prepared from three clones (cl.33-28, cl.33-30 and cl.45-28) at the early (~20 days), middle (~40 days) and later (~60 days) stages of cultivation, together with those from DT40 cells as a control. Site-specific antibodies for various acetylated and methylated Lys residues of core histones H2A, H2B, H3 and H4 were used (Fig. 7). Concerning histone H2A, bulk acetylation levels of K5 and K9 or K7 residues remained unchanged or slightly changed during cultivation as a whole. Among the tested Lys residues of histone H2B, the bulk acetylation level of K16 was certainly reduced at the early stage and thereafter remained unchanged until the later stage, whereas those of remaining Lys residues unchanged at any cultivation stages. The bulk acetylation level of K5 of histone H4 was elevated at the early stage and remained unchanged until the later stage, but those of remaining Lys residues of histone H4 unchanged as a whole. Interestingly, concerning histone H3, bulk acetylation levels of K9, K14, K18, K23 and K27 were obviously elevated from the early throughout middle to later stages, whereas the bulk methylation level of K9 did not change. However, these results were somewhat uncertain yet because some antibodies used were relatively low specificities.

Cloning of proximal ~4.9 kb 5'-upstream region of chicken Pax5 gene

As will be mentioned in our next paper [40], in spite of insufficient information concerning the Pax5 gene existing on Z sex chromosome that is monosomy in chickens, fortunately, we could generate the Pax5-deficient DT40 mutant cell line Pax5(-), and clarify severe and moderate influences of its deficiency on gene expressions of IgM H- and L-chains and others. In order to know the way of the Pax5 gene expression, sufficient information on its proximal 5'-upstream region should be essential. However, as mentioned above, nucleotide sequences of the gene were not fully deposited in any database, whereas those on its homologues could be cited from GenBank. Therefore, we directly cloned the proximal ~4.9 kb 5'-upstream region of the Pax5 gene from DT40 genomic DNA by our original gene walking

techniques, including Southern blotting, colony hybridization and sub-cloning. Finally, the nucleotide sequences of 4950 bp 5'-upstream fragment and 241 bp ORF fragment of the chicken Pax5 gene were determined (GenBank accession number: LC060666) (Fig. 8). As a matter of course, the nucleotide sequences of XbaI/ClaI fragment fully agreed with those of BamHI/ClaI fragment from positions -1995 to +241. Our nucleotide sequences from positions -635 to +46 were surely coincident with those recently reported [12]. Dual-luciferase assay using various 3'-deletion and 5'-deletion mutant vectors in both HeLa and DT40 cells preliminarily suggested not only that as a whole the proximal ~4.9 kb 5'-upstream region negatively and very slightly affected on the Pax5 gene expression, but also that two 5'-upstream regions from positions -1561 to -974 and positions -521 to -40 were possibly involved in negative and bare regulation of the Pax5 gene expression, though these findings are uncertain (our unpublished data).

Acetylation levels of Lys-9 residues of histone H3 within proximal ~2.0 kb 5'-upstream chromatin region of chicken Pax5 gene in HDAC2(-/-) DT40 mutants probably change during continuous cultivation

In general, the chromatin structure surrounding the proximal 5'-upstream region of a certain gene is thought to directly and closely participate in its gene expression, regardless of the presence or absence of transcriptional element(s) within the region. The above-mentioned unpublished data obtained by the dual-luciferase assay suggested that at least the proximal ~1.6 kb 5'-upstream region of the Pax5 gene is necessary for its gene expression, though the interpretation is still uncertain. Furthermore, remarkably, as shown in Figure 6B, in HDAC2(-/-) mutants the gene expression of Pax5 was decreased at the early cultivation stage and thereafter increased until the later stage. Therefore, to know whether the chromatin structure surrounding the 5'-upstream region of the Pax5 gene changes during cultivation, we carried out chromatin immuno-precipitation (ChIP) assay, using site-specific antibody for acetylated Lys-9 residue of histone H3 (K9/H3) and appropriate primers (Table II), on its proximal ~2.0 kb 5'-upstream chromatin region in three clones (cl.33-28, cl.33-30 and cl.45-28) at the early (~20 days) and later (~60 days) cultivation stages and in DT40 cells as a control (Fig. 9). Acetylation levels of K9/H3 within the chromatin surrounding the ~2.0 kb 5'-upstream and ORF regions (from positions -1923 to +30) of the Pax5 gene were high in DT40 cells. Noticeably, in spite of the HDAC2-deficiency, in these mutant clones, as a whole, acetylation levels of K9/H3 within the ~2.0 kb 5'-upstream chromatin region of the gene were slightly decreased at the early stage and thereafter increased at the later stage, whereas those within the ORF region remained unchanged. In addition, in clone cl.33-28, acetylation levels of K14/H3 and K18/H3 within the proximal ~2.0 kb 5'-upstream chromatin region of the gene were observed to slightly change at the early and later stages (our unpublished data).

These findings suggested that acetylation levels of at least K9/H3 within the proximal ~2.0 kb 5'-upstream chromatin region of the Pax5 gene in HDAC2(-/-) mutants were reduced at the early

cultivation stage and thereafter at the later stage elevated to almost the same levels in DT40 cells. Thus, these results qualitatively agreed with those on alterations in the gene expression of Pax5 during cultivation.

Discussion

The modulation of chromatin topology with epigenetic modifications, such as acetylation, methylation, phosphorylation, ubiquitination, sumoylation, ADP ribosylation and others is one of the most important ways to express normal cell functions in eukaryotes. For the last several decades, mechanisms to modulate the chromatin structure with such epigenetic modifications have been intensively studied without interruptions in diverse life science fields, e.g., gene expression (transcription), DNA replication, differentiation, development, memory, pluri-potency, clinical medicine, etc. [1-3, 14-23, 45-72].

We have systematically studied *in vivo* roles of numerous members of histones, histone chaperones, HATs, HDACs and transcription factors [24-33, 37-41], using gene targeting techniques in the chicken B cell line DT40 possessing homologous recombination with a very high frequency [34-36]. To know individual roles of the changed transcription factors associated with the HDAC2-deficiency on the regulation of gene expressions of IgM H- and L-chains, we generated homozygous DT40 mutant cell lines EBF1(-/-), Aiolos(-/-), E2A(-/-) plus Helios(-/-) [30, 31], and also Pax5-deficient mutant cell line Pax5(-) [40] plus Ikaros-down mutant cell line Ikaros(-/+ (its generation will be shown elsewhere). Systematic analyses of these resultant mutants showed not only that Pax5, EBF1, Aiolos plus Ikaros down-regulate gene expressions of IgM H- and L-chains, and E2A up-regulates those of the two immunoglobulin proteins, but also that HDAC2 indirectly regulates gene expressions of IgM H- and L-chains through opposite regulations of gene expressions of Pax5, EBF1, Aiolos plus Ikaros, and E2A [30, 31].

In this study, throughout qualitative analyses of HDAC2(-/-) mutant cells collected at different cultivation periods, we accidentally noticed following surprising phenomena. 2D-PAGE showed not only that amounts of IgM H- and L-chains are dramatically and certainly elevated at the early stage of cultivation, and thereafter the elevated amounts of these two immunoglobulin proteins are gradually reduced and finally at the later stage reach comparable levels in DT40 cells (Fig. 1). On the other hand, cellular levels of most of other major proteins insignificantly change during cultivation. Western blotting, using the chicken IgM L-chain antibody that cross-reacts with IgM H-chain (Figs. 2 and 3), sufficiently confirmed the above-mentioned results obtained by 2D-PAGE. Immuno-electron microscopy using antibody specific for chicken IgM H-chain also showed that in HDAC2(-/-) the immunoglobulin proteins are clearly accumulated in endoplasmic reticula at the early stage, and thereafter the accumulated proteins are reduced at the later stage to almost similar level in DT40 cells (Fig. 4).

Further, RT-PCR using primers IgM Hc and Hs showed that whole and secreted forms of IgM H-chain mRNA are considerably and dramatically increased at the early stage, and thereafter the increased amounts of these two type mRNAs are gradually decreased through the middle to later stages to very close levels in DT40 cells, although IgM L-chain mRNA moderately changes during cultivation (Fig. 5). These results mean that gene expressions of IgM H- and L-chains are dramatically and slightly increased at the early stage and thereafter obviously decreased through the middle to later stages to almost similar levels in DT40 cells.

Interestingly, RT-PCR using appropriate primers specific for various genes encoding chromatin-modifying enzymes (HDACs and HATs) and transcription factors showed noticeable results as follows (Fig. 6). Concerning HATs, in HDAC2(-/-) the gene expression of PCAF is dramatically increased from the early through middle to later stages of cultivation. On the other hand, regarding HDACs, the gene expression of HDAC9 is surely increased from the early to middle stages and thereafter slightly decreased at the later stage, contrary to this, that of HDAC7 is slightly decreased at the early stage and remains unchanged during further cultivation. As for transcription factors, in HDAC2(-/-) the gene expression of EBF1 is almost completely decreased at the early stage and thereafter remains as an undetectable level until the later stage, but that of E2A is slightly increased at the early stage and remains unchanged during further cultivation. Surprisingly, the gene expression of Pax5 is obviously decreased at the early stage and thereafter certainly increases through the middle to later stages. That is, as a whole, in HDAC2(-/-) the changing pattern of the Pax5 gene expression during cultivation is in anti-parallel with those of IgM H- and L-chain gene expressions. Moreover, immuno-blotting using site-specific antibodies for various acetylated Lys residues (K) of core histones H2A, H2B, H3 and H4 showed that bulk acetylation levels of K9/H3, K14/H3, K18/H3, K23/H3 and K27/H3 are gradually elevated during cultivation, although insignificant or slight alterations are observed in those of remaining Lys residues of core histones (except significant changes of K16/H2B and K5/H4) (Fig. 7).

To explore the way of the gene expression of Pax5, first, we directly cloned and sequenced the proximal ~4.9 kb 5'-upstream region of the gene from DT40 genomic DNA by our original gene walking techniques, including Southern blotting, colony hybridization and sub-cloning (Fig. 8). Dual-luciferase assay showed that the proximal ~4.9 kb 5'-upstream region and especially the two regions from positions -1561 to -974 and positions -521 to -40 are thought to be possibly necessary for negative control of the Pax5 gene expression (our unpublished data). But these data are still uncertain and critical promoter region(s) or element(s) has not been determined yet. Finally, to know how does the gene expression level of Pax5 change in HDAC2(-/-) during cultivation, we analyzed the chromatin structure surrounding the proximal ~2.0 kb 5'-upstream region from positions -1923 to +30 of the gene at the early and later stages by the ChIP assay. Interestingly, acetylation levels of K9/H3 within some limited segments surrounding the proximal ~2.0 kb 5'-upstream chromatin region of the gene are reduced as a whole at the

early stage and thereafter elevated at the later stage to almost similar levels in DT40 cells (Fig. 9).

Taken together, these results indicated that the HDAC2-deficiency dramatically and considerably induces accumulations of IgM H- and L-chains depending on dramatic and considerable increases in their gene expression (mRNA) levels at the early stage of cultivation and thereafter the accumulated immunoglobulin mRNAs and proteins are gradually reduced throughout various generations during continuous cultivation. Such alterations in gene expressions of IgM H- and L-chains are coupled with those of gene expressions of some of Pax5, Aiolos, EBF1, E2A, PCAF, HDAC7 and HDAC9 and others. In addition, changes in acetylation levels of K9/H3 within the chromatin structure surrounding the proximal ~2.0 kb 5'-upstream region of the Pax5 gene in HDAC2(-/-) should be a basis for changes in its gene expression level through the chromatin structural change, resulting in alterations in gene expressions of IgM H- and L-chains. However, unfortunately, the early cultivation stage used in both RT-PCR and ChIP assay was considerably delayed from that used in other four experiments (2D-PAGE, Western blotting, immuno-blotting and immuno-electron microscopy), since the mutant cells used for the first two experiments had been cultivated longer periods than those used for the last four experiments. Therefore, to further elucidate the above-mentioned interesting biological phenomena, we must newly generate HDAC2-deficient DT40 mutant cells, collect large amounts of them (enough for all experiments) at the early (as earlier as possible), middle and later cultivation stages, and analyze their cellular characteristics at these three stages in detail.

Acknowledgments

The authors are grateful to Dr. T. Suganuma for immuno-electron microscopy study. And the authors thank Dr. Y. Takami for experimental support.

Conflict of interest statement

There are no conflicts of interest.

References

- [1] Kornberg, R. D.: Structure of chromatin. *Annu. Rev. Biochem.* 46, 931-954, 1977.
- [2] van Holde, K. E.: *Chromatin*. New York: Springer-Verlag, 1989.
- [3] Workman, J. L. and Kingston, R. E.: Alteration of nucleosome structure as a mechanism of transcriptional regulation. *Annu. Rev. Biochem.* 67, 545-579, 1998.
- [4] Peterson, M. L.: RNA processing and expression of immunoglobulin genes. In *Handbook of B and T Lymphocytes* (Snow, E. C. ed.) pp321-342, 1994, Academic press, San Diego.
- [5] Busslinger, M.: Transcriptional control of early B cell development. *Annu. Rev. Immunol.* 22, 55-79, 2004.

- [6] Nutt, S. L. and Kee, B. L.: The transcriptional regulation of B cell lineage commitment. *Immunity* 26, 715-725, 2007.
- [7] Ramirez, J., Lukin, K. and Hagman, J.: From hematopoietic progenitors to B cells: mechanisms of lineage restriction and commitment. *Curr. Opin. Immunol.* 22, 177-184, 2010.
- [8] Funahashi, J., Okufuji, T., Ohuchi, H., Noji, S., Tanaka, H. and Nakamura, H.: Role of Pax-5 in the regulation of a mid-hindbrain organizer's activity. *Dev. Growth Differ.* 41, 59-72, 1999.
- [9] Stevens, S., Ong, J., Kim, U., Eckhardt, L. A. and Roeder, R. G.: Role of OCA-B in 3'-IgH enhancer function. *J. Immunol.* 164, 5306-5312, 2000.
- [10] Savitsky, D. and Calame, K.: B-1 B lymphocytes require Blimp-1 for immunoglobulin secretion. *J. Exp. Med.* 203, 2305-2314, 2006.
- [11] Nera, K.-P., Kohonen, P., Narvi, E., Peippo, A., Mustonen, L., Terho, P., Koskele, K., Buerstedde, J.-M. and Lassila, O.: Loss of Pax5 promotes plasma cell differentiation. *Immunity* 24, 283-293, 2006.
- [12] Fujita, T. and Fujii, H.: Species-specific 5'-genomic structure and multiple transcription start sites in the chicken Pax5 gene. *Gene* 477, 24-31, 2011.
- [13] John, L. B. and Ward, A. C.: The Ikaros gene family: transcriptional regulators of hematopoiesis and immunity. *Mol. Immunol.* 48, 1272-1278, 2011.
- [14] Brownell, J. E., Zhou, J., Rannali, T., Kobayashi, R., Edmondson, D. G., Roth, S. Y. and Allis, C. D.: Tetrahymena histone acetyltransferase A: a homolog to yeast Gcn5 linking histone acetylation to gene activation. *Cell* 84, 843-851, 1996.
- [15] Ogryzko, V. V., Schiltz, R. L., Russanova, V., Howard, B. H. and Nakatani, Y.: The transcriptional coactivators p300 and CPB are histone acetyltransferases. *Cell* 87, 953-959, 1996.
- [16] Taunton, J., Hassig, C. A. and Schreiber, S. L.: A mammalian histone deacetylase related to the yeast transcriptional regulator Rpd3p. *Science* 272, 408-411, 1996.
- [17] Brown, C. E., Lechner, T., Howe, L. and Workman, J. L.: The many HATs of transcription coactivators. *Trends Biochem. Sci.* 25, 15-19, 2000.
- [18] Cheung, W. L., Briggs, S. D. and Allis, C. D.: Acetylation and chromosomal functions. *Curr. Opin. Cell Biol.* 12, 326-333, 2000.
- [19] Turner, B. M.: Histone acetylation and an epigenetic code. *Bioessays* 22, 836-845, 2000.
- [20] Jenuwein, T. and Allis, C. D.: Translating the histone code. *Science* 293, 1074-1080, 2001.
- [21] Roth, S. Y., Denu, J. M. and Allis, C. D.: Histone acetyltransferases. *Annu. Rev. Biochem.* 70, 81-120, 2001.
- [22] Carrozza, M. J., Utley, R. T., Workman, J. L. and Cote, J.: The diverse functions of histone acetyltransferase complexes. *Trend. Genet.* 19, 321-329, 2003.
- [23] Yang, X. J. and Seto, E.: Collaborative spirit of histone deacetylases in regulating chromatin structure and gene expression. *Curr. Opin. Genet. Dev.* 13, 143-153, 2003.

- [24] Takami, Y., Kikuchi, H. and Nakayama, T.: Chicken histone deacetylase-2 controls the amount of the IgM H-chain at the steps of both transcription of its gene and alternative processing of its pre-mRNA in the DT40 cell line. *J. Biol. Chem.* 274, 23977-23990, 1999.
- [25] Takami, Y. and Nakayama, T.: N-terminal region, C-terminal region, nuclear export signal and deacetylation activity of histone deacetylase-3 are essential for the viability of the DT40 chicken B cell line. *J. Biol. Chem.* 275, 16191-16201, 2000.
- [26] Nakayama, T. and Takami, Y.: Participation of histones and histone-modifying enzymes in cell functions through alterations in chromatin structure. *J. Biochem.* 129, 491-499, 2001.
- [27] Takeuchi, S., Adachi, M. and Nakayama, T.: Chicken HDAC2 down-regulates IgM light chain gene promoter activity. *Biochem. Biophys. Res. Commun.* 299, 263-267, 2002.
- [28] Kikuchi, H., Takami, Y. and Nakayama, T.: GCN5: a supervisor in all-inclusive control of vertebrate cell cycle progression through transcription regulation of various cell cycle-related genes. *Gene* 347, 83-97, 2005.
- [29] Kikuchi, H., Barman, H. K., Nakayama, M., Takami, Y. and Nakayama, T.: Participation of histones, histone modifying enzymes and histone chaperones in vertebrate cell functions. *Reviews and Protocols in DT40 Research*, Springer-Verlag, Berlin, pp225-243, 2006.
- [30] Nakayama, M., Suzuki, H., Yamamoto-Nagamatsu, N., Barman, H. K., Kikuchi, H., Takami, Y., Toyonaga, K., Yamashita, K. and Nakayama, T.: HDAC2 controls IgM H and L-chain gene expressions via EBF1, Pax5, Ikaros, Aiolos and E2A gene expressions. *Genes Cells* 12, 359-373, 2007.
- [31] Kikuchi, H., Barman, H. K., Nakayama, M., Takami, Y. and Nakayama, T.: Studies on epigenetic control of B cell functions using the DT40 cell line. *Advances in Genetics Research* 2, Urbano K. V. (Ed.), Nova Science Publishers, Inc. NY, pp153-166, 2010.
- [32] Kikuchi, H., Nakayama, M., Kuribayashi, F., Imajoh-Ohmi, S., Nishitoh, H., Takami, Y. and Nakayama T.: GCN5 is involved in regulation of immunoglobulin heavy chain gene expression in immature B cells. *Gene* 544, 19-24, 2014.
- [33] Escaffit, F., Vaute, O., Chevillard-Briet, M., Segui, B., Takami, Y., Nakayama, T. and Trouche, D.: Cleavage and cytoplasmic relocalization of histone deacetylase 3 are important for apoptosis progression. *Mol. Cell. Biol.* 27, 554-567, 2007.
- [34] Baba, T. W., Giroir, B. P. and Humphries, E. H.: Cell lines derived from avian lymphomas exhibit two distinct phenotypes. *Virology* 144, 139-151, 1985.
- [35] Buerstedde, J.-M. and Takeda, S.: Increased ratio of targeted to random integration after transfection of chicken B cell lines. *Cell* 67, 179-188, 1991.
- [36] Takeda, S., Masteller, E. L., Thompson, C. B. and Buerstedde, J.-M.: RAG-2 expression is not essential for chicken immunoglobulin gene conversion. *Proc. Natl. Acad. Sci. USA* 89, 4023-4027, 1992.

[37] Takami, Y., Takeda, S. and Nakayama, T.: An approximately half set of histone genes is enough for cell proliferation and a lack of several histone variants causes protein pattern changes in the DT40 chicken B cell line. *J. Mol. Biol.* 265, 394-408, 1997.

[38] Takami, Y. and Nakayama, T.: One allele of the major histone gene cluster is enough for cell proliferation of the DT40 chicken B cell line. *Biochim. Biophys. Acta* 1354, 105-115, 1997.

[39] Takami, Y. and Nakayama, T.: A single copy of linker H1 genes is enough for proliferation of the DT40 chicken B cell line, and linker H1 variants participate in regulation of gene expression. *Genes Cells* 2, 711-723, 1997.

[40] Nakayama, M. and Nakayama, T.: Generation of Pax5-deficient DT40 mutant cells, Pax5(-), and protein and mRNA levels of IgM H- and L-chains artificially and excessively accumulated in Pax5(-) DT40 mutants are rapidly and dramatically reduced through various generations during continuous cultivation. pp. 1-28, 2018. The revised article (the original paper was uploaded in 2017 and is available from following URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10145176>) will be uploaded in 2018.

The revised article is also the modified version of Chapter 3 of the revised monograph (the original monograph was published in 2015 and is available from following URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10123995>) as follows: Nakayama, M. and Nakayama, T.: Generation of Pax5-deficient DT40 mutants, Pax5(-), and protein and mRNA levels of IgM H- and L-chains artificially accumulated in Pax5(-) are rapidly and dramatically reduced through various generations during continuous cultivation. In: *Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations*, Nakayama, T. and Nakayama, M. (Eds.), pp. 45-71, 2018. The revised monograph is available from following URL:

<http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10169365>

[41] Takami, Y., Nishi, R. and Nakayama, T.: Histone H1 variants play individual roles in transcription regulation in the DT40 chicken B cell line. *Biochem. Biophys. Res. Commun.* 268, 501-508, 2000.

[42] Bendayan, M. and Zollinger, M.: Ultrastructural localization of antigenic sites on osmium-fixed tissues applying the protein A-gold technique. *J. Histochem. Cytochem.* 31, 101-109, 1983.

[43] Chomczynski, P. and Sacchi, N.: Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* 162, 156-159, 1987.

[44] Southern, E. M.: Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 98, 503-517, 1975.

[45] Biel, M., Wascholowski, V. and Giannis, A.: Epigenetics - an epicenter of gene regulation: histones and histone-modifying enzymes. *Angew. Chem. Int. Ed. Engl.* 20, 3186-3216, 2005.

[46] Margueron, R., Trojer, P. and Reinberg, D.: The key to development: interpreting the histone code?

Curr. Opin. Genet. Dev. 15, 163-176, 2005.

[47] Saha, A., Wittmeyer, J. and Cairns, B. R.: Chromatin remodelling: the industrial revolution of DNA around histones. *Nat. Rev. Mol. Cell Biol.* 7, 437-447, 2006.

[48] Bernstein, B. E., Meissner, A. and Lander, E. S.: The mammalian epigenome. *Cell* 128, 669-681, 2007.

[49] Goldberg, A. D., Allis, C. D. and Bernstein, B. E.: Epigenetics: a landscape takes shape. *Cell* 128, 635-638, 2007.

[50] Espada, J. and Esteller, M.: Epigenetic control of nuclear architecture. *Cell. Mol. Life Sci.* 64, 449-457, 2007.

[51] Shahbazian, M. D. and Grunstein, M.: Functions of site-specific histone acetylation and deacetylation. *Annu. Rev. Biochem.* 76, 75-100, 2007.

[52] Kouzarides, T.: Chromatin modifications and their function. *Cell* 128, 693-705, 2007.

[53] Allis, C. D., Berger, S. L., Cote, J., Dent, S., Jenuwien, T., Kouzarides, T., Pillus, L., Reinberg, D., Shi, Y., Shiekhatar, R., Shilatifard, A., Workman, J. and Zhang, Y.: New nomenclature for chromatin-modifying enzymes. *Cell* 131, 633-636, 2007.

[54] Lee, K. K. and Workman, J. L.: Histone acetyltransferase complexes: one size doesn't fit all. *Nat. Rev. Mol. Cell Biol.* 8, 284-295, 2007.

[55] Berger, S. L.: The complex language of chromatin regulation during transcription. *Nature* 447, 407-412, 2007.

[56] Esteller, M.: Cancer epigenomics: DNA methylomes and histone-modification maps. *Nat. Rev. Genet.* 8, 286-298, 2007.

[57] Suganuma, T. and Workman, J. L.: Crosstalk among histone modifications. *Cell* 135, 604-607, 2008.

[58] Kohn, K. W., Aladjem, M. I., Weinstein, J. N. and Pommier, Y.: Chromatin challenges during DNA replication: A systems representation. *Mol. Biol. Cell* 19, 1-7, 2008.

[59] Clapier, C. R. and Cairns, B. R.: The biology of chromatin remodeling complexes. *Annu. Rev. Biochem.* 78, 273-304, 2009.

[60] Selvi, R. B. and Kundu, T. K.: Reversible acetylation of chromatin: implication in regulation of gene expression, disease and therapeutics. *Biotech. J.* 4, 375-390, 2009.

[61] Talbert, P. B. and Henikoff, S.: Histone variants-ancient wrap artists of the epigenome. *Nat. Rev. Mol. Cell Biol.* 11, 264-275, 2010.

[62] Javierre, B. M., Hemando, H. and Ballestar, E.: Environmental triggers and epigenetic deregulation in autoimmune disease. *Discov. Med.* 12, 535-545, 2011.

[63] Bannister, A. J. and Kouzarides, T.: Regulation of chromatin by histone modifications. *Cell Res.* 21, 381-395, 2011.

[64] Verrier, L., Vandromme, M. and Trouche, D.: Histone demethylases in chromatin cross-talks. *Biol.*

Cell 103, 381-401, 2011.

[65] Butler, J. S., Koutelou, E., Schibler, A. C. and Dent, S. Y.: Histone-modifying enzymes: regulators of developmental decisions and drivers of human disease. *Epigenomics* 4, 163-177, 2012.

[66] Kooistra, S. M. and Helin, K.: Molecular mechanisms and potential functions of histone deacetylases. *Nat. Rev. Mol. Cell. Biol.* 13, 297-311, 2012.

[67] Oliner, J. R. and Winston, F.: Chromatin and transcription in yeast. *Genetics* 190, 351-387, 2012.

[68] Narlikar, G. J.: Mechanisms and functions of ATP-dependent chromatin-remodeling enzymes. *Cell* 154, 490-503, 2013.

[69] Graff, J. and Tsai, L.-H.: Histone acetylation: molecular mnemonics on the chromatin. *Nat. Rev. Neurosci.* 14, 97-111, 2013.

[70] Chen, T. and Dent, S. Y. R.: Chromatin modifiers and remodellers: regulators of cellular differentiation. *Nat. Rev. Genet.* 15, 93-106, 2014.

[71] T. W.-W. and Reinberg, D.: Chromatin futures and the epigenetic regulation of pluripotency states in ESCs. *Development* 141, 2376-2390, 2014.

[72] Morgan, M. A. and Shilatifard, A.: Chromatin signatures of cancer. *Genes and Dev.* 29, 238-249, 2015.

Figure legends

Figure 1. Alterations in amounts of IgM H- and L-chain proteins in HDAC2(-/-) DT40 mutants during continuous cultivation.

Total cellular proteins were prepared from HDAC2(-/-) mutant clone (cl.33-12) at the early (E; ~10 days), middle (M; ~30 days) and later (L; ~60 days) stages of cultivation and compared with those from DT40 (W) by 2D-PAGE. Iso-electrofocusing in the first dimension (pI) and SDS-PAGE in the second dimension (MW) were performed on Immobiline DryStrip gel (pH 4-7) and ExcelGel XL SDS gel (gradient 12-14), followed by the fluorostaining method. H and L indicate IgM H-chain and L-chain (detected as two spots), respectively.

Figure 2. Alterations in protein levels of IgM H- and L-chains in HDAC2(-/-) DT40 mutants during continuous cultivation.

Total cellular proteins were prepared from HDAC2(-/-) mutant clone (cl.33-28) at indicated cultivation periods and compared with those from DT40 (W) by Western blotting. Aliquots of cell extracts were subjected to 12% SDS-PAGE and transferred to membrane filter. Proteins were detected with anti-chicken IgM L-chain antiserum (as primary antibody) that cross-reacts with IgM H-chain. Relative levels of IgM H-chain and large (high) and small (low) forms of IgM L-chain are shown in the lower panel. IgM H and IgM L indicate IgM H- and L-chains, respectively.

Figure 3. Alterations in protein levels of IgM H- and L-chains in two other HDAC2(-/-) DT40 mutant clones during continuous cultivation.

Total cellular proteins were prepared from two other HDAC2(-/-) mutant clones (cl.33-30 and cl.45-28) at indicated cultivation periods and analyzed as in Figure 2. However, these periods were not coincident with those in Figure 2.

Figure 4. Alterations in amounts of IgM H-chain protein in HDAC2(-/-) DT40 mutants during continuous cultivation.

HDAC2(-/-) mutant clone (cl.33-28) at the early (E; ~10 days) and later (L; ~60 days) stages of cultivation and DT40 (W) were fixed, and immuno-electron microscopy with anti-chicken IgM H-chain antiserum was performed. A) Study in wide range. A number of cells were observed in (W), (E) and (L), but accumulated IgM H-chain proteins were detected only in mutant cells in (E). B) Study for single cell. Each of cells indicated by arrows in A) was magnified and shown in (W), (E) and (L). Large amounts of accumulated IgM H-chain protein were observed only in (E).

Figure 5. Alterations in gene expressions of IgM H- and L-chains, and core histones in

HDAC2(-/-) DT40 mutants during continuous cultivation.

Total RNAs were extracted from three HDAC2(-/-) mutant clones (cl.33-28, cl.33-30 and cl.45-28) at the early (E; ~20 days), middle (M; ~40 days) and later (L; ~60 days) stages of cultivation and from DT40 (W). RT-PCR was performed on equal amounts of total RNAs, using appropriate primers for HDAC2, whole (IgM Hc), secreted (IgM Hs) plus membrane-bound (IgM Hm) forms of IgM H-chain and IgM L-chain (IgM L), and four core histones (H2A, H2B, H3 and H4). Chicken β -actin was used as a control. Some data for DT40 (W) and HDAC2(-/-) at the early (E) stage were identical with those in our previous paper [30].

Figure 6. Alterations in gene expressions of various members of HDACs, HATs and transcription factors in HDAC2(-/-) DT40 mutants during continuous cultivation.

Total RNAs used were the same ones as in Figure 5. RT-PCR was performed in the same procedure, using appropriate primers for various members of HDACs and HATs (A) and transcription factors (B), instead of those used in Figure 5. Some data for DT40 (W) and HDAC2(-/-) at the early (E) stage were identical with those in our previous paper [30].

Figure 7. Alterations in bulk acetylation and methylation levels of Lys residues of core histones in HDAC2(-/-) DT40 mutants during continuous cultivation.

Whole cellular proteins were extracted from three HDAC2(-/-) mutant clones (cl.33-28, cl.33-30 and cl.45-28) at the early (E; ~20 days), middle (M; ~40 days) and later (L; ~60 days) stages of cultivation and from DT40 (W) (three separate cultures) and subjected to 15% SDS-PAGE. Bulk acetylation levels of specific Lys residues (K) (indicated by appropriate designations) of core histones H3, H4, H2A and H2B were measured by immunoblotting using various site-specific anti-acetylated histone antibodies. The bulk methylation level of K9/H3 was measured using site-specific anti-methylated histone antibody. Most data for DT40 (W) and HDAC2(-/-) at the early (E) stage were identical with those in our previous paper [30].

Figure 8. Nucleotide sequences of the proximal ~4.9 kb 5'-upstream region of chicken Pax5 gene.

Nucleotide sequences of the proximal 5'-upstream region of the chicken Pax5 gene up to the position -4950 (including exon 1, putative transcription start site and translation initiation triplet) are shown.

Figure 9. Alterations in acetylation levels of Lys-9 residue of histone H3 within proximal 5'-upstream chromatin region of Pax5 gene in HDAC2(-/-) DT40 mutants during continuous cultivation.

Chromatin immunoprecipitation (ChIP) assay was carried out using Chromatin Immunoprecipitation

Assay kit. Cross-linked chromatin prepared from cell lysates of three HDAC2(-/-) mutant clones (cl.33-28, cl.33-30 and cl.45-28) at the early (E; ~20 days) and later (L; ~60 days) stages of cultivation and from DT40 (W) were co-precipitated by antiserum specific for acetylated Lys-9 residue of histone H3 (K9/H3). After de-crosslinking, co-precipitated chromatin was amplified by PCR using appropriate primers (see Table II) for segments A to Q of the proximal ~2.0 kb 5'-upstream region of the Pax5 gene (see Fig. 8). PCR products were analyzed by 1.5% agarose gel electrophoresis.