

Memoirs of a life scientist: Interpretations and opinions on life phenomena, and experiences and contributions in life science field ~ The chromatin conformation change code (4C) theory and previous researches (4th Edition) ~

Tatsuo Nakayama

Section of Biochemistry and Molecular Biology, Department of Medical Sciences, Faculty of Medicine, Department of Life Science, Frontier Science Research Center, University of Miyazaki, 5200, Kihara, Kiyotake, Miyazaki, 889-1692, Japan

Corresponding to E-mail: tnakayam@med.miyazaki-u.ac.jp

Footnote

This article is the 4th Edition of the Paper-7 on the Memoirs/Interpretations (the 1st, 2nd and 3rd Editions of the Paper were published in 2019, 2020 and 2021 [198, 223] are respectively available from following new URLs: <http://hdl.handle.net/10458/6614>, <http://hdl.handle.net/10458/6949> and also <http://hdl.handle.net/10458/6949>).

All figures used in this article are identical with corresponding those in the 1st, 2nd and 3rd Editions [198, 223], and also are originally selected from those in the 1st and 2nd Editions of my Retirement Commemorative Monograph published in 2015 and 2018 [138], respectively, which are available from following new URLs: <http://hdl.handle.net/10458/5293> and <http://hdl.handle.net/10458/6460>.

This article also contains a part of Chapter-7 of the Monograph (2nd Edition) [138] and also the 5th Edition (published in 2021) of the Paper-6 [222] as follows: Nakayama, T. and Nakayama, M.: A comprehensive and detailed review on the chromatin conformation change code (4C) theory: A theory on ways to gain un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through numerous generations in higher eukaryotes. pp.1-85, 2021. The 5th Edition of the Paper-6 is available from following new URL: <http://hdl.handle.net/10458/6902>.

This article is fundamentally the 4th-revised Edition of Chapter-8 of the 2nd Edition of the Retirement Commemorative Monograph [138] as follows: Nakayama, T.: Personal closing remarks on the occasion of retirement: The real course and history of our research prior to the 4C theory, personal interpretation on life phenomena, and experiences and opinions as a life scientist. In: Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations. Nakayama, T. and Nakayama, M. (Eds.), pp.1-303, 2018. The 2nd Edition of the Monograph (the 1st Edition was published in 2015 and is available from following new URL: <http://hdl.handle.net/10458/5293>) is available from following new URL: <http://hdl.handle.net/10458/6460>.

Summary

We advanced the chromatin conformation (structure) change code (4C) theory for the bio-system to gain un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through numerous generations. The 4C theory had been studied under the worst research environments, i.e., without any research grants, in our spare time from my main research projects and that by only a womanpower. The 4C theory is the typical fruit of my serendipity and the outcome of some 10 years of my small research group. In this article, I described the detail of the 4C theory and the outline on real course of my researches prior to the theory, my personal interpretations of the concept on life phenomena, and experiences and opinions as an ordinary life scientist.

In eukaryotes, both genome information and nuclear function are mainly protected by the nuclear membrane that acts as a nucleus-cytoplasm barrier. Additionally, communication and signal transduction between the nucleus and cytoplasm, both of which are essential for expressions of normal cell functions, are preferentially carried out by the to and fro passage of large and/or small molecules via the nuclear pore that acts as a guard station at the barrier. By contrast, both the signal transduction concerning an abnormal, unexpected, uncomfortable and/or disadvantageous change in the intra- and/or extra-cellular environment and the transport of useless molecules to the nucleus are usually prevented by this barrier system. Then, how do the higher eukaryotic cells cope with and/or overcome an abnormal, unexpected and/or disadvantageous environment change, when they firstly encounter it? Generally, this is possible by adhering to four typical countermeasures. First, in the case of a very severe change, cells die because it is far ahead of their adaptation ability. Second, in the case of a considerably severe change, cells cope with or overcome it by means of alterations in genome information, such as point mutation, insertion, deletion, duplication and multiplication on the DNA molecule through numerous generations. This mode is a basis for evolution of species. Of course, the mode also occurs in all of organisms, including virus, prokaryotes, lower eukaryotes, higher eukaryotes and others. Third, in the case of a moderate change, cells cope with or overcome it by means of irreversible creation of chromatin structure plasticity caused by successive chromatin conformational (structural) changes with epigenetic modifications through numerous generations (or cell divisions). This mode is a basis for the development and differentiation of higher eukaryotic cells. Fourth, in the case of a minor change, cells respond to it only through already acquired regulation mechanisms.

As will be detailed later, we accidentally discovered that IgM H- and L-chains artificially and excessively accumulated in HDAC2(-/-) DT40 mutant cells (and Pax5(-) DT40 mutant cells) at the early cultivation stage are dramatically decreased through numerous generations (cell divisions) during

continuous cultivation. This surprising and remarkable biological event, which should be one of the most fundamental and important ways for the adaptation to the environment change and life conservation of higher eukaryotes, is really an example of the third case of the above-mentioned countermeasures, which are naturally classified in my own way. Based on these findings, we advanced the chromatin conformation (structure) change code (4C) theory for the bio-system to gain un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through numerous generations (cell divisions).

Since chemical modifications of histones with acetyl and methyl groups were first proposed to regulate RNA synthesis [1], the modulation of chromatin topology has been thought to be one of the most fundamental and important events for expression of normal cell functions in eukaryotes [2-4]. The ways to modulate the chromatin structure with acetylation, methylation, phosphorylation, ubiquitination, sumoylation, ADP-ribosylation and others have been intensively studied in a variety of life science fields. Of these epigenetic modifications, acetylation and deacetylation of specific Lys residues of core histones (H2A, H2B, H3 and H4) cooperatively controlled by chromatin-modifying enzymes, such as histone acetyltransferases (HATs) and histone deacetylases (HDACs), are undoubtedly the major ones [5-32]. For the past several decades, countless numbers of research papers on the acetylation and deacetylation (and other epigenetic modifications) are accumulating without interruption in more diverse life science fields, e.g. transcription/gene expression, DNA replication, repair plus recombination, development, differentiation, memory, pluri-potency, clinical medicine, etc. [33-44].

Naturally, alterations in the chromatin structure are remarkably involved in regulations of the lymphocyte development and differentiation [45-51], which are closely related to one of my latter research projects as will be mentioned later. In addition, various transcription factors, including Ikaros, PU.1, E2A, GATA-3, EBF, Pax5 and others, are involved in regulations of the development and differentiation of lymphocytes [52-61]. And then, the regulation of IgM H-chain gene expression requires USF, TFEB, Ig/EBP, NF-IL6, OCA-b, etc. as promoter binding proteins, and Ig/EBP, NF-IL6, YY-1, E2A, PU.1, etc. as intron enhancer binding proteins.

However, my research group had mainly studied on structures (nucleotide sequences, transcriptional elements and genome organizations) of core histones and linker histone H1 of chickens until ~25 years ago (~1995) [62-72]. Prior to these researches, during more past ~25 years (started from the master course of graduate school), I had mainly studied on determination of amino acid sequences of protein (cytochrome c of bonito), in vitro and in vivo acetylation and synthesis of protein (ovalbumin) in hen's oviduct, RNA synthesis by RNA polymerase of *B. subtilis* during sporulation (the reasons for which were as follows), 2D-PAGE analysis of proteins, and structure and function of DNA binding proteins and so on

[73-90]. At that time, 1) sporulation of *B. subtilis* with alterations in both gene expression and morphology was considered as a simple model system for differentiation of higher eukaryotic cells, and 2) I had the hope as that in the future I would like to study both individual roles of several kinds of RNA polymerases and complicated mechanisms of varied cell-specific gene expressions in higher eukaryotes. Figuratively speaking, if I compare my whole researches to one game, those must be the pre-run of the hop, step and jump.

In the middle of the above-mentioned studies concerning structures of histone genes, the DT40 cell line, which was derived from chicken pre-B lymphocyte cells [91], was reported to possess an ability of homologous recombination with a very high frequency and to be a powerful system for gene targeting techniques [92, 93]. I thought that an approach using the gene targeting techniques in DT40 cells is very advantageous for clarifying fundamental and/or common phenomena in higher eukaryotic cells, besides specific and/or limited functions in the B lymphocyte cell lineage. I was confident that even my small group could continue any of researches using the ALV-transformed chicken DT40 cell line, although both the start and continuance of the researches using model animals (mice, rats, *C. elegans*, *D. melanogaster*, *Xenopus*, etc.) should be very hard for us because of insufficient grants and manpower. In addition, I was fully convinced that the peculiar knowledge on basic bio-systems obtained in DT40 cells could be universally extended to understanding fundamental and complex bio-systems in higher eukaryotes. Because almost all concepts of the most essential life phenomena, including “central dogma”, which had been elucidated in phage, prokaryote (especially *E. coli*) and/or yeast systems, could be thoroughly applicable to those of higher eukaryotes.

Based on the above-mentioned mature consideration, in parallel with studies on in vitro functions of histone chaperones (CAF-1 and HIRA) [94-102], I had immediately changed my main research projects to the elucidation of in vivo functions (and structures) of several chromatin-modifying enzymes (HATs and HDACs) and transcription factors, as well as histones and histone chaperones (CAF-1, HIRA, ASF1 and others), using the gene targeting techniques in the DT40 cell line. Of course, I had no hesitation in using the DT40 cell line as an experimental system for my research projects at all, because I had become accustomed to the chicken system through studies on both bio-synthesis and post-translational (i.e., epigenetic) modification such as acetylation and glycosylation of ovalbumin as briefly mentioned above. First, to assess individual roles of members of histones and histone chaperones, we had comprehensively generated numerous homozygous DT40 mutant cell lines, all of which lacked their corresponding specific genes, respectively. The results obtained from resultant DT40 mutants, combined, showed that this approach was useful and powerful to know individual functions of these specific genes in vivo really [103-113].

In parallel and/or continuously, based on these positive results and my intrinsic interest on universal

and global roles of histone acetylation and/or deacetylation, in order to assess individual roles of members of HATs and HDACs, we had systematically generated many homozygous DT40 mutant cell lines, each of which was devoid of a specific member of HATs and HDACs, such as GCN5, PCAF, HAT1, HDAC1, HDAC2, HDAC3 and others [114-131]. Our qualitative analyses of the HDAC2-deficient mutant cell line HDAC2(-/-), which was initially generated by us [114], revealed not only that HDAC2 regulates the amount of IgM H-chain at the two steps of its gene expression and alternative pre-mRNA processing [114], but also that it down-regulates IgM L-chain gene promoter activity [117]. The lacking of HDAC2 also represses gene expressions of HDAC7, Pax5, Aiolos, Ikaros and EBF1, elevates gene expressions of HDAC4, HDAC9, PCAF and E2A, and changes bulk acetylation levels of several Lys residues of core histones [121]. Further, we generated Pax5(-), Aiolos(-/-), Ikaros(-/-/+), EBF1(-/-), E2A(-/-), Helios(-/-) and other mutants [121, 132-134], and thereby clarified specific functions of these transcription factors in B lymphocytes [121, 129, 132-138]. In consequence, these results, together with others [139], revealed that Pax5, EBF1, Aiolos and Ikaros down-regulate gene expressions of IgM H- and L-chains, and inversely E2A and probably OBF1 up-regulate these two immunoglobulin gene expressions. Therefore, we concluded that HDAC2 as a supervisor indirectly regulates gene expressions of IgM H- and L-chains through opposite regulation of gene expressions of Pax5, EBF1, Aiolos, Ikaros, E2A and OBF1 [121, 124, 138].

However, until then, I had never met my own exciting research that takes me captive and makes my heart beat fast with joy, and I had not been throbbing heavily with almost all so-called brilliant and/or fashionable achievements in life science field with a few exceptions. I chiefly owed such feelings to the fact that I have been fascinated by the word “life” from boyhood; therefore, I had a great interest in fundamental concepts of life phenomena but not so much interest in any of individual vital substances, reactions and/or functions, even though they are very important. Surely, it goes without saying that still now countless important questions remain to be resolved in the life science field. By the way, I previously reported not only that extracellular ovalbumin and egg-white proteins inhibit ovalbumin and protein syntheses in hen’s oviduct, respectively [76, 78], but also that structural changes of specific histone genes possibly result in differences in their transcriptional regulations in chicken different tissues, the reason for which is still unknown [66, 67]. In addition, by gene targeting techniques in chicken DT40 cells, we reported that all histone gene families have the inherent ability to keep constant protein levels of respective histone subtypes, all of which are very important for life conservation, when a certain gene member of the corresponding subtype is deleted [103-108, 115]. Therefore, for these above-mentioned reasons, as will be described later in detail, I had much continued to have an interest in adaptability of higher eukaryotes to environment change (one of most fundamental concepts of life phenomena), including both the structural changes of the specific genes among different cells (tissues)

and the compensation ability. Because the regulation ways of ovalbumin and egg-white protein syntheses could be considered as a response ability of hen's oviduct to extracellular environment change, the structural alterations of specific genes (and naturally their chromatin) among different tissues should be thought as a basis for determination of cell type-specificity, and also the compensation ability should be thought as a response to intracellular environment change, i.e., the decrease in protein levels of specific histone member.

In the middle of re-constitution (re-expression) experiments to study the participation of HDAC2 in the regulation of gene expressions of IgM H- and L-chains in detail, we became aware of the surprising phenomena as follows. We first introduced the tetracycline- (tet-) repressive HDAC2 cDNA into HDAC2(-/-) mutants, and isolated the transformed cell line HDAC2(-/-)tetHDAC2(+), lacking two alleles of the endogenous HDAC2 gene but possessing the artificial tet-repressive HDAC2 cDNA. Contrary to our expectation, surprisingly, protein and mRNA levels of IgM H- and L-chains were very low in HDAC2(-/-)tetHDAC2(+) even in the case of non-existence of HDAC2 due to the presence of tet, as well as both in the case of existence of HDAC2 due to the absence of the drug and in wild-type DT40 cells having the enzyme (our unpublished data).

Concerning these findings, members of my research group each had different interpretations. One of colleagues intensely asserted not only that the accumulation of IgM H- and L-chains in HDAC2(-/-) was not truly based on the HDAC2-deficiency and resulted from only unknown artificial cause of beneath our notice, but also that the accumulated immunoglobulin proteins were only reduced spontaneously during cultivation to construct HDAC2(-/-)tetHDAC2(+). Since the colleague was bigoted toward a vague biased (but ordinary in a sense) view that several properties of higher eukaryotic cells are easily changeable during cultivation. Similarly, another colleague conservatively insisted that the accumulation and diminution of IgM H- and L-chains in HDAC2(-/-) were not directly due to the HDAC2-lacking but based on only somatic mutations occurred within the genome locus of the IgM H-chain gene during cultivation to construct the transformants, since the locus was generally considered as a hot spot for the mutations [93]. Thus, they were not interested in these findings no more.

However, intuitively, I believed that the accumulation of IgM H- and L-chains in HDAC2(-/-) was really caused by the HDAC2-deficiency itself, and their diminution during cultivation to construct HDAC2(-/-)tetHDAC2(+) should be attributable to unknown important biological mechanisms. Such my own opinion was supported in part by our following findings. First, several re-experiments (2D-PAGE, Western blotting and Northern blotting) showed that protein and mRNA levels of IgM H-chain (and also L-chain) were firstly increased in several independent HDAC2(-/-) mutant clones and thereafter reproducibly decreased, whenever they were cultivated for a long period (our unpublished data). Second, by contrast, insignificant changes (increases and/or decreases) in protein and mRNA levels of

IgM H- and L-chains were observed in numerous DT40 mutant cell lines (approximately 30), each of which was devoid of a member of HDACs (except HDAC2), HATs, transcription factors, histones and histone chaperones, etc. [103-109, 121, our unpublished data].

Meanwhile, in the process of the above-mentioned consideration and re-examinations, I noticed some following mistaken speculations due to qualitative results in our previous paper [114], besides those on the insufficient and uncertain results in our other papers [84, 86-90]. Re-experiments showed that protein and mRNA levels of HDAC2 in the heterozygous mutants HDAC2(-/+) were obviously decreased to ~50% of those in DT40 cells (our unpublished data), whereas these two levels had been reported to be kept constant by means of the compensation mechanism [103-109]. Re-examined RNase protection assay (and Northern blotting) showed that the amounts of IgM H-chain protein and ratio of secreted to membrane-bound forms of its mRNA were certainly increased in HDAC2(-/+), compared with those in DT40 cells (our unpublished data), though both the amounts and the ratio had been reported to remain unchanged. The growth rate of initially generated HDAC2(-/-) mutant clones re-tested (our unpublished data), as well as that of most clones of secondly generated HDAC2(-/-) mutants [121], was slightly but truly different from that of DT40 cells, though the growth rate of the initial mutants had been reported to be essentially identical with that of DT40 cells [114]. Pulse-labeling/chase experiment and Northern blotting re-examined showed that protein and mRNA of IgM L-chain really existed as large and small forms and the amounts of the two forms were surely increased associated with the HDAC2-deficiency (our unpublished data), though the small form of IgM L-chain had been considered as a specific proteolytic product of the large form at the C-terminal region.

Based on my consideration and enthusiasm as mentioned above, I asked again some colleagues of my research group to study mechanisms of the increases and decreases in protein and mRNA levels of IgM H- and L-chains in HDAC2(-/-) as a part of their own research works. However, several kinds of approaches gave only negative (unpublished) data as follows. RT-PCR performed on various transcription factors (including some promoter and intron enhancer binding proteins mentioned above) for the IgM H-chain gene expression showed only insignificant (or no) changing patterns in their mRNA levels between DT40 cells and HDAC2(-/-) mutants at any cultivation stages. MNase protection assay carried out on several regions within the chromatin surrounding the IgM H-chain gene showed insignificant changing patterns in the nucleosome ladders between DT40 cells and HDAC2(-/-) mutants at any cultivation stages. In addition, MNase protection assay performed on the chromatin of several genes encoding transcription factors (including Pax5, Aiolos, Ikaros and EBF1) also showed insignificant changing patterns in the nucleosome ladders between DT40 cells and HDAC2(-/-) mutants at any cultivation stages.

On the other hand, interestingly, Northern blotting using the specific probe for the IgM H-chain gene sometimes showed a mere change in amounts of small positive-like RNA molecules having a possibility

of alternative splicing product of IgM H-chain mRNA between DT40 cells and HDAC2(-/-) mutants at the early and later cultivation stages (our unpublished data). Since small RNA molecules (such as small nuclear RNA, microRNA, small interference RNA, anti-sense RNA, etc.) were watched as important players in the regulation of cell functions at that time, we temporarily focused our attention on these molecules. However, unfortunately, these findings were not reproducible; therefore, I decided that in our case the small RNA molecules were not worth studying ever since.

For all that, I believed that these negative results should be caused only by the following facts: 1) IgM H- and L-chain genes are indirect but not direct targets of HDAC2 [121]. 2) MNase protection assay was regrettably carried out only on coding regions (open reading frame (ORF) regions) but not on proximal 5'-upstream regions of these transcription factor genes. 3) HDAC2(-/-) mutants regarded as samples at the early stage must be collected really at vague early cultivation stage, i.e., at later and varied cultivation stages fairly, because they had been already cultivated for a long time to use for other experiments until then. Therefore, I resolved to analyze again properties of the initially generated HDAC2(-/-) mutants collected at different cultivation periods as correctly as possible. Further, particularly, I should like to emphasize that we must newly generate HDAC2(-/-) mutants, collect them at different cultivation periods from virtual early (as early as possible) until later stages, and systematically and closely analyze their various cellular properties, in order to solve the above-mentioned problems.

Then, I thought that a full-time researcher, but not a part-time one, must be absorbed in the project to clarify mechanisms for both the accumulation and the diminution of proteins and mRNAs of IgM H- and L-chains in HDAC2(-/-) mutants through numerous generations (cell divisions) during continuous cultivation. I asked Dr. Masami Nakayama, my wife, to work with me as a colleague (postdoctoral researcher). She had worked on this project and others listed in the References section full-time, in spite of a part-time contract (few hours per day for several years and several hours per day for only one year). Further, another PhD joined my group as an assistant professor for collaboration on the project. This plan, however, came to a deadlock in a short time, because contrary to my expectation the new colleague did not contribute towards the achievement of this project at all. Based on the deliberations and results obtained throughout the above-mentioned processes, we had started and continued the research, which made my heart beat first with joy, even under the following worst research environments. Without any direct research grants, in our spare time from main research projects (supported by some small grants) and that by only a womanpower (of M. N.), we had undertaken the dreamiest study on the chromatin conformation change code (4C) theory for gaining un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through numerous generations (cell divisions) [138]. At the start of this research, I had firmly resolved that my small research group (T. N. and M. N.) made some contribution (even if a sole and small concept) toward recent

rapid development of the life science field, which had already become a big science, as well as the space science, experimental elementary particle physics, geophysics, gravity physics/gravitational wave astronomy and others. On the other hand, it was also true that we had to principally continue our ordinary projects (most of which were supported by some small grants) [98-102, 109-113, 115-130, 132-143] and collaborations (with other groups) [131, 144-163], similarly to previous our own plus collaborative researches [62-90, 94, 103-108, 114, 164-187] with ~20 unlisted reviews and others (in English and in Japanese), since minimum research conditions were essential to execute the research on the 4C theory. Throughout whole research periods, of course, we had referred to numerous papers. Only about sixty selected randomly from them were first listed as references [1-61, 91-93] in this article, although many of these referred papers were cited in our previous papers [62-90, 94-130, 132-143]. If I had not come across this biological phenomenon, besides my failure in formation of excellent research group, my life as a scientist must be ended in imperfect combustion. Anyhow, we had started the exciting research on the 4C theory with sickening anxiety, quiet fighting spirit and a wish that it will be a forerunner in future biology (life science) beyond Biochemistry, Molecular Biology, Cell Biology and others, all of which have mostly targeted bio-systems already acquired during evolution since the birth of living things, under a following motto hidden inside:

Exploring only one, pursuing only it, by only one, and Creator of concept, rather than only one, rather than number one

The 4C theory, which is the fruit of my small serendipity and the outcome of some 10 years of intellectual investigation spirit, intuition plus insight (of T. N.) and enthusiasm, outstanding technique plus diligent work (of M. N.), is a creative concept on ability to gain un-programmed and new cell functions for the life conservation of higher eukaryotes.

In 2015, we published the 1st Edition of my retirement commemorative monograph (Monograph) at my own expense [138]. As shown later, of the Monograph, Chapters-2, 3, 4 and 5 respectively described results obtained from 1st HDAC2(-/-), Pax5(-), 2nd HDAC2(-/-) and NotchIP assay, Chapter-6 first proposed the 4C theory, Chapter-7 all-inclusively reviewed the 4C theory, and Chapter-8 mentioned my personal memoirs. In 2017, we published the 1st Editions of five original articles and one review article (Papers-1 ~ 6) on the 4C theory [188-193], respectively, which were the revised Editions of Chapters-2 ~ 7 of the 1st Edition of the Monograph (published in 2015) [138]. In addition, we published three review articles (RT/BR/208, RT/BR/209 and RT/BR/210) in Current Topics in Biochemical Research (Reviews-1, 2 and 3) [194-196], all of which reviewed the studies in the Papers-3, 4 and 5, respectively [190-192]. In 2018, we published the 2nd Edition of the Monograph (the 1st Edition was published in 2015) [138] and the 2nd Editions of the Papers-1 ~ 6 on the 4C theory (the 1st Editions were

published in 2017) [188-193]. In 2019, we published the 3rd Edition [197] of the Paper-6 (Comprehensive Review) (the 2nd Edition was published in 2018) [193], and the 1st Edition of the Paper-7 (Memoirs: the memoirs of a life scientist) [198], which was a revised Edition of Chapter-8 of the Monograph (the 2nd Edition was published in 2018) [138]. In addition, I published the 1st Edition of the Paper-8 (Review in Japanese), which was the translated Edition of the 3rd Edition of the Paper-6 published in 2019, with some fundamental concepts on life phenomena [197]. In 2020, we published the 4th Edition [199] of the Paper-6 (Comprehensive Review) (the 3rd Edition was published in 2019) [197], and the 2nd Edition of the Paper-7 (Memoirs) [199], which was a revised Edition of Chapter-8 of the Monograph (the 2nd Edition was published in 2018) [138]. In addition, I published the 2nd Edition of the Paper-8 (Review in Japanese), which was the revised Edition of the 1st Japanese Edition published in 2019, and the translated Edition of the 4th Edition of the Paper-6 published in 2020, with some fundamental concepts on life phenomena [199]. All of Figures 1 ~ 23 used in this 4th Edition are identical with corresponding those in the 1st, 2nd and 3rd Editions of the Paper-6 [198, 223], all of which were selected from those shown in the 1st and 2nd Editions of the Monograph [138] and the Papers-1 ~ 5 (the above-mentioned five original articles) [188-192]. Because I did not explain in detail the contents of these Figures here, please look at the URLs (listed in Supplementary Remarks) of the above-mentioned Monograph, Papers and Reviews on the 4C theory.

In any case, because the 4C theory is the only visible and biggest evidence for me as a scientist, here, I would like to re-describe closely the 4C theory [192, 193] by referring our recent review articles [194-196]. Based on our new results [138, 188-197] and previous results [114, 117, 119, 121, 124], we first proposed an all-inclusive hypothetic way for gaining un-programmed and new cell function to diminish excessively accumulated IgM H- and L-chains by means of irreversible creation of varied chromatin structure plasticity of Pax5, Aiolos, EBF1 and OBF1 genes with epigenetic modifications in individual clones of HDAC2(-/-) DT40 mutants through numerous generations (cell divisions) during continuous cultivation [138, 192, 193, 196, 197, 199].

First, we supposed that environment change recognition receptor/site (ECRR/ECRS) participates in the recognition of accumulation of IgM H- and L-chains as an abnormal, unexpected and/or unfavorable environment change (and probably acts in part in the signal transduction concerning the accumulation to the chromatin structure) (Fig. 20). In addition, chromatin conformation (structure) change complex (4C) machinery, which is diverse and contains a member of each of HATs plus HDACs and other factors, is supposed to directly and irreversibly create plasticity of the chromatin structure surrounding proximal 5'-upstream regions of the above-mentioned specific transcription factor genes (and to act in part in the signal transduction) (Figs. 20 and 21). Using the ECRR/ECRS, the 4C machinery and other components, the chain reaction of response to the abnormal and/or unfavorable environment change (i.e., the

accumulation of IgM H- and L-chains) occurs as follows.

First, in wild-type DT40 cells, in which various members of HATs and HDACs are largely expressed [64, 92], the 4C machinery for each of Pax5, Aiolos, EBF1 and OBF1 genes probably contains HDAC2 as a deacetylation activity, a specific member of HATs as an acetylation activity and other factors (Fig. 22) [92, 98, 101]. However, the functions of both the ECRR/ECRS and the 4C machinery may be basically not necessary for DT40 cells, because the protein and mRNA levels of IgM H- and L-chains are very low as mentioned above.

The other hand, the case of HDAC2(-/-) mutants is very different from that of DT40 cells as follows. Repeatedly, large amounts of IgM H- and L-chains artificially synthesized by their excessively increased gene expressions induced by the HDAC2-deficiency (Fig. 9) are first accumulated probably within the endoplasmic reticulum of HDAC2(-/-) mutant cells (Figs. 18 and 19). Most of the accumulated IgM H- and L-chains (which probably exist as a high molecular weight complex with each other [121]) are gradually secreted into the cultivation media (i.e., the outside of cells), whereas a part of them is transported to the nuclear envelope but not the inside of nucleus, and kept at the peri-nuclear space at the early and later cultivation stages (Figs. 18, 19 and 20). The excessively accumulated immunoglobulin proteins at the peri-nuclear space bind to the ECRR/ECRS localized at the inner nuclear membrane (where hetero-chromatin is possibly located) (Fig. 20). After the ECRR/ECRS recognizes the accumulation of IgM H- and L-chains as an abnormal and/or unfavorable environment change, the signal concerning the environment change is genome-widely transmitted to the chromatin structure surrounding proximal 5'-upstream regions of numerous genes (probably existing on several distinct chromosomes) encoding transcription factors, chromatin-modifying enzymes, and related factors and enzymes. Following the initial signal transduction, a spontaneous unbalanced response to the abnormal and/or unfavorable environment change is consecutively and separately converged to the proximal 5'-upstream regions of Pax5, Aiolos, EBF1, OBF1 and other genes in individual clones of HDAC2(-/-) mutants (Fig. 22).

In all individual clones of HDAC2(-/-) mutants, at the very early stage of cultivation (just soon after their birth by gene targeting techniques), bulk conformation of the 4C machinery dramatically changes to remove or drastically reduce HAT activity (of the assumed member of HATs), attributed to the HDAC2-deficiency. Throughout the above-mentioned process and following continuous cultivation, the 4C machinery newly contains a different member of HDACs, a specific (same or different) member of HATs and other factors, and thereby becomes varied. The diversity of alterations in the chromatin structure surrounding proximal 5'-upstream regions of these transcription factor genes is preferentially attributed to varied acetylation and deacetylation levels of one or more of the five specific Lys residues (K9/H3, K14/H3, K18/H3, K23/H3 and K27/H3) at the N-terminal tail of histone H3 [191, 195] caused by the collaboration of proper member of HATs and HDACs in the protean 4C machinery. These

successive epigenetic modifications of K9/H3 and K27/H3 (and K14/H3, K18/H3 and K23/H3) with acetyl group lead to irreversible creation of distinct chromatin structure plasticity surrounding proximal 5'-upstream regions of Pax5, Aiolos, EBF1 and OBF1 genes (Figs. 13, 14, 15, 16, 17 and 22). In generally, the chromatin structure surrounding proximal 5'-upstream regions possessing hyper-acetylation levels of one or more of the specific Lys residues of histone H3 is in the loose (open) form based on its no binding ability to DNA, but the chromatin structure surrounding proximal 5'-upstream regions possessing hypo-acetylation levels of one or more of the specific Lys residues of histone H3 is in the tight (closed) form based on its binding ability to DNA (Figs. 17, 21 and 23). By contrast, probably, the 4C machinery cannot change the chromatin structure surrounding ORF regions (coding regions) of these specific transcription factor genes by much [191, 192, 195-197]. As the need arises, the transcription factor complex (TFC) machinery (which consists of RNA polymerase, proper transcription factor(s), certain members of HATs and HDACs and other factors) can bind to promoter regions (or elements) within the loose (but not tight) form of the chromatin structure surrounding proximal 5'-upstream regions of Pax5, Aiolos, EBF1 and OBF1 genes (which have become latently active state), and thereby initiates their gene expressions (Figs. 21 and 23).

As a concrete result, tested individual clones (cl.2-1, cl.2-2, cl.2-3, cl.2-4, cl.2-5 and cl.2-6) of HDAC2(-/-) mutants gain the same, un-programmed and new cell function to reduce increased gene expressions of IgM H- and L-chains (resulting in their decreased protein levels) in almost the same changing pattern through increased or decreased gene expressions of Pax5, Aiolos plus EBF1 or OBF1 in three distinct ways (i.e., OBF1-dependent type, Pax5- and Aiolos-dependent type, and Pax5-, Aiolos- and EBF1-dependent type) through numerous generations during simple continuous cultivation under the same conditions (Fig. 10) [121, 138, 188, 190, 194]. Such distinct ways are not under the control of HDAC2 but originally based on irreversible creation of their distinct chromatin structure plasticity with epigenetic modifications during continuous cultivation [138, 190-199]. Remarkably, these tested six individual clones of HDAC2(-/-) mutants classified into the above-mentioned three different cell types should exhibit more distinct functions, respectively. Because they show three dissimilar ways for gene expressions of transcription factors and chromatin-modifying enzymes (such as Pax5, Aiolos, EBF1, OBF1, Blimp1, PCAF, HDAC9, etc.), all of which are easily considered to participate in gene expressions of numerous different genes, besides those of IgM H- and L-chains at the later cultivation stage (Figs. 1 and 10). For instance, in DT40 cells, Pax5 regulates dramatically or moderately gene expressions of PCAF, HDAC7, HDAC9, Aiolos, OBF1, Ikaros, E2A, EBF1 and PU.1, and further Pax5 isoforms A and B regulate differentially other B cell development-related factors [134, 138, 189]. In addition, Aiolos regulates pre-mature B cell apoptosis mediated by BCR signaling [141]. E2A regulates gene expressions of survivin, IAP2 and caspase-8 [140]. Helios regulates the gene expression of protein kinase Cs [139]. EBF1 regulates dramatically gene expressions of Blimp-1 and protein kinase C θ [129,

142]. Besides, in one or more of these six mutant clones, respective gene expression patterns of PCAF, HDAC7, HDAC9, Ikaros and OBF1 are spontaneously and complicatedly reversed in the middle of simple continuous cultivation, although those of Pax5, Aiolos, EBF1, E2A, PU.1 and Blimp-1 do not change while passing from the early to later cultivation stages. Furthermore, if other individual mutant clones obtained [190, 194] are analyzed, there is a possibility that additional distinct cell types might exist. These results indicate that there are many distinct ways (molecular mechanisms) for HDAC2(-/-) mutants to newly gain a same un-programmed cell function as if there exist many distinct mountain trails to climb the top of Mt. Fuji. Consequently, individual clones of HDAC2(-/-) mutants acquire flexible, elastic and pluri-potential ability not only to adapt to an abnormal, unexpected and/or unfavorable environment change in distinct ways but also to branch off into diverse derivative cell types, which may exhibit varied characteristics (functions), even though they are originally the same cell type and the established (or tumor) cell line [192, 193, 196, 197, 199].

The above-mentioned way that HDAC2(-/-) DT40 mutants gain un-programmed and new cell function to diminish excessively accumulated IgM H- and L-chains by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through numerous generations during continuous cultivation is obviously different in molecular mechanisms from the so-called endoplasmic reticulum (ER) stress response/unfolded protein response as quality control to eliminate defective products and others. Naturally, the way to eliminate unnecessary products (large amounts of IgM H- and L-chains) is clearly different from the many well-known life phenomena, e.g., the alterations in the colors of snowshoe hare's hairs and ptarmigan's feathers between summer and winter seasons, and the mimicries of *Philomachus pugnax* and *Papilio polytes* probably based on supergenes, because these are programmed functions in response to environment change and their related genome information is unchanged.

We expanded the above-mentioned hypothetic way (the 4C theory) on the exclusion of IgM H- and L-chains excessively accumulated in HDAC2(-/-) mutants to a universal hypothetic concept on the way for gaining un-programmed and new cell functions by means of irreversible creation of varied chromatin structure plasticity surrounding proximal 5'-upstream regions of specific transcription factor and chromatin-modifying enzyme genes with epigenetic modifications through numerous generations (cell divisions) in higher eukaryotes. The details of the universal concept on the 4C theory are as follows.

When higher eukaryotic cells firstly encounter an intra- and/or extra-cellular environment change in their lives, they gradually acquire the ability to adapt themselves to the new environment change or exclude it (if unnecessary, abnormal, unexpected, uncomfortable and/or painful) through numerous generations (cell divisions). Using the ECCR/ECRS, the 4C machinery and other components, the eukaryotic cells start a chain reaction of response to the new environment change as follows (Figs. 20 and

21). First, the new environment change is recognized by means of the ECRR/ECRS, which may be localized nearby the nuclear membrane as a nucleus-cytoplasm barrier (probably at the inner nuclear membrane where hetero-chromatin is possibly located) (Fig. 20). There is a possibility that putative specific molecule(s) acts as the intermediary sensor at this step to recognize the new environment change. Naturally, there is certainly another possibility that ECRR/ECRS should be newly created through gathering of its components based on the fluid mosaic model of membrane, if they exist separately within the lipid bilayer membrane. Next, putative signal(s) concerning the new environment change is genome-widely transmitted to the chromatin structure within the nucleus through numerous generations (cell divisions). Following the initial acceptance of the putative signal(s), the 4C machinery induces a slight alteration in the chromatin structure surrounding proximal 5'-upstream regions of numerous genes encoding chromatin-modifying enzymes, transcription factors, and related enzymes and factors with epigenetic modifications. The transduction of the signal(s) and spontaneous unbalanced response to the new environment change are successively repeated and converged into the restricted chromatin structure surrounding proximal 5'-upstream regions of various specific members of the above-mentioned factor and enzyme genes. Finally, this successive signal transduction concerning the new environment change causes various epigenetic modifications of histones and/or DNA within the restricted chromatin regions of the aforesaid specific genes with acetyl, methyl, phosphate, ubiquitin and ADP-ribose groups and/or others. Of these various epigenetic modifications, acetylation and deacetylation of several specific Lys residues of core histones H2A, H2B, H3 and H4 may be the major ones.

The 4C machinery, which consists of a specific member of each of the HATs and HDACs and other factors, preferentially participates in the acetylation and deacetylation among such epigenetic modifications by rights (Fig. 21). Positions of specific Lys residues and kinds of core histones are diverse. For instance, in the above-mentioned case [190-199], the acetylation and deacetylation of Lys-9, Lys-14, Lys-18, Lys-23 and Lys-27 residues of histone H3 are prominent. Consequently, epigenetic modifications of one or more of these specific Lys residues of histone H3 with acetyl group change within the chromatin structure surrounding proximal 5'-upstream regions of the above-mentioned target genes through numerous generations. Varied functions of the protean 4C machinery on such acetylation and deacetylation levels are mainly based on distinct combinations of each member of HATs and HDACs as the components, because any HAT and HDAC members' own activities are probably unchangeable. By contrast, the protean 4C machinery cannot alter the chromatin structure surrounding ORF regions (coding regions) of corresponding genes by much. The binding ability of the N-terminal tail of histone H3 to DNA is tentatively and qualitatively deduced from the acetylation and deacetylation levels of one or more of the specific Lys residues, though which Lys residue(s) really and/or mainly participates in the binding is still undefined. That is, hyper- (high) or hypo- (low or no) acetylation levels induce no binding or full binding ability, resulting in the loose (open) or tight (closed) form of the chromatin structure (Fig. 21).

Thus, chromatin structure plasticity is irreversibly created based on successive conformation changes with epigenetic modifications. These ways to gradually and tardily create chromatin structure plasticity for gaining un-programmed and new cell functions are obviously different from the ways to immediately and rapidly cause the chromatin conformation change for expressing programmed and ordinary cell functions.

Whenever the need arises, the TFC machinery can bind to promoter regions (or elements) within the loose form of the chromatin structure surrounding proximal 5'-upstream regions of the above-mentioned specific target genes (which are in the latently active state), followed by initiation of their gene expressions (Fig. 21). By contrast, the TFC machinery cannot bind to promoter regions (or elements) within the tight form of the chromatin structure surrounding proximal 5'-upstream regions of un-target genes (which are in the latently inactive state), and thereby cannot initiate their gene expressions. Consequently, the loose or tight form of the chromatin structure surrounding proximal 5'-upstream regions causes high or low (or no) transcription levels of corresponding genes (Fig. 23). Notably, there is a great possibility that the ways for irreversible creation of chromatin structure plasticity are varied among individual cells of the same type, even though the new environment change and signal(s) on the environment change are the same for all of them. That is, in order to gain un-programmed and new cell function(s), individual cells possess the ability not only to complicatedly and diversely alter the chromatin structure surrounding proximal 5'-upstream regions of various specific genes but also to separately alter the chromatin structure of the same proximal 5'-upstream region of a certain gene into varied forms. Thus, gene expressions of the specific chromatin-modifying enzymes and transcription factors diversely alter among individual cells through numerous generations, despite the same environment change. As a result, individual somatic cells of the same type can newly gain the same and/or distinct un-programmed cell function(s) in different ways, in order to accommodate themselves to a new environment change.

In conclusion, in order to gain un-programmed and new cell function(s), somatic cells (and even tumor cells) of higher eukaryotes become pluri-potent, elastic and flexible, all of which basically originated from pluri-potency, elasticity and flexibility of the chromatin structure. That is, in order to adapt to an intra- and/or extra-cellular environment change, individual somatic cells of higher eukaryotes possess the ability to newly gain the same and/or distinct un-programmed cell function(s) in different ways by means of irreversible creation of varied chromatin structure plasticity with epigenetic modifications, i.e., from the loose to tight forms or vice versa of the chromatin structure surrounding proximal 5'-upstream regions of specific transcription factor and chromatin-modifying enzyme genes (Fig. 23). Such a loose or tight form of the chromatin structure surrounding the proximal 5'-upstream region is in the latently active or inactive state for transcription of the corresponding gene, although the proximal 5'-upstream region as mere nucleotide sequences is in the silent state for expressions of most of the genome functions. Creation of varied chromatin structure plasticity in individual somatic cells is

triggered by the spontaneous unbalanced response to the new environment change when they firstly encounter with it and then irreversibly accomplished by the successive unbalanced convergence of the response through numerous generations (cell divisions). The different ways to create varied chromatin structure plasticity in individual somatic cells are certainly dependent on their antecedents. Moreover, chromatin structure plasticity, regardless of whether its creation is in the course or was already completed, is inherited to descendant generations associated with or without additional conformation change through cell divisions. Remarkably, irreversible creation of chromatin structure plasticity occurs in descendant cells but not in the cell which initially accepts the signal on the environment change, although reversible regulations of ordinary gene expressions and enzyme reactions occur in the cell itself which accepts proper signal (Fig. 22). Probably, irreversible creation of chromatin structure plasticity, with the intention of adapting to the environment change, occurs inevitably but not incidentally and/or neutrally.

The proximal 5'-upstream chromatin region (as the loose or tight form) directs the switch (on or off) for latent transcription ability of the corresponding gene by means of irreversible creation of chromatin structure plasticity; therefore, the 5'-upstream chromatin region can be regarded as "director for gene expression" from a functional point of view, besides as "notch of chromatin" from a structural point of view as mentioned above. The notch (or director) covers specific nucleotide sequences of transcriptional elements (such as promoter, operator, enhancer, etc.) and their neighboring nucleotide sequences. Remarkably, the real recipient of the signal concerning the environment change may be just the steric/three-dimensional (as if loose rope-like) chromatin structure possessing self-conversion ability of conformation (we named here), which changes dynamically between the loose and tight forms, but not mere the linear/one-dimensional (as if rod-like) chromosome structure and the planar/two-dimensional (as if thread-like) chromatin structure, the two of which are in the unchangeable static state. That is, the chromatin structure of proximal 5'-upstream region (notch or director), as a dynamic and changeable three-dimensional conformation, possesses two fundamental abilities, i.e., to receive the signal concerning the intra- and/or extra-cellular environment change and to direct the switch (on or off) for latent transcription ability of the corresponding gene by means of its irreversible chromatin conformation change that is in response to the signal. The self-conversion ability of conformation, which is based on the 4C theory to irreversibly create plasticity of three-dimensional chromatin conformation (structure) in higher eukaryotes, is obviously different from the well-known rearrangement of antibody genes and others, because these are somatic recombination (rearrangement) of DNA structures (base/nucleotide sequences) exactly. Naturally, concrete data are not enough to support these ideas. As a result of these complicated biological events, higher eukaryotes acquire pluri-potential, elastic and flexible ability to create diverse derivative cell types possessing varied characteristics/functions (e.g., varied component cells of diversely differentiated organs) from the same type of somatic cells in distinct ways, in order to accommodate themselves to new intra- and/or extra-cellular environment. In some cases, even

neighboring cells derived from the same cell type are possibly regarded as extra-cellular environment for themselves. Moreover, in higher eukaryotes diverse kinds of somatic cells share and express orderly and cooperatively vital functions with each other. For concrete typical examples, 1) there are innumerable cell types that make up the brain and nerves. In addition, there are other examples for the maintenance of homeostasis that is also one of the vital functions as follows. 2) hypothalamus synthesizes and secretes corticotropin-releasing hormone (CRH), growth hormone-releasing hormone (GHRH), luteinizing hormone-releasing hormone (LHRH), and thyrotropin-releasing hormone (TRH), 3) anterior pituitary synthesizes and secretes growth hormone (GH), luteinizing hormone (LH) and adrenocorticotrophic hormone (ACTH), and 4) posterior pituitary synthesizes and secretes antidiuretic hormone (ADH), vasopressin and oxytocin (OT). Moreover, 5) A (α), B (β), D (δ) and PP cells of Langerhans islet of pancreas synthesize and secrete glucagon, insulin, somatostatin and pancreatic polypeptide, respectively.

Finally, as mentioned above, we named our hypothetic theory on such a bio-system that gains un-programmed and new cell function(s) by means of irreversible creation of chromatin structure plasticity with epigenetic modifications, and is one of the most fundamental and important ways for life conservation and cell type determination of higher eukaryotes, as the chromatin conformation (structure) change code (4C) theory [138, 192, 193, 196, 197, 199]. By the way, here, I want to briefly explain on the simple concepts and molecular mechanisms of the two well-known code theories (i.e., the genetic code and the histone code), which are closely correlated to chromatin (or DNA) functions (especially gene expression/transcription) but different from the 4C theory, because I did not mention about these two until now in this article at all. First, the genetic code is the biochemical basis of heredity and is based on codon (triplet: three consecutive base/nucleotide sequences) in both DNA and mRNA, and the codons determine the specific amino acid sequences in proteins (and polypeptides) at the translation step on the ribosome. The translation/protein synthesis occurs through the base pair formation by the hydrogen bond between the codon in mRNA and the anticodon in aminoacyl-tRNA. Of course, the genetic code appears to be uniform for nearly all known organisms; therefore, in life phenomena it is one of the most fundamental and important concepts, as well as the double helix structure of DNA. Second, the histone code, which is an epigenetic code as well as DNA methylation and others, will be explained a little more, since it is a bit like the 4C theory apparently. The histone code is a hypothesis that the transcription of genetic information encoded in DNA is in part regulated by chemical modifications of histones within their flexible N-terminal tails protruding from the nucleosome. Namely, as well known, many of histone modifications obviously participate in chromatin structure change and thereby change expression levels of numerous genes significantly. The critical concept of the histone code hypothesis is that the histone modifications serve to recruit other enzymes and/or factors by specific recognition of the modified

histone (as a mark/modifier) via protein domains specialized for such purposes, rather than through simply stabilizing or destabilizing the interaction between the histone and the specified DNA region. These recruited enzymes and/or factors then act to alter chromatin structure actively or to promote transcription. In other words, the molecular mechanism of the histone code is based on the facts that histones are modified (marked) by several small molecules, and the mark summons (or is recognized and accessed by) various enzymes and/or factors, which participate in the expression of various chromatin functions (especially gene expression). Finally, I repeat the description on the 4C theory with a slight modification as follows. The molecular mechanism of the 4C theory is based on that the plasticity of proximal 5'-upstream chromatin regions of various genes are irreversibly created by the 4C machinery through epigenetic modification with acetyl group (and others), and the loose (open) or tight (closed) form of chromatin structure summons or rejects (or is recognized and accessed by, or is reversely un-recognized and un-accessed by) various transcription factors and/or chromatin-modifying enzymes, which participate in the expression of numerous chromatin functions (including gene expression). Therefore, needless to mention again, regarding the expression of genome (chromatin) functions including the gene expression, the 4C theory obviously functions upstream both the histone code and the genetic code. Probably, the supposed number of codes in the 4C theory, which determines complicated and varied characteristics (functions) of higher eukaryotic cells, can be roughly estimated based on the combination (multiplication) of the number of candidate genes and that of codes for each of these genes. The most influential candidates are various specific genes encoding transcription factors, chromatin-modifying enzymes, and related factors and enzymes, all of which are necessary for gaining varied cell functions and specificities of higher eukaryotes. The number of codes for each of these candidate genes is probably two. This is because the proximal 5'-upstream chromatin region (the loose or tight form), as latently active or inactive state, directs the switch (on or off) for transcription ability of the corresponding gene. On the other hand, almost all programmed, complicated and diverse cell functions, which were acquired through a long evolutionary process, are orderly and systematically expressed throughout their development and differentiation. Naturally, the 4C theory is suitable as an explanation for the development and differentiation (and so-called robustness) of higher eukaryotes, because action of putative signal(s) concerning environment changes seems to fairly resemble that of certain players (such as hormone, cytokine, nerve-transmission substance, etc.), which participate in cell-cell, tissue-tissue and/or organ-organ interactions (communications) through these two fundamental life phenomena.

Here, I would like to change the topic from the 4C theory to recent research results on important life phenomena. First, I wanted to describe briefly my impressions on genome editing techniques, which have been recently developed in life science field very rapidly and widely. Because the techniques make

impacts of right and wrong for mankind without fail, all of natural science, social science and humanities must cooperatively solve the troublesome problems, which must obviously occur now and in near future. However, I did not comment further on the techniques like recombinant DNA techniques. By the way, from old to present, a lot of ways to newly acquire cell functions, which are somewhat similar with the 4C theory apparently, are being actively studied by many scientists in various organisms (prokaryotes and eukaryotes) as follows [200-204]. There are the experimental evolutions on phenotypic changes based on mutations (genome information changes) through enormous generations during long continuous cultivation and breeding periods in *Escherichia coli* (*E. coli*) and *Drosophila melanogaster* (*D. melanogaster*). For example, *E. coli* cells propagated for numerous generations obtained the property to use citrate in media as a nutrient, although those cannot usually use citrate as a nutrient source. In addition, the temperature adaptability of two natural populations of *D. melanogaster* living in two areas with different annual temperatures was obviously different from each other after bred under the same temperature environment.

Recently, the very important life phenomena (e.g., inter-tissue communication, intestine-to-germline transmission of epigenetic information and soma-to-germ communication, etc.), which are closely related to alterations in chromatin structure, are being rapidly researched in *Caenorhabditis elegans* (*C. elegans*), *D. melanogaster* and others [205-220]. Here, by borrowing a part of description in the remarkable paper [205], I mention briefly recent achievements on these important life phenomena as follows. Epigenetic modifications of chromatin structure (such as methylation, acetylation and others of histones H2A, H2B, H3 and H4) act as intermediary between environment and gene expression, followed by participation in various organismal phenomena. The results obtained are as follows. For example, in *C. elegans* the alterations in epigenetic states caused by the deficiency of the histone H3K4me3 modifier ASH-2 in the intestine or germline increase organismal stress resistance abrogated by knockdown of the H3K4 demethylase RBR-2. The increase in stress resistance induced by the ASH-2 deficiency in the intestine is abrogated by RBR-2 knockdown in the germline, suggesting the intestine-to-germline transmission of epigenetic information. This communication from intestine to germline in the parental generation increases stress resistance in the next generation. In addition, in various species the epigenetic alterations induced by genetic manipulation and environmental trajectory can sometimes be taken over through many generations, even when the first stimulus no longer exists, e. g., the increased stress resistance induced by low-level stress exposure during development in *C. elegans* can be transmitted to the descendant generations through soma-to-germ communication. Remarkably, the systemic regulation of epigenetic information by means of small RNAs is reported to be essential for transgenerational epigenetic inheritance, which induces phenotypic alteration in descendant generations. These results, together with others, revealed that inter-tissue communication of epigenetic information provides mechanisms for intergenerational regulation of systemic stress resistance, indicating that acquired

characteristics (traits) of somatic cells of higher eukaryotes are intergenerationally inherited into descendant generations.

Returning to the original subject again, I listed below with partial repeat both the general concepts of the 4C theory and the unsolved crucial issues on the exclusion of excessive IgM H- and L-chains in bullet points. Outline of the 4C theory, which is one of the most fundamental and important ways for life conservation and cell type determination of higher eukaryotes, is concretely as follows. 1) Somatic cells of higher eukaryotes possess the pluri-potency, elasticity and flexibility, which are basically originated from those of the chromatin structure, for gaining un-programmed and new cell functions, in order to cope with and/or overcome environment change. 2) Somatic cells gradually acquire the un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity surrounding the proximal 5'-upstream regions of specific transcription factor and chromatin-modifying enzyme genes through numerous cell divisions (generations). 3) Chromatin structure plasticity (from the tight to loose forms or vice versa) is continuously and irreversibly created based on the chromatin conformation change with epigenetic modifications through numerous cell divisions (generations). 4) Diversity of chromatin structure plasticity in individual somatic cells of the same type is triggered by spontaneous unbalanced response to the environment change and thereafter accomplished by its successive convergence through numerous cell divisions (generations). 5) Irreversible creation of chromatin structure plasticity depends on both antecedents of somatic cells and successive response to the environment change, and completes in descendent cells but not in the cell which initially meets with the environment change. 6) Irreversible creation of chromatin structure plasticity probably occurs inevitably but not incidentally and/or neutrally. 7) The environment change is recognized by putative environment change recognition repetitor/site (ECRR/ECRS), and chromatin structure plasticity is irreversibly and directly created by putative chromatin conformation change complex (4C) machinery. 8) The chromatin structure of the proximal 5'-upstream region(s) of specific transcription factor and/or chromatin-modifying gene(s), as dynamic and changeable three-dimensional conformation, receives the signal on the environment change. 9) The chromatin structure (the loose or tight form) of the proximal 5'-upstream region of the specific gene directs the switch (on or off) for its latent gene expression ability. 10) The supposed number of codes in the 4C theory, which determines both complicated cell functions and diverse cell types of higher eukaryotes, may be estimated based on combination of the number of candidate genes and that (probably two) of codes for each of these specific candidate genes.

There are several crucial issues to be clarified in the 4C theory on the exclusion of excessive IgM H- and L-chains artificially accumulated in HDAC2(-/-) DT40 mutant cells [138, 188-199, 221-225]. 1) Despite the HDAC2-deficiency, why do the acetylation levels of one or more of K9, K14, K18, K23 and K27 residues of histone H3 within the chromatin structure (of ~10 nucleosomes) surrounding proximal

5'-upstream regions of Pax5, Aiolos and EBF1 genes decrease at the early stage of cultivation in HDAC2(-/-) mutants? 2) Why do the decreased acetylation levels of one or more of the specific Lys residues of histone H3 within proximal 5'-upstream chromatin regions of these three genes increase during cultivation? 3) Why do the acetylation levels of one or more of K9, K14, K18, K23 and K27 residues of histone H3 within the chromatin structure (of ~10 nucleosomes) surrounding the proximal 5'-upstream region of the OBF1 gene decrease during cultivation? 4) Which Lys residue(s) of K9, K14, K18, K23 and K27 of histone H3 is really and/or mainly involved in its binding to DNA within proximal 5'-upstream chromatin regions of Pax5, Aiolos, EBF1 and OBF1 genes must be determined. 5) Functional and steric differences between the loose and tight forms (based on hyper- and hypo-acetylation levels of one or more of the specific Lys residues of histone H3) of the chromatin structure surrounding proximal 5'-upstream regions of these four genes must be clarified more precisely. 6) Why do the changes in acetylation levels of the specific Lys residues of histone H3 for individual transcription factor gene during cultivation differ among individual mutant clones? 7) Why do the changes in acetylation levels of the specific Lys residues of histone H3 for individual transcription factor genes during cultivation differ within individual mutant clone? 8) How does the 4C machinery (which acts in irreversible creation of chromatin structure plasticity of proximal 5'-upstream region) differ from the well-known chromatin-modifying machinery, although the TFC machinery (which acts in the transcription of ORF region of the corresponding gene) may be the same as the well-known transcription machinery? 9) To demonstrate the 4C theory, it is essential to establish both the putative ECRR/ECRS as the first player to recognize the environment change and the putative 4C machinery as the final player to directly and irreversibly create chromatin structure plasticity.

In connection with the 4C theory, here, I wish to briefly express my personal interpretations concerning the conception of complicated and diverse life phenomena. First, I would like to re-describe the issues mentioned in the first part of this personal article on the Memoirs/Insights with a slight modification as follows. Higher eukaryotic cells possess four possible countermeasures to cope with and/or overcome an abnormal, unexpected and disadvantageous environment change, when they firstly encounter it in their lives. 1) When cells encounter a very severe change, they die because it is far ahead of their adaptation ability. 2) When cells encounter a considerably severe change, they cope with or overcome it by means of alterations in genome information, such as point mutation, insertion, deletion, duplication and multiplication on the DNA molecule through numerous generations. 3) When cells encounter a moderate change, they cope with or overcome it by means of irreversible creation of chromatin structure plasticity caused by successive chromatin conformation changes with epigenetic modifications through numerous generations (cell divisions). 4) When cells encounter a minor change, they respond to it only through the already acquired regulation mechanism(s). As described above, IgM

H- and L-chains artificially and excessively accumulated in HDAC2(-/-) DT40 mutants (and in Pax5(-) DT40 mutants) are dramatically decreased through numerous generations (cell divisions) during continuous cultivation. Although the above-mentioned classification is in my own way, the aforesaid remarkable biological event, which is a typical example of the most fundamental and important ways for adaptation to the environment change and for life conservation of higher eukaryotes, seems to be really an example of the third case of the above-mentioned countermeasures [138, 188-199, 221-225].

In addition, I have a following harmless insight about the development and differentiation of higher eukaryotes, which are certainly and closely related to irreversible creation of chromatin structure plasticity. For instance, when ovum and sperm are fertilized, all components of the chromatin (or chromosomes) (except DNA) existing in the former are generally considered to be evenly and symmetrically divided into two chromatin sets. However, in fact, an asymmetrical allotment of the chromatin components must spontaneously become in a few unbalances; i.e., two daughter cells after the first cell division of fertilized ovum have already somewhat of quantitative and qualitative differences in the constituent elements of their chromatin (chromosomes). Successively, the unbalanced distribution of the chromatin components is repeated and converged to create varied chromatin structure plasticity surrounding proximal 5'-upstream regions of different sets of numerous specific genes in individual cells through numerous cell divisions. Furthermore, such a spontaneous and asymmetrical allotment of the chromatin components at first cell division (followed by its successive convergence) universally occurs in all fertilized ova. Consequently, the complexity and diversity of chromatin structure plasticity cause those of cell specificity, i.e., distinct functions of the chromatin, nucleus and/or cell, leading to irreversible creation of variety of specific cell (or tissue) types of higher eukaryotes.

Well, genome information, which is inherited to descendant generations through cell divisions (and naturally, fertilization or pollination of generative cells), is generally preserved in nucleotide sequences of DNA and translated into amino acid sequences of protein (or polypeptide) through transcription into nucleotide sequences of mRNA. In eukaryotes, as mentioned above, both the genome information and the nuclear function are mainly protected by the nuclear membrane that acts as a cytoplasm-nucleus barrier; naturally, there are two other typical and important barrier systems for conservation (protection) of life, e.g., a blood-brain barrier for conservation of individuals (and/or maintenance of homeostasis) in higher animals and an insemination (sperm-ovum fertilization or stamen-pistil pollination) barrier for conservation of species in higher animals or plants. Moreover, both the communication and the signal transduction between the nucleus and cytoplasm, which are essential for normal expressions of cell functions, are preferentially performed by the to and fro passage of large and/or small molecules via the nuclear pore that acts as a guard station at the barrier. By contrast, both the transport of useless molecules and the signal transduction concerning an abnormal, unexpected and/or unfavorable change in the intra- and/or extra-cellular environment to the nucleus are usually shielded by this barrier system.

Hereafter, I want to largely digress from the subject concerning researches, to change it to my personal experiences, interpretations and opinions, and then to randomly mention each of these as follows. First, I organized “The 9th MBSJ Spring Symposium in Miyazaki” held on 2009 (MBSJ: The Molecular Biology Society of Japan). The theme was “New Quickening of Molecular Biology ~ Its Dawn of Daybreak from Miyazaki (in Japanese) ~”. As a matter of fact, I secretly and individually included the meaning of the 4C theory, which was still immature and unpublished at that time, in the theme of the formal symposium. Anyhow, I am not a little proud of the 4C theory [138, 188-199, 221-225] besides our other researches concerning histones, histone chaperones, chromatin-modifying enzymes and transcription factors in the DT40 cell line by gene targeting techniques [94-130, 132-137, 139-143]. It is because all these studies were originally started and achieved only by myself and my small group, except some technical helps, and not directly succeeded to those of any related scientists and/or groups. Particularly, I have a credit for the 4C theory, a new fundamental concept on life phenomena, which was created based on a few and insufficient data in our solitary step, i.e., by a very small group (M. N. and T. N.), in spare time from our other main projects and without any research grants.

Incidentally, as easily perceived by the above-mentioned facts, prior to the publication of the first paper on the iPS cells, in fact, we had started the research on the 4C theory concerning the exclusion of excessive IgM H- and L-chains artificially accumulated in HDAC2(-/-) DT40 mutants through numerous generations during simple continuous cultivation. However, while I was in active service, all of data on the 4C theory for gaining un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity could not be published as papers in formal Journals and presentations in any Conferences/Congress, Symposium and Meeting. Because I thought it was the minimum requirement that besides all our results obtained by that time, more additional data should be necessary to publish at the same time, in order to win recognition of scientist community. Therefore, we first published the data concerning the 4C theory as the retirement commemorative monograph (Monograph) at my own expense in 2015, instead of original papers in formal Journals. Further, when I retired, to my regret, nobody took over the research concerning the 4C theory. And I had to abandon all my own old samples, finally submitted manuscripts and Experimental Notes (containing raw data, most of which had been already published), which were kept from graduate school (except some samples and almost all data of M. N. concerning the 4C theory), because I lost both the academic and private spaces to keep them. It goes without saying that my colleagues probably keep their own samples and raw data concerning our other research projects. On my thorough retirement, for the first time, as mentioned above, I had seriously started writing of rough drafts and manuscripts for most of Chapters of the Monograph on the 4C theory in the form of original papers.

Meanwhile, concerning the acquisition of the pluri-potency of higher eukaryotic cells, following the iPS cells, the analogous cells triggered by simple stress of various reagent-treatments were reported in rapid succession. Nevertheless, to my great surprise, two papers on the STAP cells were splendidly and showily reported toward the end of the writing of our manuscripts on the 4C theory. However, after a little while these two papers were retracted and thereby the existence of the STAP cells came to naught. Regrettably and sorrowfully, the so-called STAP scandal was degraded itself to one of the worst injustices in the whole science world, and thereby the Japanese science world, especially the Japanese life science field, was got a violent blow. Naturally, I was afraid that the scandal must have indirect effects (such as severe criticisms of reviewers) on our research concerning the 4C theory not a little, while it probably has not direct effects at all.

Anyhow, regardless of the STAP scandal, I decided to publish the Monograph at my own expense in the worst case as a scientist out of office at least, since I could not obtain additional data anymore and thereby feared that our papers should be rejected by severe referees in the peer review process, when those are submitted. Based on these my own experiences, concerning a means for further efficient development of science, I should like to express my brief opinion as follows. When scientists retire thoroughly, I guess that most of them have a lot of certain but immature and unpublished data, which are taken over by nobody. In order to prevent that the enormous unquestionable data remain unpublished and thereby come to naught secretly, it is necessary for scientist community to make suitable resorts for publishing such unpublished data (of course, their reliabilities are essential) as the gemstone with a hidden potential, besides so-called formal Journals, in which mature results are published as papers and/or reports in complete forms, because open-access Journals of bad reputation rapidly go on increasing now. Although this idea may be contrary to the ordinary opinion of the community and the general public now, especially after the historic STAP scandal and other injustices, the real reason why I thought the idea is as follows. If this idea of mine can be realized, in future, someone is able to initiate and develop new research projects using some of these authentic data just as the start line without waste of time, effort and grant, even though those are the mixture of wheat and chaff, because of incomplete and immature publication.

Here, I wished to change the subject to life scientist's types (or styles) having no direct connection with the 4C theory. Since old times, roughly speaking, there are three typical researcher's types (or styles) based on means of taking influential and important data in life science field as follows. The first type is a category of researchers who find new biological concepts and/or phenomena based on morphological nature observation, like C. R. Darwin. Most of primate biologists and ecologists (in fieldwork) are classified into this category. The second type is a category of researchers who find new biological concepts, phenomena, functions and/or molecules based on their own experimental results, like G. J. Mendel. Most of experimental biologists (in wet laboratory) are classified into this category. The

third type is a category of researchers who find new biological concepts, phenomena, functions and/or molecules based on data (results) of other scientist(s), like J. D. Watson and F. H. C. Crick, and M. Kimura. Most of molecular evolutionists and informatics biologists (in dry/silico laboratory) are classified into this category. Of course, I belonged thoroughly to the second category; therefore, I could not add any more data to the 4C theory after my thorough retirement. To return from the digression, I should like to emphasize once again that these facts were just the reason and/or the excuse for the above-mentioned decision, i.e., I decided to publish our data concerning the 4C theory as the Monograph at my own expense, followed by registration as the institutional repository (of University of Miyazaki), as an amateur or an old fossil-like and analog-type scientist out of office.

In 2021, on the describing the 3rd Edition of the Paper-7 of the Memoirs/Insights (the 1st and 2nd Editions were published in 2019 and 2020) [198, 223], I added following personal comments on the publication processes of Monograph, Papers and Reviews on the 4C theory (with repeating a part of the above descriptions). In 2015, I first negotiated with Springer Japan about the publication of the retirement commemorative monograph (Monograph) on the 4C theory at my own expense several times. However, the publisher could not consider my proposal, probably because it did not match with the guidelines, due to that all of data were unpublished yet and belonged only to my group. Therefore, in the same year, I immediately published the 1st Edition of the Monograph on the 4C theory through a small printing office in Miyazaki at my own expense. Thereafter, I registered the 1st Edition of the Monograph as the institutional repository and published (uploaded) it on Website, which is now available from following new URL (but not from old URL) [138]). Then, I sent the 1st Edition of the Monograph with my compliments to many Japanese scientists in the chromatin field and other life science fields and also informed old URL to many foreign scientists in the chromatin field soon after the registration. Unfortunately, contrary to my expectation, regardless of whether comments were positive or negative, there were no responses with the except for a slight comment. However, after a little while of the publication (uploading) of the Monograph concerning the 4C theory as the repository [138], as mentioned above, some scientists suggested that we have better to submit the contents concerning the 4C theory to certain ordinary/formal Journals. Following their suggestions, we prepared the six manuscripts of the five original and one review articles, all of which were the modified versions of Chapters-2 ~ 7 of the Monograph, in a formal paper form with a relatively simple description. Then, as the first attempt I successively tried to submit one of the manuscripts (the revision of Chapter-2) to two open-access Journals (F1000Research and bioRxiv). However, the two Journals were not able to consider our submission, because our manuscripts did not match with their guidelines (policies), i.e., in the case of F1000Research the raw data for important findings must be submitted at the time of submission and in the case of bioRxiv the content must be unpublished, respectively. The reasons why I could not answer

to their guidelines were as follows. 1) Almost all figures of the Monograph and the Papers-1 ~ 6 (the five original and one review articles) consisted of PCR product bands cut off from more than 5,000 (including ~200 of the Paper-1) original photographs of gel electrophoretic patterns (i.e., raw data) of ChIP and NotchIP assays and others. Unfortunately, we were not able to provide the raw data that F1000Research means, because all the equipment used were already production stoppage and went wrong, and their license could not be used now; therefore, we had only printed matters and useless data files of all original gel electrophoretic patterns even now. 2) As mentioned above, the data concerning the 4C theory had been already published as the Monograph and uploaded as the institutional repository available from new URL (but not from old URL) [138]; therefore, all contents in the manuscripts of the six articles (Papers-1 ~ 6) were regarded as the published ones by bioRxiv. These facts strongly suggested that any ordinary/formal Journals should not accept our manuscripts of Papers-1 ~ 6 concerning the 4C theory. Therefore, in 2017, I decided to publish the modified versions of Chapters 2-7 of the Monograph concerning the 4C theory as the Papers-1 ~ 5 (original articles) and the Paper-6 (review article) with corrections in quite some detail and to register them as the institutional repositories, which are available from corresponding new (not old) URLs [188-193]. Fortunately, at length we had the opportunity to publish the researches on the 4C theory (especially the researches in Refs. 190-192) as three review articles in *Current Topics in Biochemical Research* (Volume 18) in 2017 [194-196]. Consequently, in 2018, we published the 2nd Editions of the Monograph and the Papers-1 ~ 6, and thereafter registered these 2nd Editions as the institutional repositories, all of which are available from the above-mentioned new (not old) URLs [138, 188-193]. In 2019, we published the 3rd Edition [197] of the Paper-6 (Comprehensive Review: Chapter-7 of the 2nd Edition of the Monograph) [138, 193], the 1st Edition of the Paper-7 (Memoirs: Chapter-8 of the 2nd Edition of the Monograph) [138, 198] and the 1st Edition of the Paper-8 (Japanese Edition of both the 3rd Edition of the Paper-6 with some fundamental concepts on life phenomena and Chapter-7 of the 2nd Edition of the Monograph), all of which are available from the corresponding new (not old) URLs [199]. In 2020, we published the 4th Edition [199] of the Paper-6 (Comprehensive Review: Chapter-7 of the 2nd Edition of the Monograph) [138, 197], the 2nd Edition of the Paper-7 (Memoirs: Chapter-8 of the 2nd Edition of the Monograph) [138, 198], and the 2nd Edition of the Paper-8 (Japanese Edition of both the 3rd Edition of the Paper-6 with some fundamental concepts on life phenomena and Chapter-7 of the 2nd Edition of the Monograph) [138, 199], all of which are available from the corresponding new (not old) URLs [197-199]. In 2021, we published the 3rd Edition of the Paper-5 on the 4C theory [221] (the 1st and 2nd Editions were published in 2017 and 2018 [192]), the 5th Edition of the Paper-6 on the Comprehensive Review [222] (the 1st, 2nd, 3rd and 4th Editions were published in 2017, 2018, 2019 and 2020 [193, 197, 199]), the 3rd Edition of the Paper-7 on the Memoirs [223] (the 1st and 2nd Editions were published in 2019 and 2020, with some fundamental concepts on life phenomena [198]), and the 3rd Edition of the Paper-8 on the Japanese

Review [224] (the 1st and 2nd Editions were published in 2019 and 2020 [199]). Of course, the 1st, 2nd and 3rd Editions of Paper-7 on the Memoirs were published only by myself. Moreover, in 2022, after published these 4 recent Papers-5, 6, 7 and 8 [221-224], we published the Comprehensive Review on the 4C theory in “Current Topics in Biochemical Research: Vol. 22, 2021” [225], which is the modified Edition of the 5th Edition (published in 2021) [222] of the Paper-6 on the Review Articles (the 1st, 2nd, 3rd and 4th Editions were published in 2017, 2018, 2019 and 2020) [193, 197, 199].

Finally, I should like to emphasize the following matters with some repetitions. The 4C theory [138, 188-199, 221-225] is obviously different in concepts and in molecular mechanisms from the two above-mentioned remarkable life phenomena, such as the experimental evolutions closely related to the alterations in genome information in *E. coli* and *D. melanogaster* [200-204] and the transgenerational epigenetic inheritances including the inter-tissue and soma-to-germ communications in *C. elegans* and *D. melanogaster* [205-220]. If we have been able to study the 4C theory in more favorable research environment, some of the above-mentioned questions concerning the exclusion of artificially accumulated IgM H- and L-chains in HDAC2(-/-) mutants during continuous cultivation must have been resolved at least. However, in these first and several revised Editions of the Monograph and the eight Papers (published only as institutional repositories), and in four Reviews in “Current Topics in Biochemical Research” [138, 188-199, 221-225], I could describe almost all the contents that I want to advocate based on only our insufficient data concerning the 4C theory, even though there must be a possibility of some over-descriptions. Naturally, as a concrete approach to generalize the 4C theory, for instance, elucidation of the influences of changes in temperature, atmosphere and/or nutrition on the ability to gain un-programmed and new cell function(s) in established cell lines through numerous generations and also in model animals (such as *C. elegans*, *D. melanogaster*, *Xenopus*, mice, rats and others) during development and differentiation is very helpful and significant. Because these influences under the varied conditions can be easily studied by various research groups, judging from the fact that previous researches in *D. melanogaster* were already conducted, although they were not related to molecular mechanisms [200, 203, 204, 209]. Hence, I earnestly crave that somebody has an interest in the universal concept on the 4C theory, takes over the theory, and elucidates some momentous questions on the theory including the above-mentioned ones in near future. When the 4C theory becomes clear, it can innovate the general notion on the nature of somatic cells and open the door for acquisition of adaptation ability of higher eukaryotes to the environment (intra- and/or extra-cellular) changes. Because the theory is one of the most fundamental and important phenomena for the life conservation and cell-type determination of higher eukaryotes through gaining un-programmed and new cell functions. In addition, I should like to just mention a following word. Future Biology that I mean must preferentially focus on the bio-systems (e. g., the 4C theory) for gaining un-programmed and new cell functions by means of

irreversible creation of chromatin structure plasticity with epigenetic modifications (which should inevitably occur) rather than alterations in genome information (which should incidentally and/or neutrally occur). Because the 4C theory is a new hypothetic concept on the bio-system to cope with, overcome and/or adapt to abnormal, unexpected, disadvantageous and/or even advantageous environment (intra- and/or extra-cellular) changes, when higher eukaryotes firstly encounter them.

In closing, first, I have an important matter, which must be described prior to acknowledgments. The biggest reason why I decided to upload this monograph to website is as follows, although I was just a mediocre life scientist. The life science is recently becoming a giant science more and more, and thereby big research groups with large grants are doing their research projects vigorously. Despite these circumstances, what I would like to tell young scientists in life science field is that even a small research group with a small grant can accomplish some small but creative concept, if the group has any of intellectual investigation spirit, intuition, insight plus enthusiasm, and quiet fighting spirit, diligent work plus outstanding techniques. Next, my special appreciation goes to Dr. Masami Nakayama, my wife, for her immeasurable contribution on the 4C theory with the enthusiasm, outstanding technique plus diligent work, and for her immense help throughout my life. In addition, I would like to express my grateful acknowledgments to Drs. T. Sukanuma and A. Sawaguchi for invaluable help in electron microscopy and immuno-electron microscopy studies. I am grateful to Drs. Y. Takami, H. Kikuchi and K. Toshimori for experimental support and to Dr. T. Hayashi for offer of office desk (regarding the 4C theory). I also thank all my laboratory members and colleagues and many (but relatively small) research grants for other research projects. And I must mention invaluable encouragement provided by my mentors, parents, brothers, sisters and children throughout my research life. All experiments on the 4C theory and others were done at Frontier Science Research Center and Faculty of Medicine, University of Miyazaki (Miyazaki Medical College).

Supplementary Remarks

All old URLs for the first and revised editions of Monograph, Papers and Reviews, all of which are related to the chromatin conformation change code (4C) theory, were changed to new URLs as follows. Therefore, all these first and several revised Editions of Monograph, Papers and Reviews, most of which were used as References in this article, are now available from new URLs but not from old URLs.

[Monograph]:

1st Edition (2015):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10123995>

New URL: <http://hdl.handle.net/10458/5293>

2nd Edition (2018):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10169365>

New URL: <http://hdl.handle.net/10458/6460>

[Papers]:

Paper-1 of 1st HDAC2(-/-):

1st Edition (2017):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10145170>

New URL: <http://hdl.handle.net/10458/5933>

2nd Edition (2018):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10170628>

New URL: <http://hdl.handle.net/10458/6500>

Paper-2 of Pax5(-):

1st Edition (2017):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10145176>

New URL: <http://hdl.handle.net/10458/5934>

2nd Edition (2018):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10170947>

New URL: <http://hdl.handle.net/10458/6503>

Paper-3 of 2nd HDAC2(-/-):

1st Edition (2017):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10145178>

New URL: <http://hdl.handle.net/10458/5935>

2nd Edition (2018):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10170948>

New URL: <http://hdl.handle.net/10458/6504>

Paper-4 of NotchIP:

1st Edition (2017):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10145181>

New URL: <http://hdl.handle.net/10458/5936>

2nd Edition (2018):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10170993>

New URL: <http://hdl.handle.net/10458/6505>

Paper-5 of the 4C Theory:

1st Edition (2017):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10145263>

New URL: <http://hdl.handle.net/10458/5937>

2nd Edition (2018):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10171004>

New URL: <http://hdl.handle.net/10458/6506>

3rd Edition (2021):

New URL: <http://hdl.handle.net/10458/6506>

Paper-6 of Comprehensive (All-inclusive) Review:

1st Edition (2017):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10146006>

New URL: <http://hdl.handle.net/10458/5941>

2nd Edition (2018):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10171009>

New URL: <http://hdl.handle.net/10458/6507>

3rd Edition (2019):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10174272>

New URL: <http://hdl.handle.net/10458/6613>

4th Edition (2020):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10182073>

New URL: <http://hdl.handle.net/10458/6902>

5th Edition (2021):

New URL: <http://hdl.handle.net/10458/6902>

Paper-7 of Memoirs (Insights):

1st Edition (2019):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10174284>

New URL: <http://hdl.handle.net/10458/6614>

2nd Edition (2020):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10185502>

New URL: <http://hdl.handle.net/10458/6949>

3rd Edition (2021):

New URL: <http://hdl.handle.net/10458/6949>

Paper-8 of Japanese Review:

1st Edition (2019):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10174562>

New URL: <http://hdl.handle.net/10458/6615>

2nd Edition (2020):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10185510>

New URL: <http://hdl.handle.net/10458/6950>

3rd Edition (2021):

New URL: <http://hdl.handle.net/10458/6950>

[Review articles in Current Topics in Biochemical Research]:

Review-1 (Current Topics in Biochemical Research, Vol.18, pp.11-25, 2017):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10168788>

New URL: <http://hdl.handle.net/10458/6375>

Review-2 (Current Topics in Biochemical Research, Vol.18, pp.33-56, 2017):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10168794>

New URL: <http://hdl.handle.net/10458/6376>

Review-3 (Current Topics in Biochemical Research, Vol.18, pp.65-86, 2017):

Old URL: <http://opac2.lib.miyazaki-u.ac.jp/webopac/TC10168800>

New URL: <http://hdl.handle.net/10458/6377>

Current Topics in Biochemical Research, Vol.18, pp.11-25, pp.33-56, pp.65-86, 2017:

Website: http://www.researchtrends.net/tia/title_issue.asp?id=40&in=0&vn=18&type=3

Review-4 (Current Topics in Biochemical Research, Vol. 22, pp.43-90, 2021):

URL: <http://hdl.handle.net/10458/00010392>

Current Topics in Biochemical Research, Vol.22, pp.43-90, 2021:

Website: http://www.researchtrends.net/tia/title_issue.asp?id=40&in=0&vn=22&type=3

Numbers of bibliographic and text references of old URLs of the Monograph, Papers and Reviews by August/2020 were as follows: Probably, text references mainly mean Download of Text and/or Figures.

Numbers of bibliographic and text references of old URL of the Monograph (1st Edition published in 2015) were 495 and 423, respectively. Numbers of bibliographic and text references of old URL of the Monograph (2nd Edition published in 2018) were 67 and 292, respectively.

Numbers of bibliographic and text references of old URL of the Paper-1 of 1st HDAC2(-/-) (1st Edition published in 2017) were 138 and 69, respectively. Numbers of bibliographic and text references of old URL of the Paper-1 of 1st HDAC2(-/-) (2nd Edition published in 2018) were 62 and 41, respectively.

Numbers of bibliographic and text references of old URL of the Paper-2 of Pax5(-) (1st Edition published in 2017) were 120 and 62, respectively. Numbers of bibliographic and text references of old URL of the Paper-2 of Pax5(-) (2nd Edition published in 2018) were 60 and 39, respectively.

Numbers of bibliographic and text references of old URL of the Paper-3 of 2nd HDAC2(-/-) (1st Edition published in 2017) were 112 and 58, respectively. Numbers of bibliographic and text references of old URL of the Paper-3 of 2nd HDAC2(-/-) (2nd Edition published in 2018) were 34 and 35, respectively.

Numbers of bibliographic and text references of old URL of the Paper-4 of NotchIP (1st Edition published in 2017) were 114 and 121, respectively. Numbers of bibliographic and text references of old URL of the Paper-4 of NotchIP (2nd Edition published in 2018) were 31 and 69, respectively.

Numbers of bibliographic and text references of old URL of the Paper-5 of the 4C Theory (1st Edition published in 2017) were 208 and 132, respectively. Numbers of bibliographic and text references of old URL of the Paper-5 of the 4C Theory (2nd Edition published in 2018) were 62 and 72, respectively.

Numbers of bibliographic and text references of old URL of the Paper-6 of Comprehensive (All-inclusive) Review (1st Edition published in 2017) were 131 and 152, respectively. Numbers of bibliographic and text references of old URL of the Paper-6 of Comprehensive (All-inclusive) Review (2nd Edition published in 2018) were 43 and 70, respectively. Numbers of bibliographic and text references of old URL of the Paper-6 of Comprehensive (All-inclusive) Review (3rd Edition published in 2019) were 32 and 59, respectively. Numbers of bibliographic and text references of old URL of the Paper-6 of Comprehensive (All-inclusive) Review (4th Edition published in 2020) were overlooked, respectively.

Numbers of bibliographic and text references of old URL of the Paper-7 of Memoirs (Insights) (1st Edition published in 2019) were 28 and 59, respectively. Numbers of bibliographic and text references of old URL of the Paper-7 of Memoirs (Insights) (2nd Edition published in 2020) were overlooked, respectively.

Numbers of bibliographic and text references of old URL of the Paper-8 of Japanese Review (1st Edition published in 2019) were 54 and 99, respectively. Numbers of bibliographic and text references of old URL of the Paper-8 of Japanese Review (2nd Edition published in 2020) were overlooked, respectively.

Numbers of bibliographic and text references of old URL of the Review-1 of Current Topics in Biochemical Research (Vol.18, pp.11-25, 2017) were 53 and 11, respectively.

Numbers of bibliographic and text references of old URL of the Review-2 of Current Topics in Biochemical Research (Vol.18, pp.-56, 2017) were 46 and 6, respectively.

Numbers of bibliographic and text references of old URL of the Review-3 of Current Topics in Biochemical Research (Vol.18, pp.65-86, 2017) were 66 and 11, respectively.

References

- [1] Allfrey, V., Faulker, R. M. and Mirsky, A. E.: Acetylation and methylation of histones and their possible role in the regulation of RNA synthesis. Proc. Natl. Acad. Sci. USA, 51, 786-794, 1964.
- [2] Kornberg, R. D.: Structure of chromatin. Annu. Rev. Biochem. 46, 931-954, 1977.
- [3] Felsenfeld, G.: Chromatin. Nature 271, 115-121, 1978.
- [4] van Holde, K. E.: Chromatin. New York: Springer-Verlag, 1989.

- [5] Brownell, J. E., Zhou, J., Rannali, T., Kobayashi, R., Edmondson, D. G., Roth, S. Y. and Allis, C. D.: Tetrahymena histone acetyltransferase A: a homolog to yeast Gcn5 linking histone acetylation to gene activation. *Cell* 84, 843-851, 1996.
- [6] Ogryzko, V. V., Schiltz, R. L., Russanova, V., Howard, B. H. and Nakatani, Y.: The transcriptional coactivators p300 and CPB are histone acetyltransferases. *Cell* 87, 953-959, 1996.
- [7] Taunton, J., Hassig, C. A. and Schreiber, S. L.: A mammalian histone deacetylase related to the yeast transcriptional regulator Rpd3p. *Science* 272, 408-411, 1996.
- [8] Wolffe, A. P.: Histone deacetylase: a regulator of transcription. *Science* 272, 371-372, 1996.
- [9] Hassig, C. A. and Schreiber, S. L.: Nuclear histone acetylases and deacetylases and transcriptional regulation: HATs off to HDACs. *Curr. Opin. Chem. Biol.* 1, 300-308, 1997.
- [10] Varga-Weisz, P. D. and Becker, P. B.: Chromatin-remodeling factors: machines that regulate? *Curr. Opin. Cell Biol.* 10, 346-353, 1998.
- [11] Workman, J. L. and Kingston, R. E.: Alteration of nucleosome structure as a mechanism of transcriptional regulation. *Annu. Rev. Biochem.* 67, 545-579, 1998.
- [12] Davie, J. R.: Covalent modifications of histones: expression from chromatin templates. *Curr. Opin. Genet. Dev.* 8, 173-178, 1998.
- [13] Luger, K. and Richmond, T. J.: The histone tails of the nucleosome. *Curr. Opin. Genet. Dev.* 8, 140-146, 1998.
- [14] Johnson, C. A. and Turner, B. M.: Histone deacetylases: complex transducers of nuclear signals. *Semin. Cell Dev. Biol.* 10, 179-188, 1999.
- [15] Brown, C. E., Lechner, T., Howe, L. and Workman, J. L.: The many HATs of transcription coactivators. *Trends Biochem. Sci.* 25, 15-19, 2000.
- [16] Cheung, W. L., Briggs, S. D. and Allis, C. D.: Acetylation and chromosomal functions. *Curr. Opin. Cell Biol.* 12, 326-333, 2000.
- [17] Strahl, B. D. and Allis, C. D.: The language of covalent histone modifications. *Nature* 403, 41-45, 2000.
- [18] Turner, B. M.: Histone acetylation and an epigenetic code. *Bioessays* 22, 836-845, 2000.
- [19] Jenuwein, T. and Allis, C. D.: Translating the histone code. *Science* 293, 1074-1080, 2001.
- [20] Khochbin, S., Verdel, A., Lemercier, C. and Seigneurin-Berny, D.: Functional significance of histone deacetylase diversity. *Curr. Opin. Genet. Dev.* 11, 162-166, 2001.
- [21] Roth, S. Y., Denu, J. M. and Allis, C. D.: Histone acetyltransferases. *Annu. Rev. Biochem.* 70, 81-120, 2001.
- [22] Carrozza, M. J., Utley, R. T., Workman, J. L. and Cote, J.: The diverse functions of histone acetyltransferase complexes. *Trend. Genet.* 19, 321-329, 2003.
- [23] Yang, X. J. and Seto, E.: Collaborative spirit of histone deacetylases in regulating chromatin

structure and gene expression. *Curr. Opin. Genet. Dev.* 13, 143-153, 2003.

[24] Wang, S., Yan-Neale, Y., Zeremski, M. and Cohen, D.: Transcription regulation by histone deacetylases. *Novartis Found Symp.* 259, 238-245, 2004.

[25] Biel, M., Wascholowski, V. and Giannis, A.: Epigenetics - an epicenter of gene regulation: histones and histone-modifying enzymes. *Angew. Chem. Int. Ed. Engl.* 44, 3186-3216, 2005.

[26] Margueron, R., Trojer, P. and Reinberg, D.: The key to development: interpreting the histone code? *Curr. Opin. Genet. Dev.* 15, 163-176, 2005.

[27] Kouzarides, T.: Chromatin modifications and their function. *Cell* 128, 693-705, 2007.

[28] Allis, C. D., Berger, S. L., Cote, J., Dent, S., Jenuwien, T., Kouzarides, T., Pillus, L., Reinberg, D., Shi, Y., Shiekhatar, R., Shilatifard, A., Workman, J. and Zhang, Y.: New nomenclature for chromatin-modifying enzymes. *Cell* 131, 633-636, 2007.

[29] Lee, K. K. and Workman, J. L.: Histone acetyltransferase complexes: one size doesn't fit all. *Nat. Rev. Mol. Cell Biol.* 8, 284-295, 2007.

[30] Goldberg, A. D., Allis, C. D. and Bernstein, B. E.: Epigenetics: a landscape takes shape. *Cell* 128, 635-638, 2007.

[31] Shahbazian, M. D. and Grunstein, M.: Functions of site-specific histone acetylation and deacetylation. *Annu. Rev. Biochem.* 76, 75-100, 2007.

[32] Berger, S. L.: The complex language of chromatin regulation during transcription. *Nature* 447, 407-412, 2007.

[33] Suganuma, T. and Workman, J. L.: Crosstalk among histone modifications. *Cell* 135, 604-607, 2008.

[34] Kohn, K. W., Aladjem, M. I., Weinstein, J. N. and Pommier, Y.: Chromatin challenges during DNA replication: A systems representation. *Mol. Biol. Cell* 19, 1-7, 2008.

[35] Selvi, R. B. and Kundu, T. K.: Reversible acetylation of chromatin: implication in regulation of gene expression, disease and therapeutics. *Biotech. J.* 4, 375-390, 2009.

[36] Javierre, B. M., Hemando, H. and Ballestar, E.: Environmental triggers and epigenetic deregulation in autoimmune disease. *Discov. Med.* 12, 535-545, 2011.

[37] Bannister, A. J. and Kouzarides, T.: Regulation of chromatin by histone modifications. *Cell Res.* 21, 381-395, 2011.

[38] Verrier, L., Vandromme, M. and Trouche, D.: Histone demethylases in chromatin cross-talks. *Biol. Cell* 103, 381-401, 2011.

[39] Butler, J. S., Koutelou, E., Schibler, A. C. and Dent, S. Y.: Histone-modifying enzymes: regulators of developmental decisions and drivers of human disease. *Epigenomics* 4, 163-177, 2012.

[40] Kooistra, S. M. and Helin, K.: Molecular mechanisms and potential functions of histone deacetylases. *Nat. Rev. Mol. Cell Biol.* 13, 297-311, 2012.

[41] Graff, J. and Tsai, L.-H.: Histone acetylation: molecular mnemonics on the chromatin. *Nat. Rev.*

Neurosci. 14, 97-111, 2013.

[42] Chen, T. and Dent, S. Y. R.: Chromatin modifiers and remodellers: regulators of cellular differentiation. *Nat. Rev. Genet.* 15, 93-106, 2014.

[43] T. W.-W. and Reinberg, D.: Chromatin futures and the epigenetic regulation of pluripotency states in ESCs. *Development* 141, 2376-2390, 2014.

[44] Morgan, M. A. and Shilatifard, A.: Chromatin signatures of cancer. *Genes and Dev.* 29, 238-249, 2015.

[45] Staudt, L. M. and Lenardo, M. J.: Immunoglobulin gene transcription. *Annu. Rev. Immunol.* 9, 373-398, 1991.

[46] Peterson, M. L.: RNA processing and expression of immunoglobulin genes. In *Handbook of B and T Lymphocytes* (Snow, E. C. ed.) pp321-342, 1994, Academic press, San Diego.

[47] Calame, K. L., Lin, K. I. and Tunyaplin, C.: Regulatory mechanisms that determine the development and function of plasma cells. *Annu. Rev. Immunol.* 21, 205-230, 2003.

[48] Busslinger, M.: Transcriptional control of early B cell development. *Annu. Rev. Immunol.* 22, 55-79, 2004.

[49] Su, I. H. and Tarakhovsky, A.: Epigenetic control of B cell differentiation. *Semin. Immunol.* 17, 167-172, 2005.

[50] Nutt, S. L. and Kee, B. L.: The transcriptional regulation of B cell lineage commitment. *Immunity* 26, 715-725, 2007.

[51] Ramirez, J., Lukin, K. and Hagman, J.: From hematopoietic progenitors to B cells: mechanisms of lineage restriction and commitment. *Curr. Opin. Immunol.* 22, 177-184, 2010.

[52] Funahashi, J., Okufuji, T., Ohuchi, H., Noji, S., Tanaka, H. and Nakamura, H.: Role of Pax-5 in the regulation of a mid-hindbrain organizer's activity. *Dev. Growth Differ.* 41, 59-72, 1999.

[53] Stevens, S., Ong, J., Kim, U., Eckhardt, L. A. and Roeder, R. G.: Role of OCA-B in 3'-IgH enhancer function. *J. Immunol.* 164, 5306-5312, 2000.

[54] Hannon, G. J.: RNA interference. *Nature* 418, 244-251, 2002.

[55] Shapiro-Shelef, M., Lin, K. I., McHeyzer-Williams, L. J., Liao, J., McHeyzer-Williams, M. G. and Calame, K.: Blimp-1 is required for the formation of immunoglobulin secreting plasma cells and pre-plasma memory B cells. *Immunity* 19, 607-620, 2003.

[56] Savitsky, D. and Calame, K.: B-1 B lymphocytes require Blimp-1 for immunoglobulin secretion. *J. Exp. Med.* 203, 2305-2314, 2006.

[57] Nera, K.-P., Kohonen, P., Narvi, E., Peippo, A., Mustonen, L., Terho, P., Koskele, K., Buerstedde, J.-M. and Lassila, O.: Loss of Pax5 promotes plasma cell differentiation. *Immunity* 24, 283-293, 2006.

[58] Magari, M., Aya, T., Ikeda, M., Todo, K., Kanayama, N. and Ohmori, H.: Enhancement of antibody production from a chicken B cell line DT40 by reducing Pax5 expression. *J. Biosci. Bioeng.* 107, 206-209,

2009.

- [59] Toman, I., Loree, J., Klimowicz, A. C., Bahlis, N., Lai, R., Belch, A., Pilarski, L. and Reiman, T.: Expression and prognostic significance of Oct2 and Bob1 in multiple myeloma: implications for targeted therapeutics. *Leuk. Lymphoma*. 52, 659-667, 2011.
- [60] Fujita, T. and Fujii, H.: Species-specific 5'-genomic structure and multiple transcription start sites in the chicken Pax5 gene. *Gene* 477, 24-31, 2011.
- [61] John, L. B. and Ward, A. C.: The Ikaros gene family: transcriptional regulators of hematopoiesis and immunity. *Mol. Immunol.* 48, 1272-1278, 2011.
- [62] Nakayama, T. and Setoguchi, Y.: Nucleotide sequence of a member of the chicken H2B histone-encoding gene family. *Gene* 98, 299-300, 1991.
- [63] Nakayama, T.: Nucleotide sequences of two members of the chicken H3 histone-encoding gene family. *Gene* 102, 289-290, 1991.
- [64] Setoguchi, Y. and Nakayama, T.: Nucleotide sequences of new members (H3-IV and H3-V) of the chicken H3 histone-encoding gene family. *Nucleic Acids Res.* 19, 6327, 1991.
- [65] Nakayama, T., Takechi, S., Ohshige, T., Kondo, K. and Yamamoto, K.: Nucleotide sequences of two members of the chicken H4 histone-encoding gene family. *Gene* 108, 311-312, 1991.
- [66] Nakayama, T. and Setoguchi, Y.: Structural change of a particular H2B histone gene possibly results in differences in its transcriptional regulation in different tissues. *Nucleic Acids Res.* 20, 1309-1314, 1992.
- [67] Nakayama, T. and Setoguchi, Y.: Evidence for possible structural changes of particular H3 and H2B histone genes in different chicken tissues (cells). *Biochem. Biophys. Res. Commun.* 184, 317-322, 1992.
- [68] Takami, Y. and Nakayama, T.: Presence of distinct transcriptional regulatory elements in the 5'-flanking region shared by the chicken H3 histone gene homopair. *Nucleic Acids Res.* 20, 3037-3041, 1992.
- [69] Ohshige, T., Takechi, S. and Nakayama, T.: Presence of particular transcription regulatory elements in the 5'-intergenic region shared by the chicken H2A-III and H2B-V pair. *Gene* 131, 193-199, 1993.
- [70] Nakayama, T., Takechi, S. and Takami, Y.: The chicken histone gene family. *Comp. Biochem. Physiol.* 104B, 635-639, 1993.
- [71] Takami, Y. and Nakayama, T.: Possible involvement of a ubiquitous and several distinct elements in the transcription regulation of the chicken H3 histone gene family. *Biochim. Biophys. Acta* 1264, 29-34, 1995.
- [72] Takami, Y., Higashio, M., Fukuoka, T., Takechi, S. and Nakayama, T.: Organization of the chicken histone genes in a major gene cluster and generation of an almost complete set of the core histone protein sequences. *DNA Res.* 3, 95-99, 1996.

- [73] Nakayama, T., Titani, K. and Narita, K.: The amino acid sequence of cytochrome c from bonito (*Katsuwonus pelamis*, Linnaeus). *J. Biochem.* 70, 311-326, 1971.
- [74] Nakayama, T., Ogata, K. and Narita, K.: Ovalbumin synthesis in a homologous cell-free system prepared from hen's oviduct. *J. Biochem.* 79, 853-869, 1976.
- [75] Nakayama, T., Narita, K. and Ogata, K.: Biological process of carbohydrate attachment to ovalbumin. *J. Biochem.* 79, 871-881, 1976.
- [76] Nakayama, T. and Narita, K.: Regulation of protein synthesis in hen's oviducts. I. Removal of extracellular and intracellular proteins and their effects on protein synthesis. *J. Biochem.* 81, 153-162, 1977.
- [77] Nakayama, T., Munoz, L. and Doi, R. H.: A procedure to remove protease activities from *Bacillus subtilis* sporulating cells and their crude extracts. *Anal. Biochem.* 78, 165-170, 1977.
- [78] Nakayama, T., Sugae, K. and Narita, K.: Regulation of protein synthesis in hen's oviducts. II. Extracellular ovalbumin as an inhibitor of ovalbumin synthesis in the oviducts. *J. Biochem.* 81, 1803-1812, 1977.
- [79] Munoz, L. E., Nakayama, T. and Doi, R. H.: Expression of spore coat protein gene, an "early sporulation gene", and its relationship to RNA polymerase modification. *Spores* 7, 213-219, 1978.
- [80] Nakayama, T., Williamson, V., Burtis, K. and Doi, R. H.: Purification and properties of two RNA polymerases from sporulating cells of *Bacillus subtilis*. *Eur. J. Biochem.* 88, 155-164, 1978.
- [81] Nakayama, T., Munoz, L., Sadaie, Y. and Doi, R. H.: Spore coat protein synthesis in cell-free systems from sporulating cells of *Bacillus subtilis*. *J. Bacteriol.* 135, 952-960, 1978.
- [82] Nakayama, T., Kurogi, Y., Irikura, M. and Matsuo, H.: Cooperative effects of Daunorubicin and Actinomycin D on the sporulation of *Bacillus subtilis*. *J. Biochem.* 87, 1619-1624, 1980.
- [83] Nakayama, T., Kurogi, Y., Irikura, M., Matsuo, H. and Sakai-Wada, A.: Fractionation and biochemical properties of the mother cell and forespore fractions from sporulating cells of *Bacillus subtilis*. *J. Biochem.* 88, 317-326, 1980.
- [84] Nakayama, T.: Purification of *B. subtilis* DNA-binding protein related to *E. coli* HU. *Biochem. Biophys. Res. Commun.* 97, 318-324, 1980.
- [85] Nakayama, T., Irikura, M., Kurogi, Y. and Matsuo, H.: Purification and properties of RNA polymerases from mother cells and forespores of sporulating cells of *Bacillus subtilis*. *J. Biochem.* 89, 681-691, 1981.
- [86] Nakayama, T., Kurogi, Y. and Irikura, M.: Purification of polypeptide P6 with a molecular weight of 6,000 from the vegetative chromosomes of *Bacillus subtilis*. *J. Bacteriol.* 145, 953-957, 1981.
- [87] Nakayama, T.: Identification of vitellogenin selector which binds tightly to vitellogenin gene-carrying DNA fragments. *Proc. Japan Acad.* 61, 52-55, 1985.

- [88] Nakayama, T., Irikura, M., Setoguchi, Y., Nakayama, M., Mochizuki, M. and Ogata, K.: Estrogen-mediated induction of a vitellogenin-specific non-histone chromatin protein in the male chicken liver. *Mol. Gen. Genet.* 201, 252-257, 1985.
- [89] Nakayama, T., Akizuki, N., Irikura, M., Setoguchi, Y. and Ogata, K.: Vitellogenin-specific non-histone chromatin protein in the liver of egg-laying hens. *Gunma Symposia on Endocrinology* 24, 105-118, 1987.
- [90] Nakayama, T., Irikura, M., Setoguchi, Y., Matsuo, H., Inoue, H., Yutsudo, M. and Hakura, A.: Purification and properties of a non-histone chromatin protein with a molecular weight of 38,000 daltons specific for transformed rat cells. *Biochem. Biophys. Res. Commun.* 149, 1156-1164, 1987.
- [91] Baba, T. W., Giroir, B. P. and Humphries, E. H.: Cell lines derived from avian lymphomas exhibit two distinct phenotypes. *Virology* 144, 139-151, 1985.
- [92] Buerstedde, J.-M. and Takeda, S.: Increased ratio of targeted to random integration after transfection of chicken B cell lines. *Cell* 67, 179-188, 1991.
- [93] Takeda, S., Masteller, E. L., Thompson, C. B. and Buerstedde, J.-M.: RAG-2 expression is not essential for chicken immunoglobulin gene conversion. *Proc. Natl. Acad. Sci. USA* 89, 4023-4027, 1992.
- [94] Ahmad, A., Takami, Y. and Nakayama, T.: WD repeats of the p48 subunit of chicken chromatin assembly factor-1 required for in vitro interaction with chicken histone deacetylase-2. *J. Biol. Chem.* 274, 16646-16653, 1999.
- [95] Nakayama, T., Takami, Y. and Ahmad, A.: Interaction of the p48 subunit of chicken chromatin assembly factor-1 with histone deacetylases. *Current Topics in Biochemical Research.* 1, 173-178, 1999.
- [96] Ahmad, A., Takami, Y. and Nakayama, T.: Distinct regions of the chicken p46 polypeptide are required for its in vitro interaction with histones H2B and H4 and histone acetyltransferase-1. *Biochem. Biophys. Res. Commun.* 279, 95-102, 2000.
- [97] Nakayama, T. and Takami, Y.: Histone deacetylase-2 controls the amount of the IgM H-chain at the steps of both transcription and alternative pre-mRNA processing in the DT40 cell line. *Trends in Comparative Biochem. & Physiol.* 7, 93-100, 2000.
- [98] Ahmad, A., Nagamatsu, N., Kouriki, H., Takami, Y. and Nakayama, T.: Leucine zipper motif of chicken histone acetyltransferase-1 is essential for in vivo and in vitro interactions with the p48 subunit of chicken chromatin assembly factor-1. *Nucleic Acids Res.* 29, 629-637, 2001.
- [99] Nakayama, T., Takami, Y. and Ahmad, A.: Interaction of the p48 subunit of chicken chromatin assembly factor-1 and the p46 polypeptide with histones and histone-modifying enzymes. *Res. Adv. in Biological Chem.* 1, 21-24, 2001.
- [100] Ahmad, A., Takami, Y. and Nakayama, T.: WD dipeptide motifs and LXXLL motif of chicken HIRA are necessary for transcription repression and the latter motif is essential for interaction with histone deacetylase-2 in vivo. *Biochem. Biophys. Res. Commun.* 312, 1266-1272, 2003.

- [101] Ahmad, A., Takami, Y. and Nakayama, T.: WD dipeptide motifs and LXXLL motif of chicken HIRA are essential for interactions with the p48 subunit of chromatin assembly factor-1 and histone deacetylase-2 in vitro and in vivo. *Gene* 342, 125-136, 2004.
- [102] Ahmad, A., Kikuchi, H., Takami, Y. and Nakayama, T.: Different roles of N-terminal and C-terminal halves of HIRA in transcription regulation of cell cycle-related genes that contribute to control of vertebrate cell growth. *J. Biol. Chem.* 280, 32090-32100, 2005.
- [103] Takami, Y., Takeda, S. and Nakayama, T.: Targeted disruption of an H3-IV/H3-V gene pair causes increased expression of the remaining H3 genes in the chicken DT40 cell line. *J. Mol. Biol.* 250, 420-433, 1995.
- [104] Seguchi, K., Takami, Y. and Nakayama, T.: Targeted disruption of 01H1 encoding a particular H1 histone variant causes changes in protein patterns in the DT40 chicken B cell line. *J. Mol. Biol.* 254, 869-880, 1995.
- [105] Takami, Y., Takeda, S. and Nakayama, T.: Targeted disruption of H2B-V encoding a particular H2B histone variant causes changes in protein patterns on two-dimensional polyacrylamide gel electrophoresis in the DT40 chicken B cell line. *J. Biol. Chem.* 270, 30664-30670, 1995.
- [106] Takami, Y., Takeda, S. and Nakayama, T.: An approximately half set of histone genes is enough for cell proliferation and a lack of several histone variants causes protein pattern changes in the DT40 chicken B cell line. *J. Mol. Biol.* 265, 394-408, 1997.
- [107] Takami, Y. and Nakayama, T.: One allele of the major histone gene cluster is enough for cell proliferation of the DT40 chicken B cell line. *Biochim. Biophys. Acta* 1354, 105-115, 1997.
- [108] Takami, Y. and Nakayama, T.: A single copy of linker H1 genes is enough for proliferation of the DT40 chicken B cell line, and linker H1 variants participate in regulation of gene expression. *Genes Cells* 2, 711-723, 1997.
- [109] Takami, Y., Nishi, R. and Nakayama, T.: Histone H1 variants play individual roles in transcription regulation in the DT40 chicken B cell line. *Biochem. Biophys. Res. Commun.* 268, 501-508, 2000.
- [110] Nakayama, T. and Takami, Y.: Participation of histones and histone-modifying enzymes in cell functions through alterations in chromatin structure. *J. Biochem.* 129, 491-499, 2001.
- [111] Sanematsu, F., Takami, Y., Barman, H. K., Fukagawa, T., Ono, T., Shibahara, K. and Nakayama, T.: Asf1 is required for viability and chromatin assembly during DNA replication in vertebrate cells. *J. Biol. Chem.* 281, 13817-13827, 2006.
- [112] Takami, Y., Ono, T., Fukagawa, T., Shibahara, K. and Nakayama, T.: Essential role of CAF-1-mediated rapid nucleosome assembly for DNA replication and cell division in vertebrate cells. *Mol. Biol. Cell* 18, 129-141, 2007.

- [113] Sonoda, E., Zhao, G.-Y., Kohzaki, M., Dhar, P. K., Kikuchi, K., Redon, C., Pilch, D. R., Bonner, W. W., Nakano, A., Watanabe, M., Nakayama, T., Takeda, S. and Takami, Y.: Collaborative roles of rH2AX and the Rad51 paralog Xrcc3 in homologous recombinational repair. *DNA Repair* 6, 280-292, 2007.
- [114] Takami, Y., Kikuchi, H. and Nakayama, T.: Chicken histone deacetylase-2 controls the amount of the IgM H-chain at the steps of both transcription of its gene and alternative processing of its pre-mRNA in the DT40 cell line. *J. Biol. Chem.* 274, 23977-23990, 1999.
- [115] Nakayama, T. and Takami, Y.: Systematic analysis of the nature of histone genes in the chicken DT40 B cell line using gene targeting techniques. *Trends in Comparative Biochem. & Physiol.* 4, 211-217, 1998.
- [116] Takami, Y. and Nakayama, T.: N-terminal region, C-terminal region, nuclear export signal and deacetylation activity of histone deacetylase-3 are essential for the viability of the DT40 chicken B cell line. *J. Biol. Chem.* 275, 16191-16201, 2000.
- [117] Takechi, S., Adachi, M. and Nakayama, T.: Chicken HDAC2 down-regulates IgM light chain gene promoter activity. *Biochem. Biophys. Res. Commun.* 299, 263-267, 2002.
- [118] Kikuchi, H., Takami, Y. and Nakayama, T.: GCN5: a supervisor in all-inclusive control of vertebrate cell cycle progression through transcription regulation of various cell cycle-related genes. *Gene* 347, 83-97, 2005.
- [119] Kikuchi, H., Barman, H. K., Nakayama, M., Takami, Y. and Nakayama, T.: Participation of histones, histone modifying enzymes and histone chaperones in vertebrate cell functions. *Reviews and Protocols in DT40 Research*, Springer-Verlag, Berlin, pp225-243, 2006.
- [120] Barman, H. K., Takami, Y., Ono, T., Nishijima, H., Sanematsu, F., Shibahara, K. and Nakayama, T.: Histone acetyltransferase 1 is dispensable for replication-coupled chromatin assembly but contributes to recover DNA damages created following replication blockage in vertebrate cells. *Biochem. Biophys. Res. Commun.* 345, 1547-1557, 2006.
- [121] Nakayama, M., Suzuki, H., Yamamoto-Nagamatsu, N., Barman, H. K., Kikuchi, H., Takami, Y., Toyonaga, K., Yamashita, K. and Nakayama, T.: HDAC2 controls IgM H and L-chain gene expressions via EBF1, Pax5, Ikaros, Aiolos and E2A gene expressions. *Genes Cells* 12, 359-373, 2007.
- [122] Kikuchi, H. and Nakayama, T.: GCN5 and BCR signaling collaborate to induce pre-mature B cell apoptosis through depletion of ICAD and IAP2 and activation of caspase activities. *Gene* 419, 48-55, 2008.
- [123] Barman, H. K., Takami, Y., Nishijima, H., Shibahara, K., Sanematsu, F. and Nakayama, T.: Histone acetyltransferase-1 regulates integrity of cytosolic histone H3-H4 containing complex. *Biochem. Biophys. Res. Commun.* 373, 624-630, 2008.

- [124] Kikuchi, H., Barman, H. K., Nakayama, M., Takami, Y. and Nakayama, T.: Studies on epigenetic control of B cell functions using the DT40 cell line. *Advances in Genetics Research* 2, Urbano K. V. (Ed.), Nova Science Publishers, Inc. NY, pp153-166, 2010.
- [125] Kikuchi, H., Kuribayashi, F., Takami, Y., Imajoh-Ohmi, S. and Nakayama, T.: GCN5 regulates the activation of PI3K/Akt survival pathway in B cells exposed to oxidative stress via controlling gene expressions of Syk and Btk. *Biochem. Biophys. Res. Commun.* 405, 657-661, 2011.
- [126] Kikuchi, H., Kuribayashi, F., Kiwaki, N., Takami, Y. and Nakayama, T.: GCN5 regulates the superoxide-generating system in leukocytes via controlling gp91-phox gene expression. *J. Immunol.* 186, 3015-3022, 2011.
- [127] Kikuchi, H., Kuribayashi, F., Imajoh-Ohmi, S., Nishitoh, Y., Takami, Y. and Nakayama, T.: GCN5 protects vertebrate cells against UV-irradiation via controlling gene expression of DNA polymerase η . *J. Biol. Chem.* 287, 39842-39849, 2012.
- [128] Kikuchi, H., Nakayama, M., Kuribayashi, F., Imajoh-Ohmi, S., Nishitoh, H., Takami, Y. and Nakayama T.: GCN5 is essential for IRF-4 gene expression followed by transcriptional activation of Blimp-1 in immature B cells. *J. Leukoc. Biol.* 95, 399-404, 2014.
- [129] Kikuchi, H., Nakayama, M., Kuribayashi, F., Imajoh-Ohmi, S., Nishitoh, H., Takami, Y. and Nakayama, T.: Protein kinase C θ gene expression is oppositely regulated by EBF1 and GCN5 in immature B cells. *FEBS Lett.* 588, 1739-1742, 2014.
- [130] Kikuchi, H., Nakayama, M., Kuribayashi, F., Imajoh-Ohmi, S., Nishitoh, H., Takami, Y. and Nakayama T.: GCN5 is involved in regulation of immunoglobulin heavy chain gene expression in immature B cells. *Gene* 544, 19-24, 2014.
- [131] Escaffit, F., Vaute, O., Chevillard-Briet, M., Segui, B., Takami, Y., Nakayama, T. and Trouche, D.: Cleavage and cytoplasmic relocalization of histone deacetylase 3 are important for apoptosis progression. *Mol. Cell. Biol.* 27, 554-567, 2007.
- [132] Kikuchi, H., Kuribayashi, F., Kiwaki, N. and Nakayama, T.: Curcumin dramatically enhances retinoic acid-induced superoxide generating activity via accumulation of p47-phox and p67-phox proteins in U937 cells. *Biochem. Biophys. Res. Commun.* 395, 61-65, 2010.
- [133] Kikuchi, H., Kuribayashi, F., Mimuro, H., Imajoh-Ohmi, S., Nakayama, M., Takami, Y., Nishitoh, H. and Nakayama, T.: Lack of GCN5 remarkably enhances the resistance against prolonged endoplasmic reticulum stress-induced apoptosis through up-regulation of Bcl-2 gene expression. *Biochem. Biophys. Res. Commun.* 463, 870-875, 2015.
- [134] Kikuchi, H., Nakayama, M., Kuribayashi, F., Mimuro, H., Imajoh-Ohmi, S., Nishitoh, H., Takami, Y. and Nakayama, T.: Paired box gene 5 isoforms A and B have different functions in transcriptional regulation of B cell development-related genes in immature B cells. *Microbiol. Immunol.* 59, 426-431, 2015.

- [135] Kikuchi, H., Nakayama, M., Kuribayashi, F., Mimuro, H., Imajoh-Ohmi, S., Nishitoh, H. and Nakayama, T.: Histone acetyltransferase PCAF is involved in transactivation of Bcl-6 and Pax5 genes in immature B cells. *Biochem. Biophys. Res. Commun.* 467, 509-513, 2015.
- [136] Kikuchi, H., Nakayama, M., Kawai, C., Kuribayashi, F., Mimuro, H., Imajoh-Ohmi, S., Nishitoh, H., Takami, Y. and Nakayama T.: The histone acetyltransferase p300/CBP-associated factor acts as an effective suppressor of secretory immunoglobulin synthesis in immature cells. *Microbiol. Immunol.* 59, 243-247, 2015.
- [137] Satrimafitrah, P., Barman, H. K., Ahmad, A., Nishitoh, H., Nakayama, T., Fukagawa, T. and Takami, Y.: RbAp48 is essential for viability of vertebrate cells and plays a role in chromosome stability. *Chromosome Res.* 24, 161-173, 2016.
- [138] Nakayama, T. and Nakayama, M.: Chromatin Conformation Change Code (4C) theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations. pp. 1-303, 2018. The 2nd Edition of the retirement commemorative monograph (Monograph) (the 1st Edition of the self-publishing Monograph published in 2015 is available from following new URL: <http://hdl.handle.net/10458/5293>) is available from following new URL: <http://hdl.handle.net/10458/6460>.
- [139] Kikuchi, H., Nakayama, M., Takami, Y., Kuribayashi, F. and Nakayama, T.: Possible involvement of Helios in controlling the immature B cell functions via transcriptional regulation of protein kinase Cs. *Results Immunol.* 1, 88-94, 2011.
- [140] Toyonaga, K., Kikuchi, H., Yamashita, K., Nakayama, M., Chijiwa, K. and Nakayama, T.: E2A participates in a fine control of pre-mature B cell apoptosis mediated by B cell receptor signaling via transcription regulations of survivin, IAP2 and caspase-8 genes. *FEBS J.* 276/5, 1418-1428, 2009.
- [141] Kikuchi, H., Yamashita, K., Nakayama, M., Toyonaga, K., Tsuneyoshi, I., Takasaki, M. and Nakayama, T.: Lacking of Aiolos accelerates pre-mature B cell apoptosis mediated by BCR signaling through elevation in cytochrome c release. *BBA-Molecular Cell Research* 1793, 1304-1314, 2009.
- [142] Kikuchi, H., Nakayama, M., Takami, Y., Kuribayashi, F. and Nakayama, T.: EBF1 acts as a powerful repressor of Blimp-1 gene expression in immature B cells. *Biochem. Biophys. Res. Commun.* 422, 780-785, 2012.
- [143] Takechi, S., Adachi, M. and Nakayama, T.: Cloning and characterization of the chick Oct binding factor OBF-1. *Biochim. Biophysica Acta* 1577, 466-470, 2002.
- [144] Kawano, J., Kotani T., Ogata, Y., Ohtaki, S., Takechi, S., Nakayama, T., Sawaguchi, A., Nagaike, R., Oinuma, T. and Sukanuma, T.: CALNUC (nucleobindin) is localized in the Golgi apparatus in insect cells. *Eur. J. Cell Biol.* 79, 208-217, 2000.
- [145] Takamatsu, H., Kodama, T., Imamura, A., Asai, K., Kobayashi, K., Nakayama, T., Ogasawara, N. and Watabe, K.: The *Bacillus subtilis* yabG gene is transcribed by SigK RNA polymerase during

sporulation, and yabG mutant spores have altered coat protein composition. *J. Bacteriol.* 182, 1883-1888, 2000.

[146] Ohi, T., Takechi, S., Itokazu, N., Shiomi, K., Sugimoto, S., Antoku, Y., Kato, K., Sugimoto, T., Nakayama, T. and Matsukura, S.: Two novel mutations in the adrenoleukodystrophy gene in two unrelated Japanese families and the long-term effect of bone marrow transplantation. *J. Neurol. Sci.* 177, 131-138, 2000.

[147] Yoshida, K., Kobayashi, K., Miwa, Y., Kang, C.-M., Matsunaga, M., Yamaguchi, H., Tojo, S., Yamamoto, M., Nishi, R., Ogasawara, N., Nakayama, T. and Fujita, Y.: Combined transcriptome and proteome analysis as a powerful approach to study genes under glucose repression in *Bacillus subtilis*. *Nucleic Acids Res.* 29, 683-692, 2001.

[148] Kawano, J., Nakayama, T., Takami, Y., Kotani, T., Sawaguchi, A., Nagaike, R., Oinuma, T. and Sukanuma, T.: A monoclonal antibody against insect CALNUC recognizes the prooncprotein EWS specifically in mammalian cells. Immunohistochemical and biochemical studies of the antigen in rat tissues. *Histochem. Cell Biol.* 115, 421-428, 2001.

[149] Seo, Y., Uchiyama, T., Matsuda, H., Shimizu, K., Takami, Y., Nakayama, T. and Takahama, K.: Mitochondrial DNA and STR typing of matter adhering to an earphone. *J. Forensic Sci.* 47, 605-608, 2002.

[150] Kawano, J., Nakayama, T., Kotani, T., Matsubayashi, H., Yamamoto, M.-T. and Sukanuma, T.: Identification and characterization of an insect homologue of the vertebrate Golgi apparatus protein 1 (MG-160/cysteine-rich fibroblast growth factor receptor/E-selectin ligand-1/latent transforming growth factor- β complex protein-1) with a Golgi-specific monoclonal antibody. *Histochem. Cell Biol.* 117, 381-389, 2002.

[151] Ohi, T., Saita, K., Takechi, S., Nabeshima, K., Tashiro, H., Shiomi, K., Sugimoto, S., Akematsu, T., Nakayama, T. Iwaki, T. and Matsukura, S.: Clinical features and neuropathological findings of familial amyotrophic lateral sclerosis with a His46Arg mutation in Cu/Zn superoxide dismutase. *J. Neurol. Sci.* 197, 73-78, 2002.

[152] Sakakibara, Y., Suiko, M., Pai, T. G., Nakayama, T., Takami, Y., Katafuchi, J. and Liu, M.-C.: Highly conserved mouse and human brain sulfotransferases: molecular cloning, expression, and functional characterization. *Gene* 285, 39-47, 2002.

[153] Seo, Y., Takami, Y., Takahama, K., Yoshizawa, M., Nakayama, T. and Yukawa, N.: A method for genotyping Y chromosome-linked DYS385a and DYS385b loci. *Legal Med.* 5, 228-232, 2003.

[154] Fukagawa, T., Nogami, M., Yoshikawa, M., Ikeno, M., Okazaki, T., Takami, Y., Nakayama, T. and Oshimura, M.: Dicer is essential for formation of the heterochromatin structure in vertebrate cells. *nature Cell Biology* 6, 784-791, 2004.

- [155] Matsushita, N., Takami, Y., Kimura, M., Tachiiri, S., Ishiai, M., Nakayama, T. and Takata, M.: Role of NAD-dependent deacetylases SIRT1 and SIRT2 in radiation and cisplatin-induced cell death in vertebrate cells. *Genes Cells* 10, 321-332, 2005.
- [156] Tanii, I., Yagura, T., Inagaki, N., Nakayama, T., Imaizumi, K. and Yoshinaga, K.: Preferential localization of rat GAPDS on the ribs of fibrous sheath of sperm flagellum and its expression during flagellar formation. *Acta Histochem. Cytochem.* 40, 19-26, 2007.
- [157] Hashimoto, H., Sonoda, E., Takami, Y., Kimura, H., Nakayama, T., Tachibana, M., Takeda, S. and Shinkai, Y.: Histone H1 variant, H1R is involved in DNA damage response. *DNA Repair* 6, 1584-1595, 2007.
- [158] Katayama, K., Jamhari, T., Mori, T., Kawahara, S., Miyake, K., Kodama, Y., Sugiyama, M., Kawahara, Y., Nakayama, T., Maruyama, M. and Muguruma, M.: Angiotensin-I converting enzyme inhibitory peptide derived from porcine skeletal muscle myosin and its antihypertensive activity in spontaneously hypertensive rats. *J. Food Sci.* 72, S702-S706, 2007.
- [159] Katayama, K., Anggraeni, H. E., Mori, T., Ahmed, A. M., Kawahara, S., Sugiyama, M., Nakayama, T., Maruyama, M. and Muguruma, M.: Porcine skeletal muscle troponin is a good source of peptides with Angiotensin-I converting enzyme inhibitory activity and antihypertensive effects in spontaneously hypertensive rats. *J. Agric. Food Chem.* 56, 355-360, 2008.
- [160] Nakade, K., Kamishima, R., Inoue, Y., Ahmed, A., Kawahara, S., Nakayama, T., Maruyama, M., Numata, M., Ohta, K., Aoki, T. and Muguruma, M.: Identification of an antihypertensive peptide derived from chicken bone extract. *Animal Science Journal* 79, 710-715, 2008.
- [161] Nishijima, H., Takami, Y., Nakayama, T., Adachi, N. and Shibahara, K.: Improved applications of the tetracycline-regulated gene depletion system. *BioScience Trends* 5, 161-167, 2009.
- [162] Hashimoto, H., Takami, Y., Sonoda, E., Iwasaki, T., Iwano, H., Tachibana, M., Takeda, S., Nakayama, T., Kimura, H. and Shinkai, Y.: H1 null vertebrate cells exhibit altered nucleosome architecture. *Nucleic Acids Res.* 38, 3533-3545, 2010.
- [163] Abe, T., Sugimura, K., Hosono, Y., Takami, Y., Akita, M., Yoshimura, A., Tada, S., Nakayama, T., Murofushi, H., Okumura, K., Takeda, S., Horikoshi, M., Seki, M. and Enomoto, T.: The histone chaperone facilitates chromatin transcription (FACT) protein maintains normal replication fork rates. *J. Biol. Chem.* 286, 30504-30512, 2011.
- [164] Nakanishi, O., Oyanagi, M., Kuwano, Y., Tanaka, T., Nakayama, T., Mitsui, H., Nabeshima, Y. and Ogata, K.: Molecular cloning and nucleotide sequences of cDNAs specific for rat liver ribosomal proteins S17 and L30. *Gene* 35, 289-296, 1985.
- [165] Yoshizaki, K., Tseng, L.T., Nakayama, T., Kishimoto, S. and Kishimoto, T.: Isolation of B cell-derived cell differentiation factor (B-BCDF). In *cellular and molecular biology of lymphokines* (eds. by C. Sorg and A. Schimpl), Academic Press, pp241-245, 1985.

- [166] Yamamoto, Y., Ogawa, T., Shinagawa, H., Nakayama, T., Matsuo, H. and Ogawa, H.: Determination of the initiation sites of transcription and translation of the *uvrD* gene of *Escherichia coli*. *J. Biochem.* 99, 1579-1590, 1986.
- [167] Kimura, S., Kotani, T., McBride, O.W., Umeki, K., Hirai, K., Nakayama, T. and Ohtaki, S.: Human thyroid peroxidase: Complete cDNA and protein sequence, chromosome mapping and identification of two alternately spliced mRNAs. *Proc. Natl. Acad. Sci. USA* 84, 5555-5559, 1987.
- [168] Tamura, S., Kuwano, U., Nakayama, T., Tanaka, S., Tanaka, T. and Ogata, K.: Molecular cloning and nucleotide sequence of cDNA specific for rat ribosomal protein L5. *Eur. J. Biochem.* 168, 83-87, 1987.
- [169] Kawano, J., Oinuma, T., Nakayama, T. and Suganuma, T.: Characterization of β 1 \rightarrow 4 galactosyltransferase purified from rat liver microsomes. *J. Biochem.* 111, 568-572, 1992.
- [170] Seo, Y., Nakayama, T. and Takahama, K.: Immunoaffinity purification and characterization of a human liver-specific antigen. *J. Legal Med* 47, 1-5, 1993.
- [171] Kotani, T., Umeki, K., Yamamoto, I., Takeuchi, M., Takechi, S., Nakayama, T. and Ohtaki, S.: Nucleotide sequence of the cDNA encoding mouse thyloid peroxidase. *Gene* 123, 289-290, 1993.
- [172] Nakamoto, M., Ozawa, M., Jacinto, S., Furukawa, T., Natori, Y., Shirahama, H., Yonezawa, S., Nakayama, T. and Muramatsu, T.: Mouse heparin binding protein-44 (HBP-44) associates with brushin, a high-molecular-weight glycoprotein antigen common to the kidney and teratocarcinomas. *J. Biochem.* 114, 344-349, 1993.
- [173] Take, M., Tsutsui, J., Obana, H., Ozawa, M., Nakayama, T., Maruyama, I., Arima, T. and Muramatsu, T.: Identification of nucleolin as a binding protein for midkine (MK) and heparin-binding growth associated molecule (HB-GAM). *J. Biochem.* 116, 1063-1068, 1994.
- [174] Yamazoe, M., Shirahige, K., Rashid, M. B., Kaneko, Y., Nakayama, T., Ogasawara, N. and Yoshikawa, H.: A protein which binds preferentially to single-stranded core sequence of autonomously replicating sequence is essential for respiratory function in mitochondria of *Saccharomyces cerevisiae*. *J. Biol. Chem.* 269, 15244-15252, 1994.
- [175] Matsuoka, Y., Takechi, S., Nakayama, T. and Yoneda, Y.: Exogenous histone H1 injection into mitotic cells disrupts synchronous progression of mitotic events by delaying chromosome decondensation. *J. Cell Sci.* 107, 693-701, 1994.
- [176] Sakakibara, Y., Takami, Y., Zwieb, C., Nakayama, T., Suiko, M., Nakajima, H. and Liu, M.-C.: Purification, characterization, and molecular cloning of a novel rat liver Dopa/tyrosine sulfotransferase. *J. Biol. Chem.* 270, 30470-30478, 1995.
- [177] Ogura, K., Takechi, S., Nakayama, T. and Yamamoto, M.-T.: Molecular structure of the transposable element *ninja* in *Drosophila simulans*. *Genes & Genet. Syst.* 71, 1-8, 1996.

- [178] Sakakibara, Y., Katafuchi, J., Takami, Y., Nakayama, T., Suiko, M., Nakajima, H. and Liu, M.-C.: Manganese-dependent Dopa/tyrosine sulfation in HepG2 human hepatoma cells: novel Dopa/tyrosine sulfotransferase activities associated with the human monoamine-form phenol sulfotransferase. *Biochim. Biophys. Acta* 1355, 102-106, 1997.
- [179] Sakakibara, Y., Takami, Y., Nakayama, T., Suiko, M. and Liu, M.-C.: Localization and functional analysis of the substrate specificity/catalytic domains of human M-form and P-form phenol sulfotransferases. *J. Biol. Chem.* 273, 6242-6247, 1998.
- [180] Yanagisawa, K., Sakakibara, Y., Suiko, M., Takami, Y., Nakayama, T., Nakajima, H., Takayanagi, K., Natori, Y. and Liu, M.-C.: cDNA cloning, expression, and characterization of the human bifunctional ATP/adenosine 5'-phosphosulfate kinase enzyme. *Biosci. Biotechnol. Biochem.* 62, 1037-1040, 1998.
- [181] Sakakibara, T., Yanagisawa, K., Takami, Y., Nakayama, T., Suiko, M. and Liu, M.-C.: Molecular cloning, expression, and functional characterization of novel mouse sulfotransferase. *Biochem. Biophys. Res. Commun.* 247, 681-686, 1998.
- [182] Sakakibara, Y., Yanagisawa, K., Katafuchi, J., Ringer, D. P., Takami, Y., Nakayama, T., Suiko, M. and Liu, M.-C.: Molecular cloning, expression, and characterization of novel human SULT1C sulfotransferases that catalyze the sulfonation of N-hydroxy-2-acetylaminofluorene. *J. Biol. Chem.* 273, 33929-33935, 1998.
- [183] Kamei, Y., Yuba, S., Nakayama, T. and Yoneda, Y.: Three classes of the α -subunit of the nuclear pore-targeting complex (importin- α) are differentially expressed in adult mouse tissues. *J. Histochem. Cytochem.* 47, 363-372, 1999.
- [184] Takamatsu, H., Kodama, T., Nakayama, T., Watabe, K.: Characterization of the *yrbA* gene of *Bacillus subtilis*, involved in resistance and germination of spores. *J. Bacteriol.* 181, 4986-4994, 1999.
- [185] Yanagita, T., Kobayashi, H., Yamamoto, R., Takami, Y., Yokoo, H., Yuhi, T., Nakayama, T. and Wada, A.: Protein kinase C and the opposite regulation of sodium channel α - and β 1-subunit mRNA levels in adrenal chromaffin cells. *J. Neurochem.* 73, 1749-1757, 1999.
- [186] Nakajima, Y., Yamamoto, T., Nakayama, T. and Nakanishi, S.: A relationship between protein kinase C phosphorylation and calmodulin binding to the metabotropic glutamate receptor subtype 7. *J. Biol. Chem.* 274, 27573-27577, 1999.
- [187] Seo, Y., Takami, Y., Nakayama, T. and Takahama, K.: Y chromosome DNA polymorphisms and their haplotypes in Japanese population. *Legal Med.* 1, 145-149, 1999.
- [188] Nakayama, M. and Nakayama, T.: Protein and mRNA levels of IgM H- and L-chains artificially and excessively accumulated in HDAC2-deficient DT40 mutants are dramatically reduced through various generations during continuous cultivation. pp. 1-34, 2018. The 2nd Edition of the Paper-1 (the 1st Edition of the Paper-1 published in 2017 is available from following new URL: <http://hdl.handle.net/10458/5933>) is available from following new URL:

<http://hdl.handle.net/10458/6500>.

The 2nd Edition of the Paper-1 is also the revised Edition of Chapter-2 of the 2nd Edition of the Monograph (the 1st Edition of the Monograph published in 2015 is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, M. and Nakayama, T.: Protein and mRNA levels of IgM H- and L-chains artificially accumulated in HDAC2-deficient DT40 mutants are dramatically reduced through various generations during continuous cultivations. In: Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations, Nakayama, T. and Nakayama, M. (Eds.), pp. 11-44, 2018. The 2nd Edition of the Monograph is available from following new URL: <http://hdl.handle.net/10458/6460>.

[189] Nakayama, M. and Nakayama, T.: Generation of Pax5-deficient DT40 mutant cells, Pax5(-), and protein and mRNA levels of IgM H- and L-chains artificially and excessively accumulated in Pax5(-) DT40 mutants are rapidly and dramatically reduced through various generations during continuous cultivation. pp. 1-28, 2018. The 2nd Edition of the Paper-2 (the 1st Edition of the Paper-2 published in 2017 is available from following new URL: <http://hdl.handle.net/10458/5934>) is available from following new URL: <http://hdl.handle.net/10458/6503>.

The 2nd Edition of the Paper-2 is also the revised Edition of Chapter-3 of the 2nd Edition of the Monograph (the 1st Edition of the Monograph published in 2015 is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, M. and Nakayama, T.: Generation of Pax5-deficient DT40 mutants, Pax5(-), and protein and mRNA levels of IgM H- and L-chains artificially accumulated in Pax5(-) are rapidly and dramatically reduced through various generations during continuous cultivation. In: Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations, Nakayama, T. and Nakayama, M. (Eds.), pp. 45-71, 2018. The 2nd Edition of the Monograph is available from following new URL: <http://hdl.handle.net/10458/6460>.

[190] Nakayama, M. and Nakayama, T.: IgM H- and L-chains artificially and excessively accumulated in HDAC2(-/-) DT40 mutants are gradually and dramatically reduced in distinct ways in individual mutant clones through various generations during continuous cultivation. pp. 1-38, 2018. The 2nd Edition of the Paper-3 (the 1st Edition of the Paper-3 published in 2017 is available from following new URL: <http://hdl.handle.net/10458/5935>) is available from following new URL: <http://hdl.handle.net/10458/6504>.

The 2nd Edition of the Paper-3 is also the revised Edition of Chapter-4 of the 2nd Edition of the Monograph (the 1st Edition of the Monograph published in 2015 is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, M. and Nakayama, T.: IgM H- and L-chains

accumulated artificially and excessively in HDAC2(-/-) DT40 mutants are dramatically reduced in distinct ways in individual mutant clones through various generations during continuous cultivations. In: Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations, Nakayama, T. and Nakayama, M. (Eds.), pp. 72-104, 2018. The 2nd Edition of the Monograph is available from following new URL: <http://hdl.handle.net/10458/6460>.

[191] Nakayama, M. and Nakayama, T.: Fundamental and distinct ways for irreversible creation of chromatin structure plasticity of proximal 5'-upstream regions of Pax5, Aiolos, EBF1 and OBF1 genes with epigenetic modifications for gain of new cell function to exclude accumulated IgM H- and L-chains in individual clones of HDAC2(-/-) DT40 mutants through various generations during continuous cultivation. pp. 1-70, 2018. The 2nd Edition of the Paper-4 (the 1st Edition of the Paper-4 published in 2017 is available from following new URL: <http://hdl.handle.net/10458/5936>) is available from following new URL: <http://hdl.handle.net/10458/6505>.

The 2nd Edition of the Paper-4 is also the revised Edition of Chapter-5 of the 2nd Edition of the Monograph (the 1st Edition of the Monograph published in 2015 is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, M. and Nakayama, T.: A fundamental way for irreversible creation of chromatin structure plasticity with epigenetic modifications for gaining new cell function to exclude IgM H- and L-chains accumulated in HDAC2(-/-) DT40 mutants through various generations during continuous cultivation. In: Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations, Nakayama, T. and Nakayama, M. (Eds.), pp. 105-167, 2018. The 2nd Edition of the Monograph is available from following new URL: <http://hdl.handle.net/10458/6460>.

[192] Nakayama, T. and Nakayama, M.: Chromatin conformation change code (4C) theory: A bio-system for gaining un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through various generations. pp. 1-43, 2018. The 2nd Edition of the Paper-5 (the 1st Edition of the Paper-5 published in 2017 is available from following new URL: <http://hdl.handle.net/10458/5937>) is available from following new URL: <http://hdl.handle.net/10458/6506>.

The 2nd Edition of the Paper-5 is also the revised Edition of Chapter-6 of the 2nd Edition of the Monograph (the 1st Edition of the Monograph was published in 2015 and is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, M. and Nakayama, T.: Chromatin conformation change code (4C) theory: A bio-system for gaining un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through various generations. In: Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed

and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations, Nakayama, T. and Nakayama, M. (Eds.), pp. 168-202, 2018. The 2nd Edition of the Monograph is available from following new URL: <http://hdl.handle.net/10458/6460>.

[193] Nakayama, T. and Nakayama, M.: An all-inclusive review: Chromatin conformation change code (4C) theory on a bio-system to gain un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through various generations in higher eukaryotes. pp. 1-75, 2018. The 2nd Edition of the Paper-6 (the 1st Edition of the Paper-6 published in 2017 is available from following new URL: <http://hdl.handle.net/10458/5941>) is available from following new URL: <http://hdl.handle.net/10458/6507>.

The 2nd Edition of the Paper-6 is also the revised Edition of Chapter-7 of the 2nd Edition of the Monograph (the 1st Edition of the Monograph published in 2015 and is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, T. and Nakayama, M.: All-inclusive review and history on the chromatin conformation change code (4C) theory: A bio-system for gaining un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through various generations. In: Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations, Nakayama, T. and Nakayama, M. (Eds.), pp. 203-260, 2018. The 2nd Edition of the Monograph is available from following new URL: <http://hdl.handle.net/10458/6460>.

[194] Nakayama, M. and Nakayama, T.: IgM H- and L-chains accumulated excessively in HDAC2(-/-) DT40 mutants are dramatically reduced in distinct ways in individual mutant clones through various generations during continuous cultivation. Current Topics in Biochemical Research. 18, 11-25, 2017.

The article is available from following Website and new URL.

http://www.researchtrends.net/tia/title_issue.asp?id=40&in=0&vn=18&type=3

and <http://hdl.handle.net/10458/6375>.

[195] Nakayama, M. and Nakayama, T.: Irreversible creation of chromatin structure plasticity of proximal 5'-upstream regions of Pax5, Aiolos, EBF1 and OBF1 genes with epigenetic modifications to exclude IgM H- and L-chains accumulated in individual clones of HDAC2(-/-) DT40 mutants through various generations during continuous cultivation. Current Topics in Biochemical Research. 18, 33-56, 2017.

The article is available from following Website and new URL.

http://www.researchtrends.net/tia/title_issue.asp?id=40&in=0&vn=18&type=3

and <http://hdl.handle.net/10458/6376>.

[196] Nakayama, T. and Nakayama, M.: Chromatin conformation change code (4C) theory: A bio-system to gain un-programmed and new cell functions by means of irreversible creation of chromatin structure

plasticity with epigenetic modifications through various generations. *Current Topics in Biochemical Research*. 18, 65-86, 2017.

The article is available from following Website and new URL.

http://www.researchtrends.net/tia/title_issue.asp?id=40&in=0&vn=18&type=3

and <http://hdl.handle.net/10458/6377>.

[197] Nakayama, T. and Nakayama, M.: A comprehensive review on the chromatin conformation change code (4C) theory: A bio-system for gaining un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through various generations in higher eukaryotes. pp.1-76, 2019. The 3rd Edition of the Paper-6 (the 1st and 2nd Editions published in 2017 and 2018 are respectively available from following new URLs: <http://hdl.handle.net/10458/5941> and <http://hdl.handle.net/10458/6507>) is available from following new URL: <http://hdl.handle.net/10458/6613>.

The 3rd Edition of the Paper-6 is also the revised Edition of Chapter-7 of the 2nd Monograph (the 1st Monograph published in 2015 is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, T. and Nakayama, M.: All-inclusive review and history on the chromatin conformation change code (4C) theory: A bio-system for gaining un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through various generations. In: *Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations*, Nakayama, T. and Nakayama, M. (Eds.), pp. 203-260, 2018. The 2nd Monograph is available from following new URL: <http://hdl.handle.net/10458/6460>.

The 1st Japanese Edition (with some fundamental concepts on life phenomena) of the 3rd Edition of the Paper-6 was published in 2019 and is available from following new URL: <http://hdl.handle.net/10458/6615>.

[198] Nakayama, T.: *Memoirs of a life scientist: Interpretations and opinions on life phenomena, and experiences and contributions in life science field ~ The chromatin conformation change code (4C) theory and previous researches*. pp. 1-79, 2020. The 2nd Edition of the Paper-7 (the 1st Edition published in 2019 is available from following new URL: <http://hdl.handle.net/10458/6614>) is available from following new URL: <http://hdl.handle.net/10458/6949>.

The 2nd Edition of the Paper-7 is also the revised Edition of Chapter-8 of the 2nd Monograph (the 1st Monograph published in 2015 is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, T.: Personal closing remarks on the occasion of retirement: The real course and history of our research prior to the 4C theory, personal interpretation on life phenomena, and experiences and opinions as a life scientist. In: *Chromatin Conformation Change Code (4C) Theory on Gain of*

Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations, Nakayama, T. and Nakayama, M. (Eds.), pp. 261-303, 2018. The 2nd Monograph is available from following new URL: <http://hdl.handle.net/10458/6460>.

[199] Nakayama, T. and Nakayama, M.: A comprehensive and detailed review on the chromatin conformation change code (4C) theory: A theory on ways to gain un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through numerous generations in higher eukaryotes. pp. 1-53, 2020. The 4th Edition of Paper-6 (1st, 2nd and 3rd Editions published in 2017, 2018 and 2019 are available from following New URLs: <http://hdl.handle.net/10458/5941>, <http://hdl.handle.net/10458/6507> and <http://hdl.handle.net/10458/6613>) is available from following New URL: <http://hdl.handle.net/10458/6902>.

The 4th Edition of the Paper-6 is also the revised Edition of Chapter-7 of the 2nd Edition of the Monograph (the 1st Edition of the Monograph published in 2015 is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, T. and Nakayama, M.: All-inclusive review and history on the chromatin conformation change code (4C) theory: A bio-system for gaining un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through various generations. In: Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations, Nakayama, T. and Nakayama, M. (Eds.), pp. 203-260, 2018. The 2nd Edition of the Monograph is available from following new URL: <http://hdl.handle.net/10458/6460>.

The 1st and 2nd Japanese Editions (with some fundamental concepts on life phenomena) of the 3rd and 4th Editions of the Paper-6 published in 2019 and 2020 are available from following new URLs: <http://hdl.handle.net/10458/6615> and <http://hdl.handle.net/10458/6950>.

[200] Zimmer, C. and Emlen, D. J.: The Japanese Version of Evolution: Making Sense of Life (2017).

[201] Cooper, V. S. and Lenski, R. E.: The population genetics of ecological specialization in evolving *Escherichia coli* populations. *Nature*, 407, 736-739, 2000.

[202] Blount, Z. D., Borland, C. Z. and Lenski, R. E.: Historical contingency and the evolution of a key innovation in a population of *Escherichia coli*. *Proc. Natl. Sci. USA*, 105, 7899-7906, 2008.

[203] Kawasaki, M.: Temperature adaptation in *Drosophila melanogaster*. *Konchu*, 44, 530-536, 1976 (in Japanese).

[204] Rice, T. and Salt, G. W.: Speciation via disruption selection on habitat preference: Experimental evidence. *The American Naturalist*, 131, 911-917, 1988.

[205] Nono, M., Kishimoto, S., Sato-Carlton, A., Carlton, P. M., Nishida, E. and Uno, M.:

Intestine-to-germline transmission of epigenetic information intergenerationally ensures systemic stress resistance in *C. elegans*. *Cell Reports* 30, 3207-3217, 2020.

[206] Wenzel, D., Palladino, F., and Jedrusik-Bode, M.: Epigenetics in *C. elegans*: facts and challenges. *Genesis* 49, 647–661, 2011.

[207] Gonzalez-Aguilera, C., Palladino, F., and Askjaer, P.: *C. elegans* epigenetic regulation in development and aging. *Brief. Funct. Genomics* 13, 223–234, 2014.

[208] Jin, C., Li, J., Green, C.D., Yu, X., Tang, X., Han, D., Xian, B., Wang, D., Huang, X., Cao, X., et al.: Histone demethylase UTX-1 regulates *C. elegans* life span by targeting the insulin/IGF-1 signaling pathway. *Cell Metab.* 14, 161–172, 2011.

[209] Siebold, A.P., Banerjee, R., Tie, F., Kiss, D.L., Moskowitz, J., and Harte, P.J.: Polycomb Repressive Complex 2 and Trithorax modulate *Drosophila* longevity and stress resistance. *Proc. Natl. Acad. Sci. USA* 107, 169–174, 2010.

[210] van Oosten-Hawle, P., Porter, R.S., and Morimoto, R.I.: Regulation of organismal proteostasis by transcellular chaperone signaling. *Cell* 153, 1366–1378, 2013.

[211] Tatum, M.C., Ooi, F.K., Chikka, M.R., Chauve, L., Martinez-Velazquez, L.A., Steinbusch, H.W.M., Morimoto, R.I., and Prahlad, V.: Neuronal serotonin release triggers the heat shock response in *C. elegans* in the absence of temperature increase. *Curr. Biol.* 25, 163–174, 2015.

[212] Greer, E.L., Maures, T.J., Hauswirth, A.G., Green, E.M., Leeman, D.S., Maro, G.S., Han, S., Banko, M.R., Gozani, O., and Brunet, A.: Members of the H3K4 trimethylation complex regulate lifespan in a germline-dependent manner in *C. elegans*. *Nature* 466, 383–387, 2010.

[213] Dias, B.G., and Ressler, K.J.: Parental olfactory experience influences behavior and neural structure in subsequent generations. *Nat. Neurosci.* 17, 89–96, 2014.

[214] Kishimoto, S., Uno, M., Okabe, E., Nono, M., and Nishida, E.: Environmental stresses induce transgenerationally inheritable survival advantages via germline-to-soma communication in *Caenorhabditis elegans*. *Nat. Commun.* 8, 14031, 2017.

[215] Vaiserman, A.M., Koliada, A.K., and Jirtle, R.L.: Non-genomic transmission of longevity between generations: potential mechanisms and evidence across species. *Epigenetics Chromatin* 10, 38, 2017.

[216] Rechavi, O., Hourri-Zeevi, L., Anava, S., Goh, W.S.S., Kerk, S.Y., Hannon, G.J., and Hobert, O.: Starvation-induced transgenerational inheritance of small RNAs in *C. elegans*. *Cell* 158, 277–287, 2014.

[217] Posner, R., Toker, I.A., Antonova, O., Star, E., Anava, S., Azmon, E., Hendricks, M., Bracha, S., Gingold, H., and Rechavi, O.: Neuronal Small RNAs Control Behavior Transgenerationally. *Cell* 177, 1814–1826.e15, 2019.

[218] Klosin, A., Casas, E., Hidalgo-Carcedo, C., Vavouri, T. and Lehner, B.: Transgenerational transmission of environmental information in *C. elegans*. *Science*, 356, 320-323, 2017.

[219] Zenk, F., Loeser, E., Schiavo, R., Kilpert, F., Bogdanovic, O. and Lovino, N.: Germ line-inherited

H3K27me3 restricts enhancer function during maternal-to-zygotic transition. *Science*, 357, 212-216, 2017.

[220] Moore, M. S., Keletsky, R. and Murphy, C. T.: Piwi/PRG-1 argonaute and TGF- β mediate transgenerational learned pathogenic avoidance. *Cell*, 177, 1827-1841, 2019.

[221] Nakayama, T. and Nakayama, M.: Chromatin conformation change code (4C) theory: A bio-system for gaining un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through numerous generations (3rd Edition). pp.1-51, 2021. The 3rd Edition of the Paper-5 (the 1st and 2nd Editions of the Paper-5 published in 2017 and 2018 are available from following new URLs: <http://hdl.handle.net/10458/5937> and <http://hdl.handle.net/10458/6506>) is available from following new URL: <http://hdl.handle.net/10458/6506>.

The 3rd Edition of the Paper-5 is also the revised Edition of Chapter-6 of the 2nd Edition of the Monograph (the 1st Edition of the Monograph was published in 2015 and is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, M. and Nakayama, T.: Chromatin conformation change code (4C) theory: A bio-system for gaining un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through various generations. In: *Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations*, Nakayama, T. and Nakayama, M. (Eds.), pp.168-202, 2018. The 2nd Edition of the Monograph is available from following new URL: <http://hdl.handle.net/10458/6460>.

[222] Nakayama, T. and Nakayama, M.: A comprehensive and detailed review on the chromatin conformation change code (4C) theory: A theory on ways to gain un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through numerous generations in higher eukaryotes (5th Edition). pp.1-85, 2021. The 5th Edition of the Paper-6 (the 1st, 2nd, 3rd and 4th Editions published in 2017, 2018, 2019 and 2020 are available from following new URLs: <http://hdl.handle.net/10458/5941>, <http://hdl.handle.net/10458/6507>, <http://hdl.handle.net/10458/6613> and <http://hdl.handle.net/10458/6902>) is available from following new URL: <http://hdl.handle.net/10458/6902>.

The 5th Edition of the Paper-6 is also the revised Edition of Chapter-7 of the 2nd Edition of the Monograph (the 1st Monograph published in 2015 is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, T. and Nakayama, M.: All-inclusive review and history on the chromatin conformation change code (4C) theory: A bio-system for gaining un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through various generations. In: *Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of*

Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations, Nakayama, T. and Nakayama, M. (Eds.), pp.203-260, 2018. The 2nd Edition of the Monograph is available from following new URL: <http://hdl.handle.net/10458/6460>.

[223] Nakayama, T.: Memoirs of a life scientist: Interpretations and opinions on life phenomena, and experiences and contributions in life science field ~ The chromatin conformation change code (4C) theory and previous researches (3rd Edition) ~. pp.1-83, 2021. The 3rd Edition of the Paper-7 (the 1st and 2nd Editions published in 2019 and 2020 are available from following new URLs: <http://hdl.handle.net/10458/6614> and <http://hdl.handle.net/10458/6949>) is available from following new URL: <http://hdl.handle.net/10458/6949>.

The 3rd Edition of the Paper-7 is also the revised Edition of Chapter-8 of the 2nd Monograph (the 1st Monograph published in 2015 is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, T.: Personal closing remarks on the occasion of retirement: The real course and history of our research prior to the 4C theory, personal interpretation on life phenomena, and experiences and opinions as a life scientist. In: Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations, Nakayama, T. and Nakayama, M. (Eds.), pp. 261-303, 2018. The 2nd Monograph is available from following new URL: <http://hdl.handle.net/10458/6460>.

[224] Nakayama, T. and Nakayama, M.: The Japanese review on the chromatin conformation change code (4C) theory: A theory on ways to acquire un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through numerous generations in higher eukaryotes (3rd Edition). pp.1-123, 2021. The 3rd Japanese Edition (with some fundamental concepts on life phenomena) of the Paper-8 (the 1st and 2nd Editions published in 2019 and 2020 are available from following new URLs: <http://hdl.handle.net/10458/6615> and <http://hdl.handle.net/10458/6950>) is available from following new URL: <http://hdl.handle.net/10458/6950>.

The 3rd Japanese Edition of the Paper-8 is also the revised Edition of Chapter-7 of the 2nd Edition of the Monograph (the 1st Monograph published in 2015 is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, T. and Nakayama, M.: All-inclusive review and history on the chromatin conformation change code (4C) theory: A bio-system for gaining un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through various generations. In: Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations, Nakayama, T. and Nakayama, M. (Eds.), pp.203-260, 2018. The 2nd Edition of the Monograph is available from

following new URL: <http://hdl.handle.net/10458/6460>.

[225] Nakayama, T. and Nakayama, M.: A comprehensive review on the chromatin conformation change code (4C) theory: A theory on ways of higher eukaryotes to gain un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through numerous generations. Current Topics in Biochemical Research, 22, pp.43-90, 2021.

This review article is available from following Website and new URL:

http://www.researchtrends.net/tia/title_issue.asp?id=40&in=0&vn=22&type=3 and
<http://hdl.handle.net/10458/00010392>

This review article is the revised Edition of the 5th Edition of Paper-6 on the comprehensive review (the 1st, 2nd, 3rd and 4th Editions published in 2017, 2018, 2019 and 2020 are available from following new URLs: <http://hdl.handle.net/10458/5941>, <http://hdl.handle.net/10458/6507>, <http://hdl.handle.net/10458/6613> and <http://hdl.handle.net/10458/6902>) is available from following new URL: <http://hdl.handle.net/10458/6902>.

This review article is also the revised Edition of Chapter-7 of the 2nd Edition of the Monograph (the 1st Monograph published in 2015 is available from following new URL: <http://hdl.handle.net/10458/5293>) as follows: Nakayama, T. and Nakayama, M.: All-inclusive review and history on the chromatin conformation change code (4C) theory: A bio-system for gaining un-programmed and new cell functions by means of irreversible creation of chromatin structure plasticity with epigenetic modifications through various generations. In: Chromatin Conformation Change Code (4C) Theory on Gain of Un-programmed and New Cell Functions by Means of Irreversible Creation of Chromatin Structure Plasticity with Epigenetic Modifications through Various Generations, Nakayama, T. and Nakayama, M. (Eds.), pp.203-260, 2018. The 2nd Edition of the Monograph is available from following new URL: <http://hdl.handle.net/10458/6460>.

Figure legends

Figure 1. Ways to regulate gene expressions of IgM H- and L-chains through regulation of gene expressions of Pax5, Aiolos, EBF1, OBF1 and other genes in wild-type DT40 cells or in all and individual clones of HDAC2(-/-) DT40 mutants at early and later stages of continuous cultivation.

In DT40 cells (W), HDAC2 indirectly regulates gene expressions of IgM H- and L-chains through opposite regulation of gene expressions of Pax5, Aiolos, OBF1 and EBF1, and Ikaros and E2A. In all clones of HDAC2(-/-) mutants, at the early (E) cultivation stage, IgM H- and L-chains are excessively accumulated attributed to their dramatically increased gene expressions caused by drastically decreased gene expressions of Pax5, Aiolos and EBF1, all of which down-regulate gene expressions of the two immunoglobulin proteins. In individual clones of HDAC2(-/-) mutants, at the later (L) cultivation stage, the accumulated IgM H- and L-chains are dramatically decreased attributed to their drastically decreased gene expressions in almost the same changing pattern, caused by dramatically increased or decreased gene expressions of Pax5, Aiolos and EBF1 or OBF1 in distinct ways. Clone cl.2-1 seems to be the OBF1-dependent type, clones cl.2-2, cl.2-3, cl.2-4 and cl.2-5 seem to be the Pax5- and Aiolos-dependent type, and clone cl.2-6 seems to be the Pax5-, Aiolos- and EBF1-dependent type. The figure is identical with Fig. 1 of Refs. 198 and 223. The figure is also identical with Fig. 1 of Refs. 193, 197, 199, 222 plus 225 and Fig. 6 of Ref. 194 and is a set of Figs. 8, 9 and 10 of Ref. 190. These figures were first presented in Ref. 138.

Figure 2. Alterations in amounts of IgM H- and L-chains in HDAC2(-/-) DT40 mutants during continuous cultivation.

2D-PAGE was performed on total cellular proteins prepared from HDAC2(-/-) mutant cells at the early (E), middle (M) and later (L) cultivation stages and from DT40 cells (W). H and L indicate IgM H- and L-chains. The figure is identical with Fig. 2 of Refs. 198 and 223. The figure is also identical with Fig. 2 of Refs. 193, 197, 199, 219 plus 225 and Fig. 1 of Ref. 196 and Fig. 1 of Refs. 192 plus 221 and Fig. 1 of Ref. 188. These figures were first presented in Ref. 138.

Figure 3. Alterations in amounts of IgM H-chain in HDAC2(-/-) DT40 mutants during continuous cultivation.

Immuno-electron microscopy, using anti-chicken IgM H-chain antiserum, was performed on HDAC2(-/-) mutant cells at the early (E) and later (L) cultivation stages and on DT40 cells (W) in wide range (A) and for a single cell (B). Large amounts of accumulated IgM H-chain were detected only at the early (E) stage in HDAC2(-/-) mutant cells. The figure is identical with Fig. 3 of Refs. 198 and 223. The figure is also identical with Fig. 3 of Refs. 193, 197, 199, 222 plus 225 and Fig. 4 of Ref. 188. These figures were first presented in Ref. 138.

Figure 4. Alterations in gene expressions of IgM H- and L-chains, and core histones in HDAC2(-/-) DT40 mutants during continuous cultivation.

Total RNAs were extracted from three individual clones of HDAC2(-/-) mutants at the early (E), middle (M) and later (L) cultivation stages and from DT40 cells (W). RT-PCR was performed on total RNAs, using appropriate primers for HDAC2, whole (IgM Hc), secreted (IgM Hs) plus membrane-bound (IgM Hm) forms of IgM H-chain and IgM L-chain (IgM L), and core histones H2A, H2B, H3 and H4. Some data for DT40 (W) and HDAC2(-/-) at the early (E) stage were identical with those of Ref. 124. The figure is identical with Fig. 4 of Refs. 198 and 223. The figure is also identical with Fig. 4 of Refs. 193, 197, 199, 222 plus 225 and Fig. 5 of Ref. 188. These figures were first presented in Ref. 138.

Figure 5. Alterations in amounts of IgM H- and L-chains in Pax5(-) DT40 mutants during continuous cultivation.

Western blotting was performed on total cellular proteins prepared from three Pax5(-) mutant clones at the early (E), middle (M) and later (L) cultivation stages and from DT40 cells (W), using anti-chicken IgM L-chain antiserum that cross-reacts with IgM H-chain. Relative levels of IgM H-chain and large (high) plus small (low) forms of IgM L-chain are shown in the lower panel. The figure is identical with Fig. 5 of Refs. 198 and 223. The figure is also identical with Fig. 5 of Refs. 193, 197, 199, 222 plus 225 and Fig. 3 of Ref. 189. These figures were first presented in Ref. 138.

Figure 6. Morphology of Pax5(-) DT40 mutant cells during continuous cultivation.

Microscopy was performed on Pax5(-) mutant cells at the early (E) and later (L) cultivation stages and on DT40 cells (W) in distinct ranges of vision with magnified visions. The figure is identical with Fig. 6 of Refs. 198 and 223. The figure is identical with Fig. 6 of Refs. 193, 197, 199, 222 plus 225 and Fig. 5 of Ref. 189. These figures were first presented in Ref. 138.

Figure 7. Alterations in gene expressions of IgM H- and L-chains in Pax5(-) DT40 mutants during continuous cultivation.

RT-PCR was performed on total RNAs prepared from three Pax5(-) mutant clones at the first (F), early (E), middle (M) and later (L) cultivation stages and from DT40 cells (W), using appropriate primers for Pax5, and whole (IgM Hc), secreted (IgM Hs) plus membrane-bound (IgM Hm) forms of IgM H-chain and IgM L-chain (IgM L). Their relative mRNA levels are shown in the lower panel. The figure is identical with Fig. 7 of Refs. 198 and 223. The figure is also identical with Fig. 7 of Refs. 193, 197, 199, 222 plus 225 and Fig. 6 of Ref. 189. These figures were first presented in Ref. 138.

Figure 8. Genomic organization of chicken HDAC2 gene, generation of HDAC2(-/-) DT40 mutants, and alterations in protein and mRNA levels of IgM H- and L-chains in individual clones of HDAC2(-/-) DT40 during continuous cultivation.

A) Schematic presentation of chicken HDAC2 genomic locus (top) with enlarged drawing of its intact allele (middle) and targeted alleles (two bottoms). Locations of exons 1-16, drug resistance cassettes and probe HDAC2 are indicated by solid boxes, white boxes and a gray box, respectively. Relevant fragments obtained from BamHI/EcoRV digestions are shown. B) Southern blotting of homologous recombination event. The BamHI/EcoRV fragments of genomic DNAs prepared from DT40, one HDAC2(-/+) mutant clone cl.2 and six HDAC2(-/-) mutant clones (cl.2-1, cl.2-2, cl.2-3, cl.2-4, cl.2-5 and cl.2-6) were analyzed with probe HDAC2. C) Western blotting. Western blotting, using anti-chicken IgM L-chain and H-chain antibodies, was performed on total cellular proteins prepared from the six HDAC2(-/-) mutant clones at the early (E), middle (M) and later (L) cultivation stages and from DT40 cells (W). IgM H and L indicate IgM H- and L-chains (two bands), respectively. D) RT-PCR. RT-PCR was performed on total RNAs extracted from the six HDAC2(-/-) mutant clones at the three cultivation stages and from DT40 cells (W), using appropriate primers for whole (IgM Hc), secreted (IgM Hs) and membrane-bound (IgM Hm) forms of IgM H-chain mRNA and IgM L-chain mRNA (IgM L). The figure is identical with Fig. 8 of Refs. 198 and 223. The figure is also identical with Fig. 8 of Refs. 193, 197, 199, 222 plus 225 and Fig. 1 of Ref. 194 and Fig. 1 of Ref. 190. These figures were first presented in Ref. 138.

Figure 9. Alterations in amounts of IgM H- and L-chains in individual clones of HDAC2(-/-) DT40 mutants during continuous cultivation.

Western blotting was performed on total cellular proteins prepared from the six HDAC2(-/-) mutant clones (cl.2-1, cl.2-2, cl.2-3, cl.2-4, cl.2-5 and cl.2-6) at the indicated cultivation periods, including the early (E), middle (M) and later (L) stages and from DT40 cells (W). Proteins were sequentially detected with chicken antibodies for IgM L-chain that cross-reacts with IgM H-chain (top and third), IgM H-chain (second), and β -actin as a control. Upper and lower bands (indicated by IgM H) correspond to IgM H-chain detected by antibodies for IgM H- and L-chains, respectively. IgM L-chain (indicated by IgM L) was detected as two bands. The figure is identical with Fig. 9 of Refs. 198 and 223. The figure is also identical with Fig. 9 of Refs. 193, 197, 199, 222 plus 225 and Fig. 2 of Ref. 196 (and Fig. 2 of Ref. 194 and Fig. 2 of Ref. 190, which were slightly modified). These figures were first presented in Ref. 138.

Figure 10. Alterations in gene expressions of IgM H- and L-chains, and specific members of HATs, HDACs and transcription factors in individual clones of HDAC2(-/-) DT40 mutants during

continuous cultivation.

RT-PCR was performed on total RNAs prepared from four HDAC2(-/-) mutant clones (cl.2-1, cl.2-2, cl.2-4 and cl.2-6) at the indicated periods of cultivation, including the early (E), middle (M) and later (L) stages and from DT40 cells (W), using appropriate primers for IgM H- and L-chains, PCAF, HDAC9, Pax5, Aiolos, EBF1, Blimp1, OBF1, HDAC7, Ikaros, E2A, PU.1 and XBP-1, gene expressions of which changed in Ref. 96. The figure is identical with Fig. 10 of Refs. 198 and 223. The figure is also identical with Fig. 10 of Refs. 193, 197, 199, 222 plus 225 and Fig. 3 of Ref. 196 (and Fig. 4 of Ref. 194 and Fig. 4 of Ref. 190, which were slightly modified). These figures were first presented in Ref. 138.

Figure 11. Alterations in amounts of IgM H-chain in individual clones of HDAC2(-/-) DT40 mutants during continuous cultivation.

Electron microscopy (upper panels) and immuno-electron microscopy using anti-chicken IgM H-chain antiserum (lower panels) were performed on four HDAC2(-/-) mutant clones (cl.2-1, cl.2-2, cl.2-4 and cl.2-6) at the early (E) and later (L) cultivation stages and on DT40 cells (W). Dense cytoplasmic fractions due to accumulated IgM H-chain were observed only at the early (E) stage in the four mutant clones (E in upper panels). Positive signals for IgM H-chains were observed only at the early (E) stage in these mutant clones (E in lower panels). The figure is identical with Fig. 11 of Refs. 198 and 223. The figure is also identical with Fig. 11 of Refs. 193, 197, 199, 222 plus 225 and Fig. 4 of Ref. 190. These figures were first presented in Ref. 138.

Figure 12. Alterations of morphology of individual clones of HDAC2(-/-) DT40 mutants during continuous cultivation.

Microscopy was performed on four HDAC2(-/-) mutant clones (cl.2-1, cl.2-2, cl.2-4 and cl.2-6) at the early (E) and later (L) stages of cultivation and on DT40 cells (W) in several different ranges of vision. The different ranges of vision within microscopy are presented. The aggregative form was observed only at the early (E) stage in these four mutant clones. The figure is identical with Fig. 12 of Refs. 198 and 223. The figure is also identical with Fig. 12 of Refs. 193, 197, 199, 222 plus 225 and Fig. 5 of Ref. 190. These figures were first presented in Ref. 138.

Figure 13. Alterations in acetylation levels of specific Lys residues (K) of histone H3 within proximal 5'-upstream chromatin regions of Pax5, Aiolos, EBF1 and OBF1 genes in clone cl.2-1 of HDAC2(-/-) DT40 mutants during continuous cultivation.

Neighboring overlapping tiling chromatin immuno-precipitation (NotchIP) assay was carried out on the chromatin surrounding proximal 5'-upstream, distal 5'-upstream and ORF regions of Pax5, Aiolos, EBF1 and OBF1 genes. Cross-linked chromatins were prepared from clone cl.2-1 of HDAC2(-/-)

mutants at the early (E), middle (M) and later (L) cultivation stages and from DT40 (W), and co-precipitated by antisera specific for acetylated K9, K14, K18, K23 and K27 residues of histone H3. After de-crosslinking, co-precipitated chromatin was amplified by PCR using appropriate primers for the indicated segments of each of Pax5, Aiolos, EBF1 and OBF1 genes. β -actin was used as a control. A portion of cell supernatants as an input was amplified once by PCR using the same primers, and PCR products were analyzed by agarose gel electrophoresis. The figure is identical with Fig. 13 of Refs. 198 and 223. The figure is also identical with Fig. 13 of Refs. 193, 197, 199, 222 plus 225 and Fig. 4 of Ref. 196 and is a set of Figs. 1, 5, 9 and 13 of Ref. 191. These figures were first presented in Ref. 138.

Figure 14. Alterations in acetylation levels of specific Lys residues (K) of histone H3 within proximal 5'-upstream chromatin regions of Pax5, Aiolos, EBF1 and OBF1 genes in clone cl.2-2 of HDAC2(-/-) DT40 mutants during continuous cultivation.

NotchIP assay was carried out on the chromatin surrounding proximal 5'-upstream, distal 5'-upstream and ORF regions of Pax5, Aiolos, EBF1 and OBF1 genes in clone cl.2-2 of HDAC2(-/-) mutants at the early (E), middle (M) and later (L) cultivation stages and in DT40 (W) as described in the legend for Fig. 13. The figure is identical with Fig. 14 of Refs. 198 and 223. The figure is also identical with Fig. 14 of Refs. 193, 197, 199, 222 plus 225 and Fig. 5 of Ref. 196 and is a set of Figs. 2, 6, 10 and 14 of Ref. 191. These figures were first presented in Ref. 138.

Figure 15. Alterations in acetylation levels of specific Lys residues (K) of histone H3 within proximal 5'-upstream chromatin regions of Pax5, Aiolos, EBF1 and OBF1 genes in clone cl.2-4 of HDAC2(-/-) DT40 mutants during continuous cultivation.

NotchIP assay was carried out on the chromatin surrounding proximal 5'-upstream, distal 5'-upstream and ORF regions of Pax5, Aiolos, EBF1 and OBF1 genes in clone cl.2-4 of HDAC2(-/-) mutants at the early (E), middle (M) and later (L) cultivation stages and in DT40 (W) as described in the legend for Fig. 13. The figure is identical with Fig. 15 of Refs. 198 and 223. The figure is also identical with Fig. 15 of Refs. 193, 197, 199, 222 plus 225 and Fig. 6 of Ref. 196 and is a set of Figs. 3, 7, 11 and 15 of Ref. 191. These figures were first presented in Ref. 138.

Figure 16. Alterations in acetylation levels of specific Lys residues (K) of histone H3 within proximal 5'-upstream chromatin regions of Pax5, Aiolos, EBF1 and OBF1 genes in clone cl.2-6 of HDAC2(-/-) DT40 mutants during continuous cultivation.

NotchIP assay was carried out on the chromatin surrounding proximal 5'-upstream, distal 5'-upstream and ORF regions of Pax5, Aiolos, EBF1 and OBF1 genes in clone cl.2-6 of HDAC2(-/-) mutants at the early (E), middle (M) and later (L) cultivation stages and in DT40 (W) as described in the legend for Fig.

13. The figure is identical with Fig. 16 of Refs. 198 and 223. The figure is also identical with Fig. 16 of Refs. 193, 197, 199, 222 plus 225 and Fig. 7 of Ref. 196 and is a set of Figs. 4, 8, 12 and 16 of Ref. 191. These figures were first presented in Ref. 138.

Figure 17. Summary of alterations in acetylation levels of specific Lys residues (K) of histone H3, chromatin structure surrounding proximal 5'-upstream regions of Pax5, Aiolos, EBF1 and OBF1 genes, and their gene expression levels in DT40 cells (W) and clones cl.2-1, cl.2-2, cl.2-4 and cl.2-6 of HDAC2(-/-) DT40 mutants at early (E), middle (M) and later (L) stages of continuous cultivation.

Alterations in acetylation levels (hyper or hypo) of specific Lys residues (K) of histone H3, the chromatin structure (loose or tight form) of proximal 5'-upstream regions of Pax5, Aiolos, EBF1 and OBF1 genes, and their mRNA (gene expression) levels (high or low) in DT40 cells (W) and four clones (cl.2-1, cl.2-2, cl.2-4 and cl.2-6) of HDAC2(-/-) mutants at the early (E), middle (M) and later (L) stages of cultivation are schematically shown. The figure is identical with Fig. 17 of Refs. 198 and 223. The figure is also identical with Fig. 17 of Refs. 193, 197, 199, 222 plus 225 and is the slightly modified version of Fig.5 of Ref. 195. These figures were first presented in Ref. 138.

Figure 18. Localization of IgM H-chain at peri-nuclear space, endoplasmic reticulum and cell surface of HDAC2(-/-) DT40 mutant cells.

Immuno-electron microscopy using anti-chicken IgM H-chain antiserum was carried out on HDAC2(-/-) mutant cells at the early stage of cultivation. A) Immuno-electron microscopy without saponin-treatment. B) ~ E) Immuno-electron microscopy with saponin-treatment. Arrows P, E and S indicate positive signals of IgM H-chain proteins localized at the peri-nuclear space, endoplasmic reticulum and cell surface, respectively. Large amounts of accumulated IgM H-chain were observed in the peri-nuclear space of all HDAC2(-/-) mutant cells at the early cultivation stage. The figure is identical with Fig. 18 of Refs. 198 and 223. The figure is also identical with Fig. 18 of Refs. 193, 197, 199, 222 plus 225 and Fig. 8 of Ref. 196. These figures were first presented in Ref. 138.

Figure 19. Localization of IgM H-chain in peri-nuclear space of HDAC2(-/-) DT40 mutants at early and later stages of continuous cultivation.

Immuno-electron microscopy using anti-chicken IgM H-chain antiserum was carried out on HDAC2(-/-) mutant cells at the early (E) and later (L) cultivation stages. A) Immuno-electron microscopy of one HDAC2(-/-) mutant cell at the early (E) stage. B) and C) Enlarged versions of parts indicated by arrows b and c in A). D), G) and J) Immuno-electron microscopy of three individual HDAC2(-/-) mutant cells at the later (L) stage. E), F), H), I) and K) Enlarged versions of parts indicated by arrows e, f, h, i and k in D), G) and J). The mutant cell in G) was the same one of Ref. 95. Large

amounts of accumulated IgM H-chain were observed in the peri-nuclear space of all HDAC2(-/-) mutant cells at the early (E) and later (L) stages. The figure is identical with Fig. 19 of Refs. 198 and 223. The figure is also identical with Fig. 19 of Refs. 193, 197, 199, 222 plus 225 and Fig. 3 of Ref. 192. These figures were first presented in Ref. 138.

Figure 20. Localization of IgM H-chain at peri-nuclear space, and a model for signal transduction concerning accumulated IgM H- and L-chains to chromatin in HDAC2(-/-) DT40 mutants during continuous cultivation.

Left panel: A portion of the peri-nuclear space (where IgM H- and L-chains were accumulated) of the HDAC2(-/-) mutant cell (indicated by an arrow P at lower position in Fig. 18C) was reversely enlarged. **Right panel:** A model for the signal transduction on the accumulation of IgM H- and L-chains. Signal concerning IgM H- and L-chains artificially accumulated in the peri-nuclear space of HDAC2(-/-) mutants was repeatedly transmitted to the chromatin structure, followed by unbalanced response for the signal and its convergence to various specific genes (such as Pax5, Aiolos, EBF1, OBF1, etc.) in individual mutant clones. ECRR/ECRS: environment change recognition receptor/site. 4C machinery: chromatin conformation change complex (4C) machinery. The figure is identical with Fig. 20 of Refs. 198 and 223. The figure is also identical with Fig. 20 of Refs. 193, 197, 199, 222 plus 225 and Fig. 9 of Ref. 196. These figures were first presented in Ref. 138.

Figure 21. Summary of alterations in gene expression levels, acetylation levels of specific Lys residues (K) of histone H3, and chromatin structure surrounding proximal 5'-upstream regions of Pax5, Aiolos, EBF1 and OBF1 genes in individual clones of HDAC2(-/-) DT40 mutants during continuous cultivation.

Alterations in mRNA (gene expression) levels (-, ++ or ++++), acetylation levels (Ac; -, ++ or ++++) of specific Lys residues (K) of histone H3 and the chromatin structure (loose or tight form) surrounding proximal 5'-upstream regions of Pax5, Aiolos, EBF1 and OBF1 genes in four HDAC2(-/-) mutant clones (cl.2-1, cl.2-2, cl.2-4 and cl.2-6) at the early (E) and later (L) stages of cultivation and in DT40 cells (W) are schematically presented. Alterations in mRNA levels (- or ++++) of IgM H- and L-chains are also presented. Chromatin conformation change complex (4C) machinery generally contains a specific member of each of HATs plus HDACs and other factors. Transcription factor complex (TFC) machinery generally contains RNA polymerase (RPase), specific transcription factors and others. The figure is identical with Fig. 21 of Refs. 198 and 223. The figure is also identical with Fig. 21 of Refs. 193, 197, 199, 222 plus 225 and Fig. 10 of Ref. 196. These figures were first presented in Ref. 138.

Figure 22. Irreversible creation of chromatin structure plasticity surrounding proximal

5'-upstream region(s) of specific gene(s) with epigenetic modifications in descendent cells, and reversible regulations of ordinary gene expression and enzyme reaction in cells that initially accept proper signal.

Upper panel: Irreversible creation of chromatin structure plasticity surrounding proximal 5'-upstream region(s) of specific gene(s) with epigenetic modifications occurs in descendant cells but not in initial cell that accepts environment change signal. Ac, Ac/2 and Ac/10 indicate qualitatively hyper-, considerably hyper- and somewhat hyper-acetylation levels of specific Lys residues (K) of core histones (e.g., histone H3), respectively. **Middle and Lower panels:** Regulations of ordinary gene expression (at transcriptional regulatory elements) and enzyme reaction reversibly occur in the cell that accepts proper signal. Ac and P indicate acetylation, phosphorylation, ADP-ribosylation and/or other chemical modifications, respectively. The figure is identical with Fig. 22 of Refs. 198 and 223. The figure is also identical with Fig. 22 of Refs. 193, 197, 199, 222 plus 225 and Fig. 12 of Ref. 196. These figures were first presented in Ref. 138.

Figure 23. Chromatin conformation change code (4C) theory: A model for gaining un-programmed and new cell function(s) by means of irreversible creation of chromatin structure plasticity surrounding proximal 5'-upstream region(s) of specific gene(s) with epigenetic modifications through various generations.

Irreversible creation of chromatin structure plasticity with epigenetic modifications occurs surrounding proximal 5'-upstream region(s) (notch of chromatin) but not ORF region(s) of specific gene(s) through various generations (cell divisions). The tight or loose form of the chromatin structure is based on hypo- or hyper-acetylation levels of specific Lys residues (K) of histone H3, and causes low or high transcription (gene expression) levels. The figure is identical with Fig. 23 of Refs. 198 and 223. The figure is also identical with Fig. 23 of Refs. 193, 197, 199, 222 plus 225 and Fig. 11 of Ref. 196. These figures were first presented in Ref. 138.