

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: [www.elsevier.com/locate/radcr](http://www.elsevier.com/locate/radcr)

## Case Report

# Hemorrhagic shock due to ruptured idiopathic intramural hematoma of the sigmoid colon—An autopsy case report<sup>☆</sup>

Norihiro Shinkawa, MD, PhD<sup>a,b,\*</sup>, Eiji Kakizaki, PhD<sup>a</sup>, Ai Sonoda, MSc<sup>a</sup>,  
Nobuhiro Yukawa, MD, PhD<sup>a</sup>

<sup>a</sup> Section of Legal Medicine, Department of Social Medicine, Faculty of Medicine, University of Miyazaki, Kiyotake-cho Kihara 5200, Miyazaki 889-1692, Japan

<sup>b</sup> Radiology Division, Faculty of Medicine, University of Miyazaki Hospital, Miyazaki, Japan

## ARTICLE INFO

## Article history:

Received 19 December 2022

Accepted 27 December 2022

## Keywords:

Intramural hematoma

Colon

Idiopathic

Forensic pathology

## ABSTRACT

We describe herein an autopsy case involving fatal hemorrhagic shock due to idiopathic sigmoid colonic intramural hematoma rupturing into the abdominal cavity. Antemortem computed tomography revealed a mass lesion in the sigmoid colon and a large amount of hemoperitoneum. On forensic autopsy, intramural hematoma of the sigmoid colon with ruptured serosa was identified, while the mucosa remained intact. Microscopically, hematomas were apparent, mainly in the muscularis propria. We diagnosed the cause of death as hemorrhagic shock due to idiopathic intramural hematoma of the sigmoid colon rupturing into the abdominal cavity. Although the patient had been receiving peritoneal dialysis, no relationship was identified between dialysis catheters and sigmoid colon intramural hematoma rupture. On computed tomography, the mass lesion was initially considered a submucosal neoplastic lesion or endometriotic lesion. Intramural hematoma should be considered as a differential diagnosis.

© 2022 The Authors. Published by Elsevier Inc. on behalf of University of Washington.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

## Introduction

Intramural hematoma of the colon is rare, with more than 90% of gastrointestinal intramural hematomas occurring in the duodenum and only 5% involving the colon [1,2]. For example, esophageal [3–5], gastric [6], and duodenal [7] intramu-

ral hematomas have been reported in the *Radiology Case Reports*. Gastrointestinal intramural hematomas mainly develop in the submucosa or muscularis propria [6]. The most common cause of intramural hematoma of the colon is blunt abdominal trauma, followed by anticoagulant therapy and blood dyscrasia [8]. We present herein the case of a woman in whom idiopathic intramural hematoma of the sigmoid colon, mainly

<sup>☆</sup> Competing Interests: None.

\* Corresponding author.

E-mail address: [norihiro\\_shinkawa@med.miyazaki-u.ac.jp](mailto:norihiro_shinkawa@med.miyazaki-u.ac.jp) (N. Shinkawa).

<https://doi.org/10.1016/j.radcr.2022.12.061>

1930-0433/© 2022 The Authors. Published by Elsevier Inc. on behalf of University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

between the inner circular and outer longitudinal muscle layers, ruptured into the abdominal cavity, causing death by hemorrhagic shock. As the patient had been receiving continuous ambulatory peritoneal dialysis (CAPD), from a forensic perspective, we had to elucidate the relationship between dialysis catheters and intramural hematoma of the sigmoid colon.

## Case report

### Case history

The patient was a premenopausal woman in her 50s who had been diagnosed with diabetes 8 years earlier. She had been receiving CAPD for diabetic nephropathy at home for about a year. In addition to diabetic nephropathy, the primary-care physician had noted hypertension and hyperlipidemia, but no abnormalities suggestive of hematological malignancies or coagulation/fibrinolytic disorders were obvious from blood test results.

The night before this event, according to dialysis records, the patient started peritoneal dialysis as usual. The next morning, she was found breathing but unconscious. At that time, drainage fluid from the peritoneal dialysis showed reddish discoloration and the dialyzer had issued an alarm and displayed the alert “poor drainage.” She was taken to the emergency room and given mechanical chest compressions, endotracheal intubation, and adrenaline injections, but died about 3 hours later.

### Findings from antemortem computed tomography

Antemortem non-contrast-enhanced computed tomography (CT) was performed in the emergency room (Fig. 1). A large amount of hemoperitoneum was evident. In addition, a mass lesion 3 cm in diameter was identified in the sigmoid colon (Figs. 1A, B, and D). The mass lesion showed homogeneous high attenuation compared to muscle tissue and a smooth margin (Figs. 1G and H). These findings for the mass lesion were considered inconsistent with cancer, but whether the lesion represented a submucosal neoplasm such as gastrointestinal stromal tumor or lymphoma or a non-neoplastic lesion such as intestinal endometriosis could not be determined.

A dialysis catheter was observed to the left of the mass lesion, separated by a 0.8-cm thickness of adipose tissue (Figs. 1A and B). Soft tissue displaying heterogeneous attenuation was seen on the right side of the mass lesion, which appeared to be a soft blood clot or fibrin (Figs. 1A, B, and D). Rupture and hemorrhage on the right side of the mass lesion were considered because soft blood clots and fibrin were unevenly distributed on the right side of the mass lesion. The tip of the dialysis catheter was located deep in the pouch of Douglas (Fig. 1F). No soft clot was seen at the site where the dialysis catheter penetrated the abdominal wall (Figs. 1C and E). Encapsulating peritoneal sclerosis is a known complication of peritoneal dialysis, but was ruled out in this case.

Although hemorrhage from the sigmoid colon was strongly suspected, as described above, the patient was taken to the

emergency room in a state of shock, and was unable to undergo other examinations, including contrast-enhanced CT and colonoscopy.

### Autopsy findings


Autopsy was performed 3 days postmortem. External examination showed that the patient was 155 cm tall and weighed 73 kg. Livor mortis was weak, and the conjunctivae and oral mucosa were pale. No abnormalities were evident at the insertion site of the dialysis catheter. No injuries other than those caused by the mechanical compressions for cardiopulmonary resuscitation were observed. No evidence of blunt trauma to the abdomen was seen.

Internal examination showed 4290 mL of bloody ascites, including soft blood clots and fibrin (Figs. 2A and B). However, since this volume also contained peritoneal dialysate, the measured amount of bleeding was considered inaccurate. A dark-red mass lesion 3 cm in diameter was identified in the sigmoid colon. The serosa over the lesion was ruptured and was considered the origin of bloody ascites (Fig. 3). The site of mass lesion rupture was far from the catheter. Further, the tip of the dialysis catheter was located deep in the pouch of Douglas. The cut surface of the sigmoid colon mass showed hematoma, and no lesions such as neoplasms were detected macroscopically. The mucosa overlying the lesion remained intact. Hemoglobin concentration was 3.1 g/dL and hematocrit was 11.4%. The patient displayed renal anemia antemortem, with a hemoglobin concentration of 7.6 g/dL and hematocrit of 20.8% at 3 weeks before her death. Blood volumes in various organs were low, and slight hemorrhage was observed in the left ventricular endocardium. These findings were considered to reflect lethal hemorrhage.

In addition, renal atrophy was observed and was considered consistent with diabetic nephropathy. Except for ruptured intramural hematoma of the sigmoid colon and bloody ascites, no lesions considered capable of causing death were obvious. Poisoning was also ruled out. CT and microscopy showed no findings suggestive of dialysis amyloidosis in the joints, spine, or other organs.

Microscopically, the mucosa of the sigmoid colonic lesion remained intact. Hematomas were localized from the submucosa to the subserosa, mainly between the inner circular and outer longitudinal muscle layers (Fig. 4). No pathology responsible for bleeding, such as neoplasm, amyloidosis, or angiodysplasia, was identified. Infiltration of inflammatory cells and fibrin (Figs. 2C and D) were confirmed, and the hematoma was considered fresh. No chronic inflammation suggestive of chronic catheter-related stimulation was found in the serosa of the sigmoid colon. Other microscopic findings included hyaline-like changes in efferent and afferent arterioles and fibrin caps in the renal glomeruli. Disrupted glomeruli were also seen. These findings were considered consistent with diabetic nephropathy.

From the above macro- and microscopic findings, we diagnosed the cause of death as hemorrhagic shock due to intramural hematoma of the sigmoid colon that had ruptured into the abdominal cavity. As for the cause of the intramural hematoma, no blunt trauma to the abdomen or underlying disease state was identified, so the case had to be considered



**Fig. 1** – Antemortem non-contrast-enhanced CT images. Circles indicate the mass lesion in the sigmoid colon. Arrowheads indicate the dialysis catheter. Arrows indicate adipose tissue interposed between the mass lesion and dialysis catheter. Dashed circles indicate soft blood clots and fibrin. (A) Coronal-view reconstruction of abdominal computed tomography (CT). (B) Axial-view image from pelvic CT. (C) Axial image taken 4.5 cm cranial from Image B. (D) Sagittal-view reconstruction of abdominal CT. (E) Sagittal-view image taken 2.1 cm to the left of Image D. (F) Sagittal-view image taken 1.2 cm to the left of Image E. (G) Enlargement of Image A. (H) Enlargement of Image B.

idiopathic. No other lesions that could have caused death were evident.


## Discussion

Intramural hematoma of the colon is rare [1], and clinical case reports have been described only sporadically. We were able to identify 28 clinical case reports in searches of PubMed and Scopus using the keywords “intramural hematoma” and “colon.” Among these 28 cases, 1 patient died. One case involved postpolypectomy complications in which hemicolectomy was performed to include an intramural hematoma le-


sion, but the patient died 4 days after surgery [9]. In addition to these searches in the literature, 1 fatal case was reported in Japanese. The patient had a history of myelodysplastic syndrome and chronic heart failure, and was warfarinized. The lesion was resected after the patient developed an intramural hematoma of the colon, but he died 34 days after surgery due to heart and respiratory failure [10].

### Etiology of intramural hematoma of the colon


The most common cause of intramural hematoma of the colon is blunt abdominal trauma, followed by anticoagulant therapy and blood dyscrasia [8]. In this case, the patient had received mechanical chest compressions but sigmoid colon



**Fig. 2** – Bloody ascites, soft blood clots, and fibrin. (A) Bloody ascites (4290 mL). (B) Soft blood clots and fibrin. (C) Microscopic views of fibrin. Hematoxylin and eosin (HE),  $\times 40$ . (D) Microscopic views of fibrin. Phosphotungstic acid hematoxylin,  $\times 40$ .



**Fig. 3** – Arrows indicate the site of rupture of the sigmoid serosa. (A) Abdominal cavity. Most of the bloody ascites has been removed. (B) Most of the soft clot and fibrin removed from photo A. (C) Whole colon removed. (D) Close-up image of the rupture site in the sigmoid serosa.



**Fig. 4 – Microscopic views of intramural hematoma from the sigmoid colon. HE. (A) The mucosa is intact, with dense hematoma in the lower layer of the inner circular muscle seen in the lower right of the photograph. Slight bleeding is also observed in the submucosa.  $\times 40$ . (B) Lower layer of Image A. Below the photograph, the serosa is ruptured.  $\times 40$ . (C) An area in which hematoma is clearly visible between the inner circular and outer longitudinal muscle layers. Hematoma is also evident in the subserosa. The serosa is not ruptured in this area.  $\times 40$ . (D) Enlargement of Image C. Inflammatory cells and fibrin are apparent.  $\times 100$ .**

injury was unlikely to have resulted as a complication of these [11]. Gastrointestinal hematomas are estimated to occur in 1 of every 250,000 anticoagulated patients [12]. The patient in our case did not show findings suggestive of blunt abdominal trauma, was not anticoagulated and had no obvious blood dyscrasias. Postpolypectomy status [9,13], endometriosis [14], and a background of acute cholecystitis [15] or intestinal pneumatosis [16] have been reported as other causes. In our case, those causes were all excluded by a comprehensive medical history and macro- and microscopic findings. We therefore had to classify this case as idiopathic, a cause representing about 1% of intramural hematomas [17].

#### Symptoms of intramural hematoma of the colon

The clinical symptoms associated with intramural hematoma of the colon are nonspecific, varying from tolerable abdominal pain to hemorrhagic shock [9]. Symptoms may differ depending on whether the bleeding remains intramural, ruptures and bleeds into the mucosa, or ruptures and bleeds outside the serosa. Cases of bleeding remaining intramural [8,18], cases of rupture and bleeding into the lumen resulting in hematochezia [19,20], and cases of rupture and bleeding into the abdominal cavity [21,22] have all been reported. Cases of intramural hematoma of the ascending colon bleeding in the

retroperitoneum [23] and massive bleeding from intramural hematoma of the ascending colon extending to the intestinal lumen, abdominal cavity, mesentery, and retroperitoneum have also been reported [11].

#### Diagnosis and treatment of intramural hematoma of the colon

Nakamura et al. reported a case of idiopathic gastric intramural hematoma diagnosed using contrast-enhanced ultrasonography [6]. Because our patient was obese, sigmoid intramural hematoma may have been difficult to diagnose on ultrasonography.

Colonoscopy is useful for diagnosis [8]. Colonoscopy shows an elevated lesion covered with normal colonic mucosa and blue mucosa around the lesion, reflecting submucosal hematoma [17].

Hematoma is visualized as a high-attenuation mass compared to muscle tissue on CT in the acute phase, but the hematoma is gradually absorbed and attenuation thus decreases over time, requiring differentiation from other lesion such as cysts, abscesses, and necrotic tumors [21]. In our case, the mass showing high attenuation compared to muscle tissue detected on CT (Fig. 1) was considered consistent with acute hematoma. The patient also underwent dialysis 3

times a day, and drainage fluid from the previous dialysis was normal according to dialysis records. This time course and histopathological findings were also consistent with acute hematoma. We considered that contrast-enhanced CT contributes to the qualitative diagnosis of mass lesions and identification of bleeding sites.

Magnetic resonance imaging (MRI) findings of intramural hematoma have been reported [17,24]. On MRI, signal changes vary depending on the postbleeding interval [17,24]. The patient in this case was taken to the emergency room in a state of shock, with no time to perform MRI.

Intramural hematomas of the colon are usually found in the submucosal layer [6,10,11,17,25] and muscularis propria [6]. A case of idiopathic bleeding in the inner circular muscle layer has also been reported [21]. In this case, hematoma was mainly localized between the inner circular and outer longitudinal muscle layers. The cause of bleeding into the muscularis propria is unknown.

Surgical treatment is considered the gold standard by some [8,19], while conservative treatment remains the first choice according to others [10,21]. Of course, surgical treatment carries risks in patients on anticoagulant therapy or with underlying blood disorders [10].

#### **Relationship between peritoneal dialysis and bleeding in this case**

Takagi et al. [26] reviewed 45 patients with peritonitis due to gastrointestinal perforation in patients with CAPD. The most common site of perforation was the sigmoid colon. Catheters were a rare cause of perforation, involving 3 cases. Takagi et al. [26] considered that the main cause of perforation was the addition of chronic catheter-related stimulation of the intestinal tract to underlying conditions of amyloidosis and diverticulitis. In our case, no catheter-related chronic inflammation, amyloidosis or diverticulitis was observed. The dialysis catheter and hematoma were clearly separate in this patient, with no adhesions between the catheter and surrounding tissue. The tip of the catheter was thus considered unlikely to have injured the sigmoid colon in this episode. In that sense, we ruled out involvement of the catheter.

Reddish drainage fluid from peritoneal dialysis is often related to menses or ovulation, particularly ovarian cyst rupture [27]. Although the reddish drainage fluid in the present case was due to intramural hematoma of the sigmoid colon fatally rupturing into the abdominal cavity, the patient had previously experienced reddish drainage fluid that had been attributed to menstrual issues.

Amyloidosis is a complication of dialysis and must therefore be considered. The prevalence of amyloidosis is similar in hemodialysis and peritoneal dialysis patients [28]. A case of gastric and colonic submucosal hematoma with amyloidosis has been reported, although that patient had not undergone dialysis [29]. Therefore, in the present case, we should consider the possibility that gastrointestinal amyloidosis occurred as a complication of dialysis and intramural hematoma occurred in the gastrointestinal tract. Although amyloidosis occurs in patients with end-stage renal failure on long-term CAPD [28], the patient in our case had a relatively short dialysis history of only 1 year, and amyloidosis due to dialysis

would thus have been unlikely. In addition, amyloid deposition as seen in the above case report [29] was not observed microscopically in our case.

In conclusion, we have presented a rare case of ruptured idiopathic intramural hematoma of the sigmoid colon. Nonenhanced CT alone was able to identify the site of bleeding and matched well with findings from autopsy and pathology. With these findings on CT, intramural hematoma should be considered among the differential diagnoses.

#### **Patient consent**

Written consent was obtained from the patient's close relative for publication of the case information.

#### **REFERENCES**

- [1] Vecchio R, Cacciola E, Figuera M, Catalano R, Giulla G, Distefano ER, et al. Idiopathic intramural hematoma of the right colon. A case report and review of the literature. *Int J Surg Case Rep* 2019;60:16–20. doi:10.1016/j.ijscr.2019.05.004.
- [2] Hughes CE III, Conn J Jr, Sherman JO. Intramural hematoma of the gastrointestinal tract. *Am J Surg* 1977;133(3):276–9. doi:10.1016/0002-9610(77)90528-1.
- [3] Stooksberry T, McHam B, Lisle M. Undiagnosed aorto-esophageal fistula causing intramural hematoma of the esophagus. *Radiol Case Rep* 2020;15(8):1128–32. doi:10.1016/j.radcr.2020.05.040.
- [4] Sharma K, Wang Y. Submucosal esophageal hematoma precipitated by chronic idiopathic thrombocytopenic purpura. *Radiol Case Rep* 2017;12(2):278–80. doi:10.1016/j.radcr.2017.01.022.
- [5] Isaac MFG, Ho CL, Leong S. Life-threatening bleeding from dissecting intramural hematoma of esophagus (IHE) treated by trans arterial embolization. *Radiol Case Rep* 2021;16(9):2474–7. doi:10.1016/j.radcr.2021.05.067.
- [6] Nakamura J, Manabe N, Kato K, Suehiro M, Ayaki M, Fujita M, et al. A case of idiopathic gastric intramural hematoma diagnosed using contrast-enhanced ultrasonography and followed up with conservative treatment. *Radiol Case Rep* 2022;17(12):4487–91. doi:10.1016/j.radcr.2022.08.105.
- [7] Amankwa AT, De Graft-Johnson E, Twum KA, Ankomah K. A case of spontaneous intramural duodenal hematoma in a young African man: imaging findings. *Radiol Case Rep* 2021;16(7):1675–8. doi:10.1016/j.radcr.2021.03.070.
- [8] Santis G, Sista F, Giuliani A, Leardi S. Idiopathic intramural hematoma of sigmoid colon. A case report. *Ann Ital Chir* 2011;82:395–7.
- [9] Voudoukis E, Spiridakis KG, Karmiris K, Georgiou G, Kontopodis N, Konstantinidis K, et al. Intramural hematoma of the ascending colon leading to intraperitoneal hemorrhage: a possible post-polypectomy complication. *Ann Gastroenterol* 2012;25(3):265–7.
- [10] Kamimura K, Hagio K, Isono T, Ueda T, Ashizawa N, Nakada H, et al. A case of intramural hematoma of the ascending colon in a patient with myelodysplastic syndrome and anticoagulant therapy. *Nihon Rinsho Geka Gakkai Zasshi (Jpn Surg Assoc)* 2014;75(10):2789–94 [in Japanese].
- [11] Milling L, Mikkelsen S, Astrup BS. Characteristics of mechanical CPR-related injuries: a case series. *J Forensic Leg Med* 2020;70:101918. doi:10.1016/j.jflm.2020.101918.

- [12] Abdel Samie A, Theilmann L. Risk factors and management of anticoagulant-induced intramural hematoma of the gastrointestinal tract. *Eur J Trauma Emerg Surg* 2013;39(2):191–4. doi:10.1007/s00068-013-0250-1.
- [13] Yu S, Wu G, Wen Z, Huang H. Intramural hematoma of the colon after polypectomy—an unusual case occurrence. *Endoscopy* 2014;46:E46. doi:10.1055/s-0033-1359122.
- [14] Wynn TE. Endometriosis of the sigmoid colon. Massive intramural hematoma. *Arch Pathol* 1971;92(1):24–7.
- [15] Hess RA. Intramural hematoma of the transverse colon with obstruction and possible impending gangrene: report of a case associated with acute hemorrhagic cholecystitis. *Am Surg* 1964;30:138–40.
- [16] Bonardi JM, Clerc G, Juif JG. Intramural hematoma of the large intestine and intestinal pneumatosis. *Pediatric* 1977;32:261–6.
- [17] Umeda I, Ohta H, Doi T, Nobuoka A, Kanisawa Y, Kawasaki R, et al. Idiopathic intramural hematoma of the colon. *Gastrointest Endosc* 2007;66:861–4. doi:10.1016/j.gie.2007.02.003.
- [18] Fernandes SR, Gonçalves AR, Correia LA. Spontaneous intramural hematoma of the colon. *Rev Esp Enferm Dig* 2016;108(8):501–2.
- [19] Figueiredo J, Borges N, Pascoalinho J, Matos R. Giant intramural hematoma of the colon in acquired factor viii inhibitor. *Acta Med Port* 2019;32(9):614–17. doi:10.20344/amp.8519.
- [20] Nagami Y, Fujiwara Y, Yamamura M, Shiba M, Watanabe T, Tominaga K, et al. Intramural hematoma of the colon caused by double-balloon enteroscopy in a patient with chronic disseminated intravascular coagulation. *Endoscopy* 2016;48:E105–6. doi:10.1055/s-0042-104276.
- [21] Imura J, Miyata K, Yuasa N, Takeuchi E, Goto Y, Miyake H, et al. A case of colonic perforation due to idiopathic intramural hematoma of the sigmoid colon. *Nihon Shokaki Geka Gakkai Zasshi (Jpn J Gastroenterol Surg)* 2012;45(5):552–8 [in Japanese]. doi:10.5833/jjgs.45.552.
- [22] Bakaeen FG, Chow M, Ghadir M, Albo D, Berger DH, Chu D. Ruptured colonic intramural hematoma with massive hemorrhage after aortic valve replacement. *Am Surg* 2010;76(6):647–8.
- [23] Tanaka H, Endoh Y, Nakagawa M, Kojima T. Intramural hematoma of the colon due to blunt trauma. *Nihon Kyukyū Igakukai Zasshi (J Jpn Assoc Acute Med) [in Japanese]* 1992;3:189–93.
- [24] Niehues SM, Denecke T, Bassir C, Hamm B, Haas M. Intramural duodenal hematoma: clinical course and imaging findings. *Acta Radiol Open* 2019;8(4). doi:10.1177/2058460119836256.
- [25] Lee SH, Lee JH, Park DH, Chung IK, Park SH, Kim SJ. Intramural colonic hematoma: complication of anticoagulation with heparin. *Gastrointest Endosc* 2005;62(5):783–4. doi:10.1016/j.gie.2005.04.045.
- [26] Takagi A, Hatta T, Ueno R, Kado H, Segawa H, Shiotsu Y, et al. A patient with perforative peritonitis undergoing peritoneal dialysis: a case report. *Nihon Toseki Igakkai Zasshi (J Jpn Soc Dial Ther)* 2011;44(10):1039–45 [in Japanese]. doi:10.4009/jsdt.44.1039.
- [27] Kuwahara T, Yoshinaga N, Hashimoto H, Tanaka A, Yahata K, Sugawara A, et al. Recurrent hemoperitoneum with different etiologies in a woman receiving continuous ambulatory peritoneal dialysis with end stage renal disease due to chronic rheumatoid arthritis and secondary. *Nihon Toseki Igakkai Zasshi (J Jpn Soc Dial Ther)* 1995;28(10):1383–7 [in Japanese]. doi:10.4009/jsdt.28.1383.
- [28] Tan SY, Baillod R, Brown E, Farrington K, Soper C, Percy M, et al. Clinical, radiological and serum amyloid P component scintigraphic features of  $\beta$ 2-microglobulin amyloidosis associated with continuous ambulatory peritoneal dialysis. *Nephrol Dial Transplant* 1999;14(6):1467–71. doi:10.1093/ndt/14.6.1467.
- [29] Sato T, Chinzei R, Funada K, Ohtsu I, Doi H, Takahashi M, et al. A case of gastric and colonic submucosal hematomas with amyloidosis. *Prog Dig Endosc* 2018;92(1):92–3. doi:10.11641/pde.92.1\_92.